

# Honours Analysis Notes

Anthony Catterwell

April 18, 2019

## Contents

### 1 The Real Number System

1.1	Introduction . . . . .	
1.2	Ordered Field Axioms . . . . .	
1.3	Completeness Axiom . . . . .	
1.4	Mathematical Induction . . . . .	
1.5	Inverse Functions and Images . . . . .	
1.6	Countable and Uncountable Sets . . . . .	

### 2 Sequences in $\mathbb{R}$

2.1	Limits of Sequences . . . . .	
2.2	Limit Theorems . . . . .	
2.3	Bolzano-Weierstrass Theorem . . . . .	
2.4	Cauchy Sequences . . . . .	
2.5	Limits Supremum and Infimum . . . . .	

# 1 The Real Number System

## 1.1 Introduction

## 1.2 Ordered Field Axioms

- **Postulate 1** *Field Axioms*

There are functions  $+$  and  $\cdot$  defined on  $\mathbb{R}^2 := \mathbb{R} \times \mathbb{R}$ , which satisfy the following properties  $\forall a, b, c \in \mathbb{R}$

- *Closure Properties*:  $a + b, a \cdot b \in \mathbb{R}$
- *Associative Properties*:  $a + (b + c) = (a + b) + c$  and  $a \cdot (b \cdot c) = (a \cdot b) \cdot c$
- *Commutative Properties*:  $a + b = b + a$  and  $a \cdot b = b \cdot a$
- *Distributive Law*:  $a \cdot (b + c) = a \cdot b + a \cdot c$
- *Existence of Additive Identity*: There is a unique element  $0 \in \mathbb{R}$  such that  $0 + a = a$  for all  $a \in \mathbb{R}$
- *Existence of Multiplicative Identity*: There is a unique element  $1 \in \mathbb{R}$  such that  $1 \neq 0$  and  $1 \cdot a = a$  for all  $a \in \mathbb{R}$
- *Existence of Additive Inverses*: For every  $x \in \mathbb{R}$  there is a unique element  $-x \in \mathbb{R}$  such that

$$x + (-x) = 0$$

- *Existence of Multiplicative Inverses*: For every  $x \in \mathbb{R} \setminus \{0\}$  there is a unique element  $x^{-1} \in \mathbb{R}$  such that

$$x \cdot (x^{-1}) = 1$$

- **Postulate 2** *Order Axioms*

There is a relation  $<$  on  $\mathbb{R} \times \mathbb{R}$  that has the following properties:

- *Trichotomy Property*: Given  $a, b \in \mathbb{R}$ , one and only one of the following statements hold:

$$a < b, b < a, \text{ or } a = b$$

- *Transitive Property*: For  $a, b, c \in \mathbb{R}$

$$a < b \text{ and } b < c \implies a < c$$

- *Additive Property*: For  $a, b, c \in \mathbb{R}$

$$a < b \text{ and } c \in \mathbb{R} \implies a + c < b + c$$

- *Multiplicative Properties*: For  $a, b, c \in \mathbb{R}$

$$a < b \text{ and } c > 0 \implies ac < bc$$

and

$$a < b \text{ and } c < 0 \implies bc < ac$$

- **Remark 1.1**

We will assume that the sets  $\mathbb{N}$  and  $\mathbb{Z}$  satisfy the following properties:

1. If  $n, m \in \mathbb{Z}$ , then  $n + m, n - m$  and  $mn$  belong to  $\mathbb{Z}$
2. If  $n \in \mathbb{Z}$ , then  $n \in \mathbb{N}$  if and only if  $n \geq 1$

3. There is no  $n \in \mathbb{Z}$  that satisfies  $0 < n < 1$

• **Definition 1.4** *Absolute Value*

The *absolute value* of a number  $a \in \mathbb{R}$  is the number

$$|a| := \begin{cases} a & a \geq 0 \\ -a & a < 0 \end{cases}$$

• **Remark 1.5** The *absolute value* is multiplicative; that is,  $|ab| = |a||b| \forall a, b \in \mathbb{R}$

• **Theorem 1.6** *Fundamental Theorem of Absolute Values*

Let  $a \in \mathbb{R}$  and  $M \geq 0$ . Then  $|a| \leq M \iff -M \leq a \leq M$ .

• **Theorem 1.7** The absolute value satisfies the following three properties:

1. *Positive Definite*: For all  $a \in \mathbb{R}$ ,  $|a| > 0$  with  $|a| = 0$  if and only if  $a = 0$ .
2. *Symmetric*: For all  $a, b \in \mathbb{R}$ ,  $|a - b| = |b - a|$ ,
3. *Triangle Inequalities*: For all  $a, b \in \mathbb{R}$ ,

$$|a + b| \leq |a| + |b| \quad \text{and} \quad ||a| - |b|| \leq |a - b|$$

• **Theorem 1.9** Let  $x, y, a \in \mathbb{R}$

1.  $x < y + \epsilon \forall \epsilon > 0 \iff x \leq y$
2.  $x > y - \epsilon \forall \epsilon > 0 \iff x \geq y$
3.  $|a| < \epsilon \forall \epsilon > 0 \iff a = 0$

### 1.3 Completeness Axiom

• **Definition 1.10** *Upper bounds*

Let  $E \subset \mathbb{R}$  be non-empty

1. The set  $E$  is said to be *bounded above* if and only if there is an  $M \in \mathbb{R}$  such that  $a \leq M$  for all  $a \in E$ , in which case  $M$  is called an *upper bound* of  $E$ .
2. A number  $s$  is called a *supremum* of the set  $E$  if and only if  $s$  is an upper bound of  $E$  and  $s \leq M$  for all upper bounds  $M$  of  $E$ . (In this case we shall say that  $E$  has a *finite supremum*  $s$  and write  $s = \sup E$ )

• **Remark 1.12** If a set has one upper bound, it has infinitely many upper bounds.

• **Remark 1.13** If a set has a supremum, then it has only one supremum.

• **Theorem** *Approximation Property for Suprema*

If  $E$  has a finite supremum and  $\epsilon > 0$  is any positive number, then there is a point  $a \in E$  such that

$$\sup E - \epsilon < a \leq \sup E$$

• **Theorem 1.15**

If  $E \subset \mathbb{Z}$  has a supremum, then  $\sup E \in E$ . In particular, if the supremum of a set, which contains only integers, exists, that supremum must be an integer.

• **Postulate 3** *Completeness Axiom*

If  $E$  is a nonempty subset of  $\mathbb{R}$  that is bounded above, then  $E$  has a finite supremum.

• **Theorem 1.16** *The Archimedean Principle*

Given real numbers  $a$  and  $b$ , with  $a > 0$ , there is an integer  $n \in \mathbb{N}$  such that  $b < na$ .

- **Theorem 1.18** *Density of Rationals*

If  $a, b \in \mathbb{R}$  satisfy  $a < b$ , then there is a  $q \in \mathbb{Q}$  such that  $a < q < b$ .

- **Definition 1.19** *Upper bounds*

Let  $E \subseteq \mathbb{R}$  be nonempty

1. The set  $E$  is said to be *bounded below* if and only if there is an  $m \in \mathbb{R}$  such that  $a \geq m$ , in which case  $m$  is called a *lower bound* of the set  $E$ .
2. A number  $t$  is called an *infimum* of the set  $E$  if and only if  $t$  is a lower bound of  $E$  and  $t \geq m$  and write  $t = \inf E$ .
3.  $E$  is said to be *bounded* if and only if it is bounded both above and below.

- **Theorem 1.20** *Reflection Principle*

Let  $E \subseteq \mathbb{R}$  be nonempty

1.  $E$  has a supremum if and only if  $-E$  has an infimum, in which case

$$\inf(-E) = -\sup E$$

2.  $E$  has an infimum if and only if  $-E$  has a supremum, in which case

$$\sup(-E) = -\inf E$$

- **Theorem 1.21** *Monotone Property*

Suppose that  $A \subseteq B$  are nonempty subsets of  $\mathbb{R}$ .

1. If  $B$  has a supremum, then  $\sup A \leq \sup B$ .
2. If  $B$  has an infimum, then  $\inf A \geq \inf B$ .

## 1.4 Mathematical Induction

- **Theorem 1.22** *Well-Ordering Principle*

If  $E$  is a nonempty subset of  $\mathbb{N}$ , then  $E$  has a least element (i.e.  $E$  has a finite infimum and  $\inf E \in E$ ).

- **Theorem 1.23**

Suppose for each  $n \in \mathbb{N}$  that  $A(n)$  is a proposition which satisfies the following two properties:

1.  $A(1)$  is true.
2. For every  $n \in \mathbb{N}$  for which  $A(n)$  is true,  $A(n+1)$  is also true.

Then  $A(n)$  is true for all  $n \in \mathbb{N}$ .

- **Theorem 1.26** *Binomial Formula*

If  $a, b \in \mathbb{R}, n \in \mathbb{N}$  and  $0^0$  is interpreted to be 1, then

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$$

## 1.5 Inverse Functions and Images

- **Definition 1.29** *Injection, Surjection, Bijection*

Let  $X$  and  $Y$  be sets and  $f : X \rightarrow Y$

1.  $f$  is said to be *injective* if and only if

$$x_1, x_2 \in X \text{ and } f(x_1) = f(x_2) \implies x_1 = x_2$$

2.  $f$  is said to be *surjective* if and only if

$$\forall y \in Y \exists x \in X \ni y = f(x)$$

3.  $f$  is called *bijective* if and only if it is both injective and surjective

• **Theorem 1.30**

Let  $X$  and  $Y$  be sets and  $f : X \rightarrow Y$ . Then the following three statements are equivalent.

1.  $f$  has an inverse;
2.  $f$  is injective from  $X$  onto  $Y$ ;
3. There is a function  $g : Y \rightarrow X$  such that

$$g(f(x)) = x \quad \forall x \in X$$

and

$$f(g(y)) = y \quad \forall y \in Y$$

Moreover, for each  $f : X \rightarrow Y$ , there is only one function  $g$  that satisfies these. It is the inverse function  $f^{-1}$ .

• **Remark 1.31**

Let  $I$  be an interval and let  $f : I \rightarrow \mathbb{R}$ . If the derivative of  $f$  is either always positive on  $I$ , or always negative on  $I$ , then  $f$  is injective on  $I$ .

• **Definition 1.33** *Image*

Let  $X$  and  $Y$  be sets and  $f : X \rightarrow Y$ . The *image* of a set  $E \subseteq X$  under  $f$  is the set

$$f(E) := \{y \in Y : y = f(x) \text{ for some } x \in E\}$$

The *inverse image* of a set  $E \subseteq Y$  under  $f$  is the set

$$f^{-1}(E) := \{x \in X : f(x) = y \text{ for some } y \in E\}$$

• **Definition 1.35** *Union, Intersection*

Let  $\mathcal{E} = \{E_\alpha\}_{\alpha \in A}$  be a collection of sets.

1. The *union* of the collection  $\mathcal{E}$  is the set

$$\bigcup_{\alpha \in A} E_\alpha := \{x : x \in E_\alpha \text{ for some } \alpha \in A\}$$

2. The *intersection* of the collection  $\mathcal{E}$  is the set

$$\bigcap_{\alpha \in A} E_\alpha := \{x : x \in E_\alpha \text{ for all } \alpha \in A\}.$$

• **Theorem 1.36** *DeMorgan's Laws*

Let  $X$  be a set and  $\{E_\alpha\}_{\alpha \in A}$  be a collection of subsets of  $X$ . If for each  $E \subseteq X$  the symbol  $E^c$  represents the set  $X \setminus E$ , then

$$\left( \bigcup_{\alpha \in A} E_\alpha \right)^c = \bigcap_{\alpha \in A} E_\alpha^c$$

and

$$\left( \bigcap_{\alpha \in A} E_\alpha \right)^c = \bigcup_{\alpha \in A} E_\alpha^c$$

• **Theorem 1.37**

Let  $X$  and  $Y$  be sets and  $f : X \rightarrow Y$ .

1. If  $\{E_\alpha\}_{\alpha \in A}$  is a collection of subsets of  $X$ , then

$$f\left(\bigcup_{\alpha \in A} E_\alpha\right) = \bigcup_{\alpha \in A} f(E_\alpha) \quad \text{and} \quad f\left(\bigcap_{\alpha \in A} E_\alpha\right) \subseteq \bigcap_{\alpha \in A} f(E_\alpha)$$

2. If  $B$  and  $C$  are subsets of  $X$ , then  $f(C) \setminus f(B) \subseteq f(C \setminus B)$

3. If  $\{E_\alpha\}_{\alpha \in A}$  is a collection of subsets of  $Y$ , then

$$f^{-1}\left(\bigcup_{\alpha \in A} E_\alpha\right) = \bigcup_{\alpha \in A} f^{-1}(E_\alpha) \quad \text{and} \quad f^{-1}\left(\bigcap_{\alpha \in A} E_\alpha\right) = \bigcap_{\alpha \in A} f^{-1}(E_\alpha)$$

4. If  $B$  and  $C$  are subsets of  $Y$ , then  $f^{-1}(C \setminus B) = f^{-1}(C) \setminus f^{-1}(B)$ .

5. If  $E \subseteq f(X)$ , then  $f(f^{-1}(E)) = E$ , but if  $E \subseteq X$ , then  $E \subseteq f^{-1}(f(E))$ .

## 1.6 Countable and Uncountable Sets

• **Definition 1.38** *Countable Uncountable* Let  $E$  be a set.

1.  $E$  is said to be *finite* if and only if either  $E = \emptyset$  or there exists an injective function which takes  $\{1, 2, \dots, n\}$  onto  $E$ , for some  $n \in \mathbb{N}$ .
2.  $E$  is said to be *countable* if and only if there exists an injective function which takes  $\mathbb{N}$  onto  $E$ .
3.  $E$  is said to be *at most countable* if and only if  $E$  is either finite or countable.
4.  $E$  is said to be *uncountable* if and only if  $E$  is neither finite nor countable.

• **Remark 1.39** *Cantor's Diagonalisation Argument*

The open interval  $(0, 1)$  is uncountable.

• **Lemma 1.40**

A nonempty set  $E$  is at most countable if and only if there is a function  $g$  from  $\mathbb{N}$  onto  $E$ .

• **Theorem 1.41**

Suppose  $A$  and  $B$  are sets.

1. If  $A \subseteq B$  and  $B$  is at most countable, then  $A$  is at most countable.
2. If  $A \subseteq B$  and  $A$  is uncountable, then  $B$  is uncountable.
3.  $\mathbb{R}$  is uncountable.

• **Theorem 1.42**

Let  $A_1, A_2, \dots$  be at most countable sets.

1. Then  $A_1 \times A_2$  is at most countable.
2. If

$$E = \bigcup_{j=1}^{\infty} A_j := \bigcup_{j \in \mathbb{N}} A_j := \{x : x \in A_j \text{ for some } j \in \mathbb{N}\},$$

then  $E$  is at most countable.

• **Remark 1.43**

The sets  $\mathbb{Z}$  and  $\mathbb{Q}$  are countable, but the set of irrationals is uncountable.

## 2 Sequences in $\mathbb{R}$

### 2.1 Limits of Sequences

- **Definition 2.1** *Convergence*

A sequence of real numbers  $\{x_n\}$  is set to *converge* to a real number  $a \in \mathbb{R}$  if and only if for every  $\epsilon > 0$  there is an  $N \in \mathbb{N}$  (which in general depends on  $\epsilon$ ) such that

$$n \geq N \implies |x_n - a| < \epsilon$$

- **Remark 2.4** A sequence can have at most one limit.

- **Definition 2.5** *Subsequence*

By a *subsequence* of a sequence  $\{x_n\}_{n \in \mathbb{N}}$ , we shall mean a sequence of the form  $\{x_{n_k}\}_{k \in \mathbb{N}}$ , where each  $n_k \in \mathbb{N}$  and  $n_1 < n_2 < \dots$ .

- **Remark 2.6**

If  $\{x_n\}_{n \in \mathbb{N}}$  converges to  $a$  and  $\{x_{n_k}\}_{k \in \mathbb{N}}$  is any subsequence of  $\{x_n\}_{n \in \mathbb{N}}$ , then  $x_{n_k}$  converges to  $a$  as  $k \rightarrow \infty$ .

- **Definition 2.7** *Bounded Sequences*

Let  $\{x_n\}$  be a sequence of real numbers.

1. The sequence  $\{x_n\}$  is said to be *bounded above* if and only if the set  $\{x_n : n \in \mathbb{N}\}$  is bounded above.
2. The sequence  $\{x_n\}$  is said to be *bounded below* if and only if the set  $\{x_n : n \in \mathbb{N}\}$  is bounded below.
3.  $\{x_n\}$  is said to be *bounded* if and only if it is bounded both above and below.

- **Theorem 2.8** Every convergent sequence is bounded.

### 2.2 Limit Theorems

- **Theorem 2.9** *Squeeze Theorem*

Suppose that  $\{x_n\}$ ,  $\{y_n\}$ , and  $\{w_n\}$  are real sequences.

1. If  $x_n \rightarrow a$  and  $y_n \rightarrow a$  as  $n \rightarrow \infty$ , and if there is an  $N_0 \in \mathbb{N}$  such that

$$x_n \leq w_n \leq y_n \text{ for } n \geq N_0$$

then  $w_n \rightarrow a$  as  $n \rightarrow \infty$ .

2. If  $x_n \rightarrow 0$  as  $n \rightarrow \infty$  and  $\{y_n\}$  is bounded, then  $x_n y_n \rightarrow 0$  as  $n \rightarrow \infty$ .

- **Theorem 2.11**

Let  $E \subset \mathbb{R}$ . If  $E$  has a finite supremum (respectively, a finite infimum), then there is a sequence  $x_n \in E$  such that  $x_n \rightarrow \sup E$  (respectively, a sequence  $y_n \in E$  such that  $y_n \rightarrow \inf E$ ) as  $n \rightarrow \infty$ .

- **Theorem 2.12**

Suppose that  $\{x_n\}$  and  $\{y_n\}$  are real sequences and that  $\alpha \in \mathbb{R}$ . If  $\{x_n\}$  and  $\{y_n\}$  are convergent, then

- 1.

$$\lim_{n \rightarrow \infty} (x_n + y_n) = \lim_{n \rightarrow \infty} x_n + \lim_{n \rightarrow \infty} y_n$$

- 2.

$$\lim_{n \rightarrow \infty} (\alpha x_n) = \alpha \lim_{n \rightarrow \infty} x_n$$

and

3.

$$\lim_{n \rightarrow \infty} (x_n y_n) = \left( \lim_{n \rightarrow \infty} x_n \right) \left( \lim_{n \rightarrow \infty} y_n \right)$$

If, in addition,  $y_n \neq 0$  and  $\lim_{n \rightarrow \infty} y_n \neq 0$ , then

4.

$$\lim_{n \rightarrow \infty} \frac{x_n}{y_n} = \frac{\lim_{n \rightarrow \infty} x_n}{\lim_{n \rightarrow \infty} y_n}$$

(In particular, all these limits exist.)

• **Definition 2.14** *Divergence*

Let  $\{x_n\}$  be a sequence of real numbers.

1.  $\{x_n\}$  is said to *diverge* to  $+\infty$  if and only if for each  $M \in \mathbb{R}$  there is an  $N \in \mathbb{N}$  such that

$$n \geq N \implies x_n > M$$

2.  $\{x_n\}$  is said to *diverge* to  $-\infty$  if and only if for each  $M \in \mathbb{R}$  there is an  $N \in \mathbb{N}$  such that

$$n \geq N \implies x_n < M$$

• **Theorem 2.15**

Suppose that  $\{x_n\}$  and  $\{y_n\}$  are real sequences such that  $x_n \rightarrow +\infty$  (respectively,  $x_n \rightarrow -\infty$ ) as  $n \rightarrow \infty$ .

1. If  $y_n$  is bounded below (respectively,  $y_n$  is bounded above), then

$$\lim_{n \rightarrow \infty} (x_n + y_n) = +\infty \quad (\text{respectively, } \lim_{n \rightarrow \infty} (x_n + y_n) = -\infty)$$

2. If  $\alpha > 0$ , then

$$\lim_{n \rightarrow \infty} (\alpha x_n) = +\infty \quad (\text{respectively, } \lim_{n \rightarrow \infty} (\alpha x_n) = -\infty)$$

3. If  $y_n > M_0$  for some  $M_0 > 0$  and all  $n \in \mathbb{N}$ , then

$$\lim_{n \rightarrow \infty} (x_n y_n) = +\infty \quad (\text{respectively, } \lim_{n \rightarrow \infty} (x_n y_n) = -\infty)$$

4. If  $\{y_n\}$  is bounded and  $x_n \neq 0$ , then

$$\lim_{n \rightarrow \infty} \frac{y_n}{x_n} = 0$$

• **Corollary 2.16**

Let  $\{x_n\}$ ,  $\{y_n\}$  be real sequences and  $\alpha, x, y$  be extended real numbers. If  $x_n \rightarrow x$  and  $y_n \rightarrow y$ , as  $n \rightarrow \infty$ , then

$$\lim_{n \rightarrow \infty} (x_n + y_n) = x + y$$

provided that the right side is not of the form  $\infty - \infty$ , and

$$\lim_{n \rightarrow \infty} (\alpha x_n) = \alpha x, \quad \lim_{n \rightarrow \infty} (x_n y_n) = xy$$

provided that none of these products is of the form  $0 \cdot \pm\infty$ .

• **Theorem 2.17** *Comparison Theorem*

Suppose that  $\{x_n\}$  and  $\{y_n\}$  are convergent sequences. If there is an  $N_0 \in \mathbb{N}$  such that

$$x_n \leq y_n \text{ for } n \geq N_0$$

then

$$\lim_{n \rightarrow \infty} x_n \leq \lim_{n \rightarrow \infty} y_n$$

In particular, if  $x_n \in [a, b]$  converges to some point  $c$ , then  $c$  must belong to  $[a, b]$ .



## 2.3 Bolzano-Weierstrass Theorem

- **Definition 2.18** *Increasing, Decreasing* Let  $\{x_n\}_{n \in \mathbb{N}}$  be a sequence of real numbers.
  1.  $\{x_n\}$  is said to be *increasing* (respectively, *strictly increasing*) if and only if  $x_1 \leq x_2 \leq \dots$  (respectively,  $x_1 < x_2 < \dots$ ).
  2.  $\{x_n\}$  is said to be *decreasing* (respectively, *strictly decreasing*) if and only if  $x_1 \geq x_2 \geq \dots$  (respectively,  $x_1 > x_2 > \dots$ ).
  3.  $\{x_n\}$  is said to be *monotone* if and only if it is either increasing or decreasing.
- **Theorem 2.19** *Monotone Convergence Theorem*  
if  $\{x_n\}$  is increasing and bounded above, or if  $\{x_n\}$  is decreasing and bounded below, then  $\{x_n\}$  converges to a finite limit.
- **Definition 2.22** *Nested*  
A sequence of sets  $\{I_n\}_{n \in \mathbb{N}}$  is said to be *nested* if and only if

$$I_1 \supseteq I_2 \supseteq \dots$$

- **Theorem 2.23** *Nested Interval Property*  
If  $\{I_n\}_{n \in \mathbb{N}}$  is a nested sequence of nonempty closed bounded intervals, then  $E := \bigcap_{n=1}^{\infty} I_n$  is nonempty. Moreover, if the lengths of these intervals satisfy  $|I_n| \rightarrow 0$  as  $n \rightarrow \infty$  then  $E$  is a single point.
- **Remark 2.24** The Nested Interval Property might not hold if “closed” is omitted.
- **Remark 2.25** The Nested Interval Property might not hold if “bounded” is omitted.
- **Theorem 2.26** *Bolzano-Weierstrass Theorem*  
Every bounded sequence of real numbers has a convergent subsequence.

## 2.4 Cauchy Sequences

- **Definition 2.27** *Cauchy*  
A sequence of points  $x_n \in \mathbb{R}$  is said to be *Cauchy* (in  $\mathbb{R}$ ) if and only if for every  $\epsilon > 0$  there is an  $N \in \mathbb{N}$  such that
 
$$n, m \geq N \implies |x_n - x_m| < \epsilon$$
- **Remark 2.28** If  $\{x_n\}$  is convergent, then  $\{x_n\}$  is Cauchy.
- **Theorem 2.29** *Cauchy*  
Let  $\{x_n\}$  be a sequence of real numbers. Then  $\{x_n\}$  is Cauchy if and only if  $\{x_n\}$  converges (to some point  $a \in \mathbb{R}$ ).
- **Remark 2.31** A sequence that satisfies  $x_{n+1} - x_n \rightarrow 0$  is not necessarily Cauchy.

## 2.5 Limits Supremum and Infimum

- **Definition 2.32** *Limit Supremum & Infimum*  
Let  $\{x_n\}$  be a real sequence. Then the *limit supremum* of  $\{x_n\}$  is the extended real number

$$\limsup_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} (\sup_{k \geq n} x_k)$$

and the *limit infimum* of  $\{x_n\}$  is the extended real number

$$\liminf_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} (\inf_{k \geq n} x_k)$$

- **Theorem 2.35**

Let  $\{x_n\}$  be a sequence of real numbers,  $s = \limsup_{n \rightarrow \infty} x_n$ , and  $t = \liminf_{n \rightarrow \infty} x_n$ . Then there are subsequences  $\{x_{n_k}\}_{k \in \mathbb{N}}$  and  $\{x_j\}_{j \in \mathbb{N}}$  such that  $x_{n_k} \rightarrow s$  as  $k \rightarrow \infty$  and  $x_j \rightarrow t$  as  $j \rightarrow \infty$ .

- **Theorem 2.36**

Let  $\{x_n\}$  be a real sequence and  $x$  be an extended real number. Then  $x_n \rightarrow x$  as  $n \rightarrow \infty$  if and only if

$$\limsup_{n \rightarrow \infty} x_n = \liminf_{n \rightarrow \infty} x_n = x$$

- **Theorem 2.37**

Let  $\{x_n\}$  be a sequence of real numbers. Then  $\limsup_{n \rightarrow \infty} x_n$  (respectively,  $\liminf_{n \rightarrow \infty} x_n$ ) is the largest value (respectively, the smallest value) to which some subsequences of  $\{x_n\}$  converges. Namely, if  $x_{n_k} \rightarrow x$  as  $k \rightarrow \infty$ , then

$$\liminf_{n \rightarrow \infty} x_n \leq x \leq \limsup_{n \rightarrow \infty} x_n$$

- **Remark 2.38** If  $\{x_n\}$  is any sequence of real numbers, then

$$\liminf_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} x_n$$

- **Remark 2.39** A real sequence  $\{x_n\}$  is bounded above if and only if  $\limsup_{n \rightarrow \infty} x_n < \infty$ , and is bounded below if and only if  $\liminf_{n \rightarrow \infty} x_n > -\infty$ .

- **Theorem 2.40**

If  $x_n \leq y_n$  for  $n$  large, then

$$\limsup_{n \rightarrow \infty} x_n \leq \limsup_{n \rightarrow \infty} y_n \quad \text{and} \quad \liminf_{n \rightarrow \infty} y_n \leq \liminf_{n \rightarrow \infty} x_n$$