

Reducing the Computation of Linear Complexities of Periodic Sequences over $GF(p^m)$

Hao Chen

Abstract—The linear complexity of a periodic sequence over $GF(p^m)$ plays an important role in cryptography and communication [12]. In this correspondence, we prove a result which reduces the computation of the linear complexity and minimal connection polynomial of a period un sequence over $GF(p^m)$ to the computation of the linear complexities and minimal connection polynomials of u period n sequences. The conditions $u|p^m - 1$ and $\gcd(n, p^m - 1) = 1$ are required for the result to hold. Some applications of this reduction in fast algorithms to determine the linear complexities and minimal connection polynomials of sequences over $GF(p^m)$ are presented.

Index Terms—Berlekamp-Massey algorithm, Games-Chan algorithm, linear complexity, minimal connection polynomial, cryptography

I. INTRODUCTION

For a period N sequence $\mathbf{a} = a_0, a_1, \dots, a_{N-1}, a_0, \dots$ over a finite field $GF(p^m)$, its linear complexity $c(\mathbf{a})$ is defined to be the length of the shortest linear feedback shift register to generate it, i.e. the smallest positive integer k such that there exist some c_1, \dots, c_k in $GF(p^m)$ and $a_{i+k} = c_1 a_{i+k-1} + \dots + c_k a_i$ hold for all $i \geq 0$. The polynomial $m(\mathbf{a}) = 1 - (c_1 x + \dots + c_k x^k)$ is called the minimal connection polynomial [12].

The linear complexity of a periodic sequence is considered as the measure of its randomness and plays an important role in the application of the sequence in cryptography and communication. There are many works [1],[2],[4],[6],[8],[9],[10],[11],[14],[15] and [16] on efficient algorithms for determining the linear complexities and minimal connection polynomials of sequences. Some authors also have interesting results about the linear complexities of some special sequences (see [3],[7] and [13]). The famous Berlekamp-Massey algorithm [11] can be used to compute the linear complexity and minimal connection polynomial of a period N sequence over $GF(p^m)$ with time complexity $O(N^2)$ (that is, at most $O(N^2)$ field operations in $GF(p^m)$). One of the main advantages of the Berlekamp-Massey algorithm is the input at the step t of the algorithm is the first t elements of the sequence. Actually, the Berlekamp-Massey algorithm only needs $2c(\mathbf{a})$ consecutive elements of the sequence to determine its linear complexity and minimal connection polynomial [12]. An adapted fast version of Berlekamp-Massey algorithm due to Blackburn [1] can be used with time complexity $O(N(\log N)^2 \log \log N)$.

In [6] Games and Chan gave a fast algorithm which can be used to determine the linear complexity and minimal

connection polynomial of a period $N = 2^t$ binary sequence with time complexity $O(N)$. This algorithm was also generalized to compute the linear complexity and minimal connection polynomial of a period $N = p^t$ sequence over $GF(p^m)$ with time complexity $O(N)$ (see [5] and [8]). Based on the Games-Chan algorithm, some authors developed fast algorithms [9], [10] and [14] for computing the k -error linear complexities of period $N = 2^t$ binary sequences and period $N = p^t$ sequences over $GF(p^m)$. G.Xiao et al. [15] and [16] gave fast algorithms to compute the linear complexities and minimal connection polynomials of period $N = p^t$ or $N = 2p^t$ sequences over $GF(q)$, when q is a primitive root modulo p^2 . For sequences of period $N = 2^t n$, where $2^t | p^m - 1$ and $\gcd(n, p^m - 1) = 1$, a fast algorithm which can be used to determine their linear complexities more efficiently was given in our paper [4].

It is well known that the linear complexity and minimal connection polynomial of a periodic sequence over $GF(p^m)$ can be understood from its generating function. For a sequence $\mathbf{a} = a_0, a_1, \dots, a_{N-1}, a_0, \dots$ over $GF(p^m)$ of period N , its generating function $A(x) = a_0 + a_1 x + \dots + a_i x^i + \dots = \sum_{i \geq 0} a_i x^i = \frac{a_0 + a_1 x + \dots + a_{N-1} x^{N-1}}{1 - x^N}$. Then the linear complexity of the sequence \mathbf{a} is $c(\mathbf{a}) = \deg(1 - x^N) - \deg(\gcd(a_0 + a_1 x + \dots + a_{N-1} x^{N-1}, 1 - x^N))$ and the minimal connection polynomial is $m(\mathbf{a})(x) = \frac{1 - x^N}{\gcd(a_0 + a_1 x + \dots + a_{N-1} x^{N-1}, 1 - x^N)}$ [12].

In this correspondence we prove a result which reduces the computation of the linear complexity and minimal connection polynomial of a period un sequence over $GF(p^m)$ to the computation of the linear complexities and minimal connection polynomials of u period n sequences. This reduction result can be combined with various known algorithms to compute the linear complexities of sequences more efficiently. The main result of this correspondence can be thought as a generalization of the result in our previous paper [4].

II. MAIN RESULT

Let m be a positive integer, p be a prime number, u be a positive integer such that u divides $p^m - 1$, and n be a positive integer such that $\gcd(n, p^m - 1) = 1$. It is clear there are u distinct u -th roots of unity x_0, \dots, x_{u-1} , where $x_0 = 1$, in $GF(p^m)$ since $u | p^m - 1$. From the condition $\gcd(n, p^m - 1) = 1$, we can find a unique $b_i \in GF(p^m)$, which is the n -th root of x_i for all $i = 0, \dots, u - 1$. The following result is the main result of this correspondence.

Theorem. Suppose $p, m, u, n, x_0, \dots, x_{u-1}, b_0, \dots, b_{u-1}$ are given as above. Let $\mathbf{a} = a_0, a_1, \dots, a_{un-1}, a_0, a_1, \dots$ be a period un sequence over $GF(p^m)$. Let \mathbf{a}^j be the period n sequence over $GF(p^m)$ with its first period $a_0 + a_n b_j^n + \dots + a_{(u-1)n} b_j^{(u-1)n}, \dots, a_i b_j^i + a_{n+i} b_j^{n+i} + \dots + a_{(u-1)n+i} b_j^{(u-1)n+i}, \dots, a_{n-1} b_j^{n-1} + a_{2n-1} b_j^{2n-1} + \dots + a_{un-1} b_j^{un-1}$, for $j = 0, 1, \dots, u - 1$. Then $c(\mathbf{a}) = c(\mathbf{a}^0) + c(\mathbf{a}^1) + \dots + c(\mathbf{a}^{u-1})$ and

H. Chen is with the Department of Computing and Information Technology, School of Information Science and Engineering, Fudan University, Shanghai, 200433, People's Republic of China

$$m(\mathbf{a})(x) = m(\mathbf{a}^0)(b_0^{-1}x)m(\mathbf{a}^1)(b_1^{-1}x) \cdots m(\mathbf{a}^{u-1})(b_{u-1}^{-1}x).$$

Proof. Let $f(x) = \sum_{i=0}^{un-1} a_i x^i$. It is clear $1 - x^{un} = \prod_{i=0}^{u-1} (x_i - x^n) = x_1 \cdots x_{u-1} (1 - x^n) \prod_{i=1}^{u-1} (1 - (b_i^{-1}x)^n)$. Any two distinct polynomials among the u polynomials $(1 - x^n), (1 - (b_1^{-1}x)^n), \dots, (1 - (b_{u-1}^{-1}x)^n)$ are coprime in $GF(p^m)[x]$. Thus $\gcd(f(x), 1 - x^{un}) = \gcd(f(x), 1 - x^n) \prod_{i=1}^{u-1} \gcd(f(x), 1 - (b_i^{-1}x)^n)$.

It is clear $\gcd(f(x), 1 - x^n) = \gcd(f_0(x), 1 - x^n)$, where $f_0(x) = \sum_{i=0}^{n-1} (a_i + a_{n+i} + \cdots + a_{(u-1)n+i}) x^i$. Thus $c(\mathbf{a}^0) = \deg(\frac{1-x^n}{\gcd(f_0(x), 1-x^n)})$ and $m(\mathbf{a}^0)(x) = \frac{1-x^n}{\gcd(f_0(x), 1-x^n)}$. For each j satisfying $1 \leq j \leq u-1$, we set $\gcd(f(x), 1 - (b_j^{-1}x)^n) = g_j(x)$ and $\gcd(f(b_j y), 1 - y^n) = h_j(y)$. Then $g_j(x) = h_j(b_j^{-1}x)$. We have $h_j(y) = \gcd(f_j(y), 1 - y^n)$, where $f_j(y) = \sum_{i=0}^{n-1} (a_i b_j^i + a_{n+i} b_j^{n+i} + \cdots + a_{(u-1)n+i} b_j^{(u-1)n+i}) y^i$. Thus $c(\mathbf{a}^j) = \deg(\frac{1-y^n}{h_j(y)})$ and $m(\mathbf{a}^j)(y) = \frac{1-y^n}{h_j(y)}$. Finally $c(\mathbf{a}) = un - [\sum_{i=0}^{u-1} \deg(\gcd(f(x), 1 - (b_i^{-1}x)^n))] = c(\mathbf{a}^0) + c(\mathbf{a}^1) + \cdots + c(\mathbf{a}^{u-1})$ and $m(\mathbf{a})(x) = m(\mathbf{a}^0)(b_0^{-1}x)m(\mathbf{a}^1)(b_1^{-1}x) \cdots m(\mathbf{a}^{u-1})(b_{u-1}^{-1}x)$. The conclusion is proved.

When $u = 2^t n$, the above result was proved in our previous paper [4].

In the reduction we need the storage of u elements $b_0 = 1, b_1, \dots, b_{u-1} \in GF(p^m)$ in advance. For a period $N = un$ sequence over $GF(p^m)$, where $u|p^m - 1$ and $\gcd(n, p^m - 1) = 1$, we need $\frac{(u-1)N}{u}$ field operations to get the sequence \mathbf{a}^0 , $(u-1)N$ field operations to get the elements $b_1, \dots, b_1^{(u-1)n-1}, \dots, b_{u-1}, \dots, b_{u-1}^{(u-1)n-1}$, and $\frac{(2u-1)(u-1)N}{u}$ field operations to get the sequences $\mathbf{a}^1, \dots, \mathbf{a}^{u-1}$. Thus the time complexity of the reduction in the main result is $3(u-1)N$ field operations in $GF(p^m)$.

III. APPLICATIONS

In this section we use the main result and some known algorithms to give fast algorithms for computing the linear complexities of sequences over $GF(p^m)$.

A. An easy example

Let p be an odd prime, m be an arbitrary positive integer and n be a positive integer such that n and $p^m - 1$ are coprime. Then we have a unique element b in $GF(p^m)$ such that $b^n = -1$. Here we note $b^{2n} = 1$. For arbitrary $a_0, \dots, a_{n-1} \in GF(p^m)$, let $\mathbf{a} = a_0, a_1, \dots, a_{n-1}, -a_0, -a_1, \dots, -a_{n-1}, a_0, \dots$ be a period $2n$ sequence over $GF(p^m)$. Set $\mathbf{a}' = 2a_0, 2a_1 b, \dots, 2a_i b^i, \dots, 2a_{n-1} b^{n-1}, 2a_0, \dots$, which is a period n sequence over $GF(p^m)$. From the main result, the linear complexity $c(\mathbf{a})$ is the same as the linear complexity $c(\mathbf{a}')$ and the minimal connection polynomial $m(\mathbf{a})(x)$ is just $m(\mathbf{a}')(bx)$. Thus the linear complexity and minimal connection polynomial of the period $2n$ sequence \mathbf{a} can be determined from the period n sequence \mathbf{a}' .

B. Combining with the generalized Games-Chan algorithm

In this subsection it is assumed that p is a prime number, m is a positive integer and u is a positive integer such that u divides $p^m - 1$. We now give a fast algorithm to compute the linear complexity $c(\mathbf{a})$ of a period $N = up^h$ sequence \mathbf{a} over $GF(p^m)$ with time complexity $O(N)$. Here u is understood as a constant not depending on the sequence. We need the storage of u elements $b_0 = 1, b_1, \dots, b_{u-1}$ in advance.

Input: A period $N = up^h$ sequence \mathbf{a} over $GF(p^m)$.

Output: The linear complexity $c(\mathbf{a})$.

Algorithm.

Perform the reduction of the main result, we get u period p^h sequences $\mathbf{a}^0, \dots, \mathbf{a}^{u-1}$.

For the period p^h sequences $\mathbf{a}^0, \dots, \mathbf{a}^{u-1}$, perform the following generalized Games-Chan algorithm **GGC**, the outputs are the linear complexities $c(\mathbf{a}^0), \dots, c(\mathbf{a}^{u-1})$.

GGC Algorithm.

- 1) Initial value: $\mathbf{s} \leftarrow \mathbf{s} = (s_0, \dots, s_{p^h-1}) \in GF(p^m)^{p^h}$, $N \leftarrow p^h$, $c \leftarrow 0$.
- 2) Repeat the following a)-c) until $h = 0$.
 - a) For a given p^h -tuple \mathbf{s} , set $\mathbf{s}^{(i)} = (s_{ip^{h-1}}, \dots, s_{(i+1)p^{h-1}-1})$ for $i = 0, \dots, p-1$, and $\mathbf{b}^{(u)} = \sum_{j=0}^{p-u-1} C_{p-j-1}^u \mathbf{s}^{(j)}$, where $u = 0, \dots, p-1$ and C_{p-u-1}^u 's are the binomial coefficients.
 - b) Find the smallest w such that $\mathbf{b}^{(0)} = \mathbf{b}^{(1)} = \dots = \mathbf{b}^{(p-w-1)} = 0$ and $\mathbf{b}^{(p-w)} \neq 0$ for a $w \in \{1, \dots, p\}$. Here if $\mathbf{b}^{(0)} \neq 0$, we set $w = p$.
 - c) Do $\mathbf{s} \leftarrow \mathbf{b}^{(p-w)}$, $c \leftarrow (w-1)p^{h-1} + c$, and goto a).
- 3) When $h = 0$ and $\mathbf{s} = (s_0) \neq 0$, then $c \leftarrow c + 1$, otherwise $c \leftarrow c$.

The final output c of **GGC** is the linear complexity $c(\mathbf{s})$ of the period p^h sequence \mathbf{s} over $GF(p^m)$.

Finally we get the linear complexity of $c(\mathbf{a}) = \sum_{i=0}^{u-1} c(\mathbf{a}^i)$ from the main result.

We refer to [5],[8] and [10] for the generalized Games-Chan algorithm. **GGC** needs at most $2p^2 N'$ field operations in $GF(p^m)$ for determining the linear complexity of a period $N' = p^h$ sequence over $GF(p^m)$. On the other hand we need at most $3(u-1)N$ field operations in the reduction for a given period $N = uN'$ sequence. Thus the above algorithm needs $3(u-1)N + u(2p^2 \frac{N}{u}) = [3(u-1) + 2p^2]N$ field operations in $GF(p^m)$, where N is the period of the input

sequence. The coefficient $3(u-1) + 2p^2$ is a fixed constant not depending on the sequence. For example, the above fast algorithm can be used to determine the linear complexities of period $N = 3 \cdot 7^h$ sequences over $GF(7^m)$ and period $N = 3 \cdot 13^h$ sequences over $GF(13^m)$.

Example. Let $\mathbf{a} = 123401520113061256331\dots$ be a period 21 sequence over $GF(7)$. We want to compute its linear complexity and minimal connection polynomial by the above algorithm. First we note $b_0 = 1$, $b_1 = 4$ and $b_2 = 2$ in $GF(7)$. Then

$$\mathbf{a}^0 = 4424645, \mathbf{a}^1 = 4366203, \mathbf{a}^2 = 2622130.$$

$$c(\mathbf{a}) = c(\mathbf{a}^0) + c(\mathbf{a}^1) + c(\mathbf{a}^2).$$

$$m(\mathbf{a})(x) = m(\mathbf{a}^0)(x)m(\mathbf{a}^1)(4x)m(\mathbf{a}^2)(2x).$$

In the case of $p = 7$ we use the generalized Games-Chan algorithm and get

$$c(\mathbf{a}^0) = 7, m(\mathbf{a}^0) = (1-x)^7,$$

$$c(\mathbf{a}^1) = 7, m(\mathbf{a}^1) = (1-x)^7,$$

$$c(\mathbf{a}^2) = 7, m(\mathbf{a}^2) = (1-x)^7.$$

Finally we have $c(\mathbf{a}) = 21$ and $m(\mathbf{a}) = (1-x)^7(1-4x)^7(1-2x)^7$.

Comparing with the Blackburn's algorithm given in [2], the reduction to the u period p^h sequences is the same as that in the Blackburn's algorithm, because in this case the u -th root of unity α in [2] is an element of $GF(p^m)$.

C. Combining with the Berlekamp-Massey algorithm

We can also apply the reduction of the main result to compute the linear complexity of a period $N = un$ ($\gcd(n, p^m - 1) = 1$) sequence \mathbf{a} over $GF(p^m)$, where u divides $p^m - 1$ and n is not a power of p . In this case, we apply the Berlekamp-Massey algorithm [11] with time complexity $O(n^2)$ (or the Blackburn's version [1] of Berlekamp-Massey algorithm with time complexity $O(n(\log n)^2 \log \log n)$) to the u period n sequences after the reduction. It is obvious that this would be more efficient than applying the Berlekamp-Massey algorithm directly to the original sequence. However when this reduction is used, we have to know the whole period of the sequence.

D. Combining with the Xiao-Wei-Lam-Imamura algorithm

Let p and q be two prime numbers. Suppose q is a primitive root modulo p^2 , that is, q is the generator of the multiplicative group of residue classes (modulo p^2) which are coprime to p , then a fast algorithm for determining the linear complexity of a period $N = p^n$ sequence over $GF(q^m)$ with

time complexity $O(N)$ was given in [16]. Combining with the reduction in our main result, we can determine the linear complexity of a period $N = up^n$ sequence over $GF(q^m)$ with time complexity $O(N)$, if u divides $q^m - 1$, q is a primitive root modulo p^2 , p and $q^m - 1$ are coprime. For example, it is easy to check that 13 is a primitive root modulo 25, thus we can determine the linear complexities of period $N = 3 \cdot 5^n$ sequences over $GF(13^m)$ (if $m \neq 0 \pmod{4}$) with time complexity $O(N)$.

IV. Conclusion

We have proved a result reducing the computation of the linear complexity of a period un sequence over $GF(p^m)$, where u divides $p^m - 1$ and $\gcd(n, p^m - 1) = 1$, to the computation of the linear complexities of u period n sequences. Based on this reduction and some known algorithms we can compute the linear complexities of period un sequences over $GF(p^m)$ more efficiently. It seems that the main result might be useful for other problems about the linear complexities of sequences over $GF(p^m)$.

Acknowledgment. The author wishes to thank the Associate Editor and the anonymous referees for their helpful comments and criticisms on the 1st version of the paper. This work was supported in part by NNSF of China under Grant 90607005 and Distinguished Young Scholar Grant 10225106.

e-mail: chen hao@fudan.edu.cn

REFERENCES

- [1] S.R.Blackburn, Fast rational interpolation, Reed-Solomon decoding and the linear complexity profiles of sequences, IEEE Trans. Inf. Theory, vol.43, no.2, pp.537-548, Mar. 1997.
- [2] S.R.Blackburn, A generalization of the Discrete Fourier Transform: determining the minimal polynomial of a periodic sequence, IEEE Trans. Inf. Theory, vol.40, no.5, pp.1702-1704, Sept. 1994.
- [3] S.R.Blackburn, The linear complexity of the self-shrinking generator, IEEE Trans. Inf. Theory, vol.43, no.5, pp.2073-2077, Sept. 1999.
- [4] H.Chen, Fast algorithms for determining the linear complexity of sequences over $GF(p^m)$ with period $2^t n$, IEEE Trans. Inf. Theory, vol.51, no.5, pp.1854-1856, May 2005.
- [5] C.Ding, G.Xiao and W.Shan, The stability theory of stream ciphers, Lecture Notes in Computer Science, vol.56, Springer-Verlag, 1991.
- [6] R.A.Games and A.H. Chan, A fast algorithm for determining the complexity of a binary sequence with period 2^n , IEEE Trans. Inf. Theory, vol.29, no.1, pp.144-146, Jan. 1983.

[7] F. Griffin and I.E.Shparlinski, On the linear complexity profile of the power generator. IEEE Trans. Inf. Theory, vol.46, no.5, pp.2159-2162, Sept.2000.

[8] K.Imamura and T.Moriuchi, A fast algorithm for determining the linear complexity of p-ary sequences with period p^n , p prime, IEICE Tech. Rep. IT 93-75, pp.73-78, 1993.

[9] T.Kaida, S.Uehara and K.Imamura, An algorithm for the k-error linear complexity of sequences over $GF(p^m)$ with period p^n , p prime, Information and Computation, vol. 151, pp.134-147, 1999.

[10] A.G.Lauder and K.G.Paterson, Computation the error linear complexity spectrum of a binary sequence with period 2^n , IEEE Trans. Inf. Theory, vol.49, no.1, pp.273-280, Jan.2003.

[11] J.L.Massey, Shift register synthesis and BCH decoding, IEEE Trans. Inf. Theory, vol.15, no.1, pp.122-127, Jan.1969.

[12] A. Menezes, P. van Oorschot and S.Vanstone, Handbook of applied cryptography, CRC Press Inc. 1997.

[13] I.E.Shparlinski, On the linear complexity of the power generator, Designs, Codes and Cryptography, 2001, vol.23, no.1, pp.5-10, Jan.2001.

[14] M.Stamp and C.F.Martin, An algorithm for the k-error linear complexity of binary sequences with period 2^n , IEEE Trans. Inf. Theory, vol.39, no.4, pp.1398-1401, July 1998.

[15] S.Wei, G.Xiao and Z.Chen, A fast algorithm for determining the minimal polynomial of a sequence with period $2p^n$ over $GF(q)$, IEEE Trans. Inf. Theory, vol.48, no.10, pp2754-2758, Oct.2002.

[16] G.Xiao, S.Wei, K.Y.Lam and K.Imamura, A fast algorithm for determining the linear complexity of a sequence with period p^n over $GF(q)$, IEEE Trans. Inf. Theory, vol.46, no.6, pp.2203-2206, Sept.2000.

Hao Chen was born in Anhui Province of China on Dec. 21, 1964. He is currently a professor in the Department of Computing and Information Technology of Fudan University, Shanghai, China. His research interests are cryptography and coding, quantum information and computation.