

Properties of the Discrete Pulse Transform for Multi-Dimensional Arrays

Roumen Anguelov and Inger Fabris-Rotelli
Department of Mathematics and Applied Mathematics
University of Pretoria
roumen.anguelov@up.ac.za
inger.fabris-rotelli@up.ac.za

ISBN: 978-1-86854-785-2

Technical Report

2010/01

1 Introduction

This report presents properties of the Discrete Pulse Transform on multi-dimensional arrays introduced earlier in [1]. The main result given here in Lemma 2.1 is also formulated in [4, Lemma 21]. However, the proof, being too technical, was omitted there and hence it appears in full in this publication.

2 The Lemma

The lemma which follows deals with two technical aspects of the Discrete Pulse Transform of a function $f \in \mathcal{A}(\mathbb{Z}^d)$ (where $\mathcal{A}(\mathbb{Z}^d)$ denotes a vector lattice). The first is that the Discrete Pulse representation of a function f , given by

$$f = \sum_{n=1}^N D_n(f),$$

can be written as the sum of individual pulses of each resolution layer $D_n(f)$. The second result in the lemma below indicates a form of linearity for the nonlinear LULU operators.

Lemma 2.1

Let $f \in \mathcal{A}(\mathbb{Z}^d)$, $\text{supp}(f) < \infty$, be such that f does not have local minimum sets or local maximum sets of size smaller than n , for some $n \in \mathbb{N}$. Then we have the following two results.

a)

$$(id - P_n)f = \sum_{i=1}^{\gamma^-(n)} \phi_{ni} + \sum_{j=1}^{\gamma^+(n)} \varphi_{nj}, \quad (1)$$

where $V_{ni} = \text{supp}(\phi_{ni})$, $i = 1, 2, \dots, \gamma^-(n)$, are local minimum sets of f of size n , $W_{nj} = \text{supp}(\varphi_{nj})$, $j = 1, 2, \dots, \gamma^+(n)$, are local maximum sets of f of size n , ϕ_{ni} and φ_{nj} are negative and positive discrete pulses respectively, and we also have that

$$\bullet V_{ni} \cap V_{nj} = \emptyset \text{ and } \text{adj}(V_{ni}) \cap V_{nj} = \emptyset, \quad i, j = 1, \dots, \gamma^-(n), \quad i \neq j, \quad (2)$$

$$\bullet W_{ni} \cap W_{nj} = \emptyset \text{ and } \text{adj}(W_{ni}) \cap W_{nj} = \emptyset, \quad i, j = 1, \dots, \gamma^+(n), \quad i \neq j, \quad (3)$$

$$\bullet V_{ni} \cap W_{nj} = \emptyset \quad i = 1, \dots, \gamma^-(n), \quad j = 1, \dots, \gamma^+(n). \quad (4)$$

b) For every fully trend preserving operator A

$$U_n(id - AU_n) = U_n - AU_n,$$

$$L_n(id - AL_n) = L_n - AL_n.$$

Proof.

a) Let $V_{n1}, V_{n2}, \dots, V_{n\gamma^-(n)}$ be all local minimum sets of size n of the function f . Since f does not have local minimum sets of size smaller than n , then f is a constant on each of these sets, by [4, Theorem 14]. Hence, the sets are disjoint, that is $V_{ni} \cap V_{nj} = \emptyset$, $i \neq j$. Moreover, we also have

$$\text{adj}(V_{ni}) \cap V_{nj} = \emptyset, \quad i, j = 1, \dots, \gamma^-(n). \quad (5)$$

Indeed, let $x \in \text{adj}(V_{ni}) \cap V_{nj}$. Then there exists $y \in V_{ni}$ such that $(x, y) \in r$. Hence $y \in V_{ni} \cap \text{adj}(V_{nj})$. From the local minimality of the sets V_{ni} and V_{nj} we obtain respectively $f(y) < f(x)$ and $f(x) < f(y)$, which is clearly a contradiction. For every $i = 1, \dots, \gamma^-(n)$ denote by y_{ni} the point in $\text{adj}(V_{ni})$ such that

$$f(y_{ni}) = \min_{y \in \text{adj}(V_{ni})} f(y). \quad (6)$$

Then we have

$$U_n f(x) = \begin{cases} f(y_{ni}) & \text{if } x \in V_{ni}, i = 1, \dots, \gamma^-(n) \\ f(x) & \text{otherwise (by [4, Theorem 9])} \end{cases}$$

Therefore

$$(id - U_n)f = \sum_{i=1}^{\gamma^-(n)} \phi_{ni} \quad (7)$$

where ϕ_{ni} is a discrete pulse with support V_{ni} and negative value (down pulse).

Let $W_{n1}, W_{n2}, \dots, W_{n\gamma^+(n)}$ be all local maximum sets of size n of the function $U_n f$. By [4, Theorem 12(b)] every local maximum set of $U_n f$ contains a local maximum set of f . Since f does not have local maximum sets of size smaller than n , this means that the sets W_{nj} , $j = 1, \dots, \gamma^+(n)$, are all local maximum sets of f and f is constant on each of them. Similarly to the local minimum sets of f considered above we have $W_{ni} \cap W_{nj} = \emptyset$, $i \neq j$, and $\text{adj}(W_{ni}) \cap W_{nj} = \emptyset$, $i, j = 1, \dots, \gamma^+(n)$. Moreover, since $U_n(f)$ is constant on any of the sets $V_{ni} \cup \{y_{ni}\}$, $i = 1, \dots, \gamma^-(n)$, see [4, Theorem 14], we also have

$$(V_{ni} \cup \{y_{ni}\}) \cap W_{nj} = \emptyset, \quad i = 1, \dots, \gamma^-(n), \quad j = 1, \dots, \gamma^+(n), \quad (8)$$

which implies (4).

Further we have

$$L_n U_n f(x) = \begin{cases} U_n f(z_{nj}) & \text{if } x \in W_{nj}, j = 1, \dots, \gamma^+(n) \\ U_n f(x) & \text{otherwise} \end{cases}$$

where $z_{nj} \in \text{adj}(W_{nj})$, $j = 1, \dots, \gamma^+(n)$, are such that $U_n f(z_{nj}) = \max_{z \in \text{adj}(W_{nj})} U_n f(z)$.

Hence

$$(id - L_n)U_n f = \sum_{j=1}^{\gamma^+(n)} \varphi_{nj} \quad (9)$$

where φ_{nj} is a discrete pulse with support W_{nj} and positive value (up pulse). Thus we have shown that

$$(id - P_n)f = (id - U_n)f + (id - L_n)U_nf = \sum_{i=1}^{\gamma^-(n)} \phi_{ni} + \sum_{j=1}^{\gamma^+(n)} \varphi_{nj}.$$

b) Let the function $f \in \mathcal{A}(\mathbb{Z}^d)$ be such that it does not have any local minimum or local maximum sets of size less than n . Denote $g = (id - AU_n)(f)$. We have

$$g = (id - AU_n)(f) = (id - U_n)(f) + ((id - A)U_n)(f). \quad (10)$$

As in a) we have that (7) holds, that is we have

$$(id - U_n)(f) = \sum_{i=1}^{\gamma^-(n)} \phi_{ni}, \quad (11)$$

where the sets $V_{ni} = \text{supp}(\phi_{ni})$, $i = 1, \dots, \gamma^-(n)$, are all the local minimum sets of f of size n and satisfy (2). Therefore

$$g = \sum_{i=1}^{\gamma^-(n)} \phi_{ni} + ((id - A)U_n)(f). \quad (12)$$

Furthermore,

$$U_n(f)(x) = \begin{cases} f(x) & \text{if } x \in \mathbb{Z}^d \setminus \bigcup_{i=1}^{\gamma^-(n)} V_{ni} \\ v_i & \text{if } x \in V_{ni} \cup \{y_{ni}\}, i = 1, \dots, \gamma^-(n), \end{cases}$$

where $v_i = f(y_{ni}) = \min_{y \in \text{adj}(V_{ni})} f(y)$. Using that A is fully trend preserving, for every $i = 1, \dots, \gamma^-(n)$ there exists w_i such that $((id - A)U_n)(f)(x) = w_i$, $x \in V_{ni} \cup \{y_{ni}\}$. Moreover, using that every adjacent point has a neighbor in V_{ni} we have that $\min_{y \in \text{adj}(V_{ni})} ((id - A)U_n)(f)(y) = w_i$. Considering that the value of the pulse ϕ_{ni} is negative, we obtain through the representation (12) that V_{ni} , $i = 1, \dots, \gamma^-(n)$, are local minimum sets of g .

Next we show that g does not have any other local minimum sets of size n or less. Indeed, assume that V_0 is a local minimum set of g such that $\text{card}(V_0) \leq n$. Since

$V_0 \cup \text{adj}(V_0) \subset \mathbb{Z}^d \setminus \bigcup_{i=1}^{\gamma^-(n)} V_{ni}$ it follows from (12) that V_0 is a local minimum set of $((id - A)U_n)(f)$. Then using that $(id - A)$ is neighbor trend preserving and using [4, Theorem 17] we obtain that there exists a local minimum set W_0 of $U_n(f)$ such that $W_0 \subseteq V_0$. Then applying again [4, Theorem 17] or [4, Theorem 12] we obtain that there exists a local minimum set \tilde{W}_0 of f such that $\tilde{W}_0 \subseteq W_0 \subseteq V_0$. This inclusion implies that $\text{card}(\tilde{W}_0) \leq n$. Given that f does not have local minimum sets of size

less than n we have $\text{card}(\tilde{W}_0) = n$, that is \tilde{W}_0 is one of the sets V_{ni} - a contradiction. Therefore, V_{ni} , $i = 1, \dots, \gamma^-(n)$, are all the local minimum sets of g of size n or less. Then using again (7) we have

$$(id - U_n)(g) = \sum_{i=1}^{\gamma^-(n)} \phi_{ni} \quad (13)$$

Using (11) and (13) we obtain

$$(id - U_n)(g) = (id - U_n)(f)$$

Therefore

$$\begin{aligned} (U_n(id - AU_n))(f) &= U_n(g) = g - (id - U_n)(f) \\ &= (id - AU_n)(f) - (id - U_n)(f) \\ &= (U_n - AU_n)(f). \end{aligned}$$

This proves the first identity. The second one is proved in a similar manner. ■

References

- [1] R Anguelov and I N Plaskitt (now Fabris-Rotelli), A Class of LULU Operators on Multidimensional arrays, University of Pretoria, Tech. Rep. UPWT2007/22, 2008, URL=<http://arxiv.org/abs/0712.2923>.
- [2] R Anguelov and I N Fabris-Rotelli, Discrete Pulse Transform of Images, Proceedings of ICISP 2008: International Conference on Image and Signal Processing, Cherbourg-Octeville, Normandy, France, 1-3 July 2008, Lecture Notes in Computer Science, **5099** (2008) 1-9.
- [3] I Fabris-Rotelli and S J van der Walt, The Discrete Pulse Transform in Two Dimensions, Proceedings of the Twentieth Annual Symposium of the Pattern Recognition Association of South Africa, 30 November - 1 December 2009 Stellenbosch, South Africa.
- [4] R Anguelov and I N Fabris-Rotelli, LULU Operators and Discrete Pulse Transform for Multi-dimensional Arrays, IEEE Transactions on Image Processing, to appear, 2010.