

# Matching Lab

INFO/STSCI/ILRST 3900: Causal Inference

15 Oct 2025


# Reminders and Announcements

- ▶ Peer reviews- make sure to review all you're assigned by tomorrow, Oct 16
- ▶ In-class quiz 3 tomorrow, Oct 16
- ▶ Project Part 1 due Monday, Oct 20
- ▶ Office hours:
  - ▶ Filippo: Thursday 4-5 pm in 321A CIS Building
  - ▶ Shira: Monday 5-6 pm in 329A CIS Building
  - ▶ Sam: Tuesday 4-5 pm, in 350 CIS Building
- ▶ Check Ed for announcements and use for HW help!

# Matching Review

- ▶ Suppose person  $i$  is in the treatment group ( $A_i = 1$ ).
- ▶ Want to compare their outcome under treatment vs control
- ▶ Fundamental problem of causal inference: I can only observe one of these
- ▶ Matching: Find a person  $j$  in the control group ( $A_j = 0$ ) that is *similar enough* to person  $i$  and compare their outcomes
- ▶ Reasoning: if people are *similar enough*, then maybe their potential outcomes are also *similar enough*
- ▶ How do we define *similar enough*?
- ▶ We can use covariates!  $\vec{L}$


What if  $\vec{L}$  is multivariate?




- ▶ Conditional exchangeability holds when conditioning on Age and Education!
- ▶ Matching: look for a group of untreated units which has a similar distribution of Age and Education to the treated group

What if  $\vec{L}$  is multivariate?


What if  $\vec{L}$  is multivariate?



What if  $\vec{L}$  is multivariate?




What if  $\vec{L}$  is multivariate?



Which untreated unit should be the match?




What if  $\vec{L}$  is multivariate?




Which untreated unit should be the match?

What if  $\vec{L}$  is multivariate?




Which untreated unit should be the match?

What if  $\vec{L}$  is multivariate?




Which untreated unit should be the match?

What if  $\vec{L}$  is multivariate? We need a **distance metric**




- Define a way to measure “distance” between two individuals as a single number

What if  $\vec{L}$  is multivariate? We need a **distance metric**




- ▶ Define a way to measure “distance” between two individuals as a single number
- ▶ Match individuals using that distance!

What if  $\vec{L}$  is multivariate? We need a **distance metric**



What if  $\vec{L}$  is multivariate? We need a **distance metric**



► Manhattan distance:


- $d(\text{Treated}, \text{Untreated 1}) = 3 + 4 = 7$
- $d(\text{Treated}, \text{Untreated 2}) = 6 + 0 = 6 \checkmark$

► Euclidean distance:  $d(i, j) = \sqrt{\sum_p (L_{pi} - L_{pj})^2}$

- $d(\text{Treated}, \text{Untreated 1}) = \sqrt{3^2 + 4^2} = 5 \checkmark$
- $d(\text{Treated}, \text{Untreated 2}) = \sqrt{6^2 + 0^2} = 6$

► Which individual to pick depends on the distance metric!

What if  $\vec{L}$  is multivariate? We need a **distance metric**



- ▶ Manhattan distance:  $d(i, j) = \sum_p |L_{pi} - L_{pj}|$ 
  - ▶  $d(\text{Treated}, \text{Untreated 1}) = 3 + 4 = 7$
  - ▶  $d(\text{Treated}, \text{Untreated 2}) = 6 + 0 = 6 \checkmark$
- ▶ Euclidean distance:  $d(i, j) = \sqrt{\sum_p (L_{pi} - L_{pj})^2}$ 
  - ▶  $d(\text{Treated}, \text{Untreated 1}) = \sqrt{3^2 + 4^2} = 5 \checkmark$
  - ▶  $d(\text{Treated}, \text{Untreated 2}) = \sqrt{6^2 + 0^2} = 6$
- ▶ Which individual to pick depends on the distance metric!



# A common distance metric: Mahalanobis distance

Motivated by two principles

- ▶ Principle 1: Address unequal variances
  - ▶ Age might range uniformly from 18 to 80
  - ▶ Education range uniformly from 0 to 16
  - ▶ We might correct for this so age doesn't dominate the distance

# A common distance metric: Mahalanobis distance

Motivated by two principles

- ▶ Principle 1: Address unequal variances
  - ▶ Age might range uniformly from 18 to 80
  - ▶ Education range uniformly from 0 to 16
  - ▶ We might correct for this so age doesn't dominate the distance
- ▶ Principle 2: Address correlations
  - ▶ Suppose we included age in years, age in months, and education
  - ▶ Suppose we included age in years and age in months are very correlated
  - ▶ We should care about a correlation-corrected distance

# Code

Let's try this out in R!

- ▶ Section 2 is worked out for you: read through, run the code blocks, and answer the questions
- ▶ Section 3 asks you to write some code (will be very similar to the code from Section 2)
- ▶ Then move on to the `matching_examples.Rmd` file on the website