

MARKOV CHAIN MONTE CARLO I

PROBABILISTIC DECISION MAKING VU

(REINFORCEMENT LEARNING VO)

Thomas Wedenig

Dec 10, 2025

Institute of Machine Learning and Neural Computation
Graz University of Technology, Austria

1. Motivation
2. Markov Chains
3. Metropolis-Hastings
4. Continuous State Spaces
5. Estimating Expectations with MCMC Samples

MOTIVATION

- Let π be a **probability distribution** on state space \mathcal{X}
 - Discrete \mathcal{X} : probability mass function $\pi : \mathcal{X} \rightarrow [0, 1]$
 - Continuous \mathcal{X} : probability density function $\pi : \mathcal{X} \rightarrow \mathbb{R}_{\geq 0}$
- Assume π is either
 - **known**, i.e., we can evaluate $\pi(\mathbf{x}) \forall \mathbf{x} \in \mathcal{X}$
 - **known up to normalization**, i.e., we can evaluate $\tilde{\pi}(\mathbf{x}) \propto \pi(\mathbf{x}) \forall \mathbf{x} \in \mathcal{X}$

Goal: Draw i.i.d. samples from π .

- In practice, it's **very common** to only know the **unnormalized** distribution $\tilde{\pi}$:

$$\pi(\mathbf{x}) = \frac{\tilde{\pi}(\mathbf{x})}{Z}, \quad Z := \int_{\mathcal{X}} \tilde{\pi}(\mathbf{x}) \, d\mathbf{x}$$

Bayesian Inference

- Prior $p(\mathbf{x})$
- Likelihood $p(\mathbf{y} \mid \mathbf{x})$
- Posterior:

$$p(\mathbf{x} \mid \mathbf{y}) = \frac{p(\mathbf{y} \mid \mathbf{x}) p(\mathbf{x})}{\int_{\mathcal{X}} p(\mathbf{y} \mid \mathbf{x}') p(\mathbf{x}') \, d\mathbf{x}'} =: \pi_{\mathbf{y}}(\mathbf{x})$$

- Hence, $\tilde{\pi}_{\mathbf{y}}(\mathbf{x}) = p(\mathbf{y} \mid \mathbf{x}) p(\mathbf{x}) \propto p(\mathbf{x} \mid \mathbf{y})$

Boltzmann Distributions

- Given an **energy function** $E : \mathcal{X} \rightarrow \mathbb{R}$
 - E can be **learned from data**
 - Sometimes, E is **known** (e.g., physics)
- $E(\mathbf{x})$ is small for likely \mathbf{x} :

$$\pi(\mathbf{x}) := \frac{\exp(-E(\mathbf{x}))}{\int_{\mathcal{X}} \exp(-E(\mathbf{x}')) \, d\mathbf{x}'}$$


Why do we care about **samples** from π in the first place? 🤔

Generative Models

- In **Generative Modeling**, **samples** are the quantities of interest !
- Generate novel molecules, images, text, audio, video ...



<https://www.nytimes.com/2023/04/08/technology/ai-photos-pope-francis.html>



Hoogeboom et al., 2023, <https://arxiv.org/pdf/2203.17003>

Why do we care about **samples** from π in the first place? 🤔


Monte Carlo Estimation

- Often, we care about $\mathbb{E}_{X \sim \pi}[f(X)]$
 - Mean $\mathbb{E}_{X \sim \pi}[X]$, variance $\mathbb{E}_{X \sim \pi}[(X - \mathbb{E}_{X' \sim \pi}[X'])^2]$, higher order moments, ...
- Importantly, in Bayesian methods, we seek the **Posterior Predictive**:

$$p(y_* | \mathbf{x}_*, \mathcal{D}) = \int \underbrace{p(y_* | \mathbf{x}_*, \boldsymbol{\theta})}_{\text{Likelihood}} \underbrace{p(\boldsymbol{\theta} | \mathcal{D})}_{\text{Posterior}} d\boldsymbol{\theta} = \mathbb{E}_{p(\boldsymbol{\theta} | \mathcal{D})}[p(y_* | \mathbf{x}_*, \boldsymbol{\theta})]$$

- Given i.i.d. samples, we could **estimate all these quantities** with Monte Carlo !

MARKOV CHAINS




- A Markov Chain (MC) consists of n RVs X_i with state space \mathcal{X}

- The joint factorizes as

$$p(x_1, x_2, \dots, x_n) = p(x_1) p(x_2 | x_1) p(x_3 | x_2) \dots p(x_n | x_{n-1})$$

- Markov property: The next state x_{t+1} only depends on the current state x_t
 - The history x_1, \dots, x_{t-1} becomes irrelevant
- Ingredients: initial distribution $p(x_1)$, transition distributions $p(x_t | x_{t-1})$




$$p(x_1, x_2, \dots, x_n) = p(x_1) p(x_2 | x_1) p(x_3 | x_2) \dots p(x_n | x_{n-1})$$

- Assume **finite state space** $\mathcal{X} := \{1, \dots, s\}$
- Let's think about the **marginal distributions** $p_t(x_t)$:

$$p_1(x_1), \quad p_2(x_2) = \sum_{x_1} p(x_2 | x_1) p_1(x_1), \quad p_3(x_3) = \sum_{x_2} p(x_3 | x_2) p_2(x_2),$$

$$\dots \quad p_n(x_n) = \sum_{x_{n-1}} p(x_n | x_{n-1}) p_{n-1}(x_{n-1})$$



$$p(x_1, x_2, \dots, x_n) = p(x_1) p(x_2 | x_1) p(x_3 | x_2) \dots p(x_n | x_{n-1})$$

How do we sample from $p(x_1, \dots, x_n)$? 🤔


- Ancestral sampling:

$$x_1 \sim p_1(x_1), \quad x_2 \sim p(x_2 | x_1), \quad x_3 \sim p(x_3 | x_2), \quad \dots \quad x_n \sim p(x_n | x_{n-1})$$

- The distribution of x_t is $p_t(x_t)$

Goal of MCMC (Informal)

Given π (or $\tilde{\pi}$), construct Markov Chain such that $p_t \approx \pi$ for large t .




- All considered chains will be **time homogeneous**:
- Transition $p(X_t = i \mid X_{t-1} = j)$ will **not depend** on t , only on i, j
- Let's collect them in a **matrix $P \in \mathbb{R}^{s \times s}$** :

$$P_{ij} := p(X_t = i \mid X_{t-1} = j)$$

i.e., the columns sum to 1.

- We can write the **marginal mass function** as a vector **$\mu_t \in \mathbb{R}^s$** :

$$\mu_t := (p_t(X_t = 1), \dots, p_t(X_t = s))^T$$



$$P_{ij} := p(X_t = i \mid X_{t-1} = j)$$

- For all t , we now have:

$$\mu_{t+1} = P\mu_t$$


- Note that

$$\mu_2 = P\mu_1, \quad \mu_3 = P\mu_2 = P(P\mu_1) = P^2\mu_1$$

where $P^2 = PP$ denotes repeated matrix multiplication

- In general, for all t :


$$\mu_{t+1} = P^t\mu_1 \quad (P^t)_{ij} = p(X_t = i \mid X_1 = j)$$



<https://ermongroup.github.io/cs228-notes/inference/sampling/>


- $\mathcal{X} = \{1, 2, 3\}$
- Transition probability matrix:

$$P = \begin{bmatrix} 0 & 0.5 & 1 \\ 1 & 0 & 0 \\ 0 & 0.5 & 0 \end{bmatrix}, \quad P_{ij} = p(X_t = i \mid X_{t-1} = j)$$



<https://ermongroup.github.io/cs228-notes/inference/sampling/>

- We pick $\mu_1 = (1/3, 1/3, 1/3)^\top$ and iteratively compute $\mu_t = P\mu_{t-1}$



- **Observation 1:** When $\mu = (0.4, 0.4, 0.2)^\top$, we had $\mu = P\mu$ 😲
- Such a μ is called a **stationary distribution** of the Markov Chain !
 - i.e., if $X_t \sim \mu$, then $X_{t+1} \sim \mu$
- **Observation 2:** We actually **converged** to this μ when starting with our μ_1 🤯
- In this Markov Chain, it turns out that
 - The **stationary distribution** is unique
 - We will converge to it for **any initial distribution** μ_1 😍

Goal of MCMC in finite state spaces (Less Informal)

Given $\pi \in \mathbb{R}^S$ (or $\tilde{\pi}$), construct P such that $\lim_{t \rightarrow \infty} \mu_t = \pi$ for any μ_1 .

- For this, π needs to be the **unique stationary distribution**

How do we ensure there exists a unique stationary distribution ?

Irreducibility

A finite-state Markov Chain is **irreducible** if

$$\forall i, j \exists t \geq 1 : (P^t)_{ij} > 0.$$

i.e., it's possible to go from any state i to any state j in finite time.


Unique stationary distribution

A finite-state Markov Chain has a **unique stationary distribution**


$$\mu = P\mu$$

if and only if it is irreducible.


Are the following Markov Chains **irreducible** ?




<https://ermongroup.github.io/cs228-notes/inference/sampling/>




<https://ermongroup.github.io/cs228-notes/inference/sampling/>





$$P = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad P_{ij} = p(X_t = i \mid X_{t-1} = j)$$

- Since it's irreducible, there exists a unique stationary distribution μ
 - Namely $\mu = (0.5, 0.5)^\top$
- Let's again compute marginal distributions μ_t , starting with $\mu_1 = (1, 0)^\top$



How do we ensure we also **converge to the stationary distribution** ?

Period

Given a finite-state Markov Chain, the **period** of state i is

$$d(i) := \gcd \left(\{t \geq 1 : (P^t)_{ii} > 0\} \right).$$

Aperiodicity


A finite-state Markov Chain is said to be **aperiodic** if $d(i) = 1$ for all i .

Convergence to stationary distribution


An **irreducible and aperiodic** finite-state Markov Chain **converges** to its unique stationary distribution μ :

$$\lim_{t \rightarrow \infty} \mu_t = \mu$$

for any initial distribution μ_1 .



<https://ermongroup.github.io/cs228-notes/inference/sampling/>



- **Important:** We write down P and (many) μ_t here for **pedagogical** reasons
 - In practice (e.g. huge s or \mathcal{X} uncountable), this is totally **impossible**
- When **sampling from the chain**, this is **not needed** !
- To sample from μ_n
 1. Sample x_1 from the **initial distribution**
 2. Repeat: Sample $x_t \sim p(X_t | X_{t-1} = x_{t-1})$ for $t = 2, 3, \dots, n$
 3. Return x_n

- So we wish to design P such that it's **irreducible** and **aperiodic**
- But how do we pick P such that the stationary distribution is π ? 🤔
- It is **nasty** to directly solve for P in

$$\pi = P\pi$$

- Instead, many MCMC algorithms design chains that satisfy **detailed balance**

Detailed Balance

A **sufficient** (*but not necessary*) condition for $\pi = P\pi$ to hold is called **detailed balance**:

$$\forall i, j : P_{ij} \pi_j = P_{ji} \pi_i$$

i.e., the amount of mass transported from $j \rightarrow i$ is the same as from $i \rightarrow j$ at stationarity.

Proof.

Assume P satisfies detailed balance at π . Then, for all i :

$$(P\pi)_i = \sum_j P_{ij} \pi_j \stackrel{\text{DB}}{=} \sum_j P_{ji} \pi_i = \pi_i \underbrace{\sum_j P_{ji}}_{=1} = \pi_i$$

and hence, $\pi = P\pi$.



$$\forall i, j: P_{ij} \pi_j = P_{ji} \pi_i$$

$$P = \begin{bmatrix} \frac{7}{12} & \frac{3}{8} & \frac{1}{2} \\ \frac{1}{4} & \frac{1}{2} & \frac{1}{4} \\ \frac{1}{6} & \frac{1}{8} & \frac{1}{4} \end{bmatrix}, \quad \boldsymbol{\pi} = \left(\frac{1}{2} \quad \frac{1}{3} \quad \frac{1}{6} \right)^{\top}, \quad \boldsymbol{\pi} = P\boldsymbol{\pi}$$

Probability Mass Flow 1 \leftrightarrow 2

- Flow 1 \rightarrow 2: $\frac{1}{4} \cdot \frac{1}{2} = \frac{1}{8}$
- Flow 2 \rightarrow 1: $\frac{3}{8} \cdot \frac{1}{3} = \frac{1}{8}$

Probability Mass Flow 1 \leftrightarrow 3

- Flow 1 \rightarrow 3: $\frac{1}{6} \cdot \frac{1}{2} = \frac{1}{12}$
- Flow 3 \rightarrow 1: $\frac{1}{2} \cdot \frac{1}{6} = \frac{1}{12}$

Probability Mass Flow 2 \leftrightarrow 3

- Flow 2 \rightarrow 3: $\frac{1}{8} \cdot \frac{1}{3} = \frac{1}{24}$
- Flow 3 \rightarrow 2: $\frac{1}{4} \cdot \frac{1}{6} = \frac{1}{24}$

METROPOLIS-HASTINGS

Markov Chain Monte Carlo (MCMC)

Any algorithm that **constructs a Markov Chain**

- that has the target π as the **unique stationary distribution**,
- **converges** to π as $t \rightarrow \infty$ (for any initial distribution),
- and **draws samples** from that Markov Chain (“**Monte Carlo Simulation**”)

is called a **Markov Chain Monte Carlo (MCMC)** algorithm.

- We will now look at the **canonical** MCMC algorithm:
- The **Metropolis-Hastings**¹ algorithm

¹Authorship controversial. Mainly developed by Arianna and Marshall Rosenbluth, while Hastings generalized it.

- For now, again consider finite state space $\mathcal{X} := \{1, \dots, s\}$
- We wish to construct an **irreducible** and **aperiodic** Markov Chain
- We will use **detailed balance** to enforce that π is the **stationary distribution**
- Of course, the chain should be **easy to sample from**
- **General idea:**
 - When in state X_t , we **propose** a new point \hat{X}_{t+1} by **sampling from a proposal distribution**
 - We will **accept \hat{X}_{t+1}** with a certain probability and **reject** it else, i.e.,

$$X_{t+1} = \begin{cases} \hat{X}_{t+1} & \text{with probability } \alpha \\ X_t & \text{with probability } 1 - \alpha \end{cases}$$

- Consider a **proposal matrix** $Q \in \mathbb{R}^{s \times s}$ where

$$Q_{ij} = q(\hat{X}_{t+1} = i \mid X_t = j)$$

where q is the **proposal distribution**

- We'll **pick a q** that is **easy to sample from** and induces an **irreducible chain**
 - e.g., q could be the uniform distribution over states “close” to X_t
- When picking $P = Q$, the stationary distribution will in general **not** be π !

- Goal: π should be the stationary distribution
- We introduce an acceptance probability $\alpha(j \rightarrow i) \in [0, 1]$
- For $i \neq j$, we set

$$P_{ij} := Q_{ij} \alpha(j \rightarrow i)$$

- To ensure the columns of P sum to 1, we set the diagonal to

$$P_{jj} := 1 - \sum_{i \neq j} P_{ij}$$

- Let's pick $\alpha(j \rightarrow i)$ such that P satisfies **detailed balance** at π , i.e., for all i, j :

$$P_{ij} \pi_j = P_{ji} \pi_i$$

- For $i \neq j$, substitute $P_{ij} = Q_{ij} \alpha(j \rightarrow i)$:

$$Q_{ij} \alpha(j \rightarrow i) \pi_j = Q_{ji} \alpha(i \rightarrow j) \pi_i$$

- Rearranging yields

$$\frac{\alpha(j \rightarrow i)}{\alpha(i \rightarrow j)} = \frac{Q_{ji} \pi_i}{Q_{ij} \pi_j}$$

- There are **infinitely** many choices of $0 \leq \alpha(j \rightarrow i), \alpha(i \rightarrow j) \leq 1$ that satisfy this
- The **Metropolis-Hastings (MH)** algorithm **picks**

$$\alpha(j \rightarrow i) = \min \left(1, \frac{Q_{ji} \pi_i}{Q_{ij} \pi_j} \right)$$

which is the **largest acceptance rate** allowed by detailed balance

$$\alpha(j \rightarrow i) = \min \left(1, \frac{Q_{ji} \pi_i}{Q_{ij} \pi_j} \right)$$

- **Note:** We can compute $\alpha(j \rightarrow i)$ even when we only have access to $\tilde{\pi}$ 🥰

$$\frac{Q_{ji} \pi_i}{Q_{ij} \pi_j} = \frac{Q_{ji} \frac{\tilde{\pi}_i}{Z}}{Q_{ij} \frac{\tilde{\pi}_j}{Z}} = \frac{Q_{ji} \tilde{\pi}_i}{Q_{ij} \tilde{\pi}_j}$$

Sample X_1 from initial distribution p_1 . Repeat for $t = 1, \dots, n$:

1. Given current state $X_t = j$, propose:

$$\hat{X}_{t+1} \sim q(\hat{X}_{t+1} \mid X_t = j)$$

2. Let $\hat{X}_{t+1} = i$. Compute acceptance probability:

$$\alpha(j \rightarrow i) = \min \left(1, \frac{Q_{ji} \tilde{\pi}_i}{Q_{ij} \tilde{\pi}_j} \right)$$

3. Accept or reject:

$$\text{Sample } U \sim \text{Uniform}(0, 1), \quad X_{t+1} = \begin{cases} \hat{X}_{t+1} & \text{if } U \leq \alpha(j \rightarrow i) \\ X_t & \text{else} \end{cases}$$

- For $i \neq j$, detailed balance holds **by construction**, and for $i = j$ it is **trivially true**
- When Q is **irreducible** (on the support of π), P is **as well**
- P is **aperiodic** because for some j , we have $P_{jj} > 0^2$ since either
 - we pick $Q_{jj} > 0$ for some j , so sometimes we propose to stay in state j , or
 - $\alpha(j \rightarrow i) < 1$, i.e., we sometimes reject the proposal and stay in state j
- Thus, π is the **unique stationary distribution**, and we will **converge to it** as $t \rightarrow \infty$

²This implies that an irreducible chain is aperiodic.

METROPOLIS-HASTINGS MCMC DEMO

CONTINUOUS STATE SPACES

So far, the state space \mathcal{X} was **finite**. What about **continuous \mathcal{X}** ? 🤔

- The target π is now assumed to be a **probability measure**
 - In practice, this will almost always admit a **density** w.r.t. Lebesgue
- Instead of $P_{ij} = \mathbb{P}(X_{t+1} = i \mid X_t = j)$, we now have a **transition kernel**

$$K(A \mid \mathbf{x}) := \mathbb{P}(X_{t+1} \in A \mid X_t = \mathbf{x})$$

which is a **probability measure** for every $\mathbf{x} \in \mathcal{X}$

- Instead of evolving $\mu_{t+1} = P\mu_t$, we **integrate against K** :

$$\mu_{t+1}(A) = \int_{\mathcal{X}} K(A \mid \mathbf{x}) \mu_t(d\mathbf{x})$$

- **Stationarity** meant $\pi = P\pi$, but now its

$$\pi(A) = \int_{\mathcal{X}} K(A \mid \mathbf{x}) \pi(d\mathbf{x})$$

- Irreducibility, aperiodicity, and **detailed balance** are defined slightly differently
 - Conceptually the same idea, just with **measures** instead of **mass functions**
- For cont. \mathcal{X} , there are irreducible & aperiodic MCs **without stationary distributions**
 - We need an **additional condition** called **positive Harris recurrence**³
 - It makes sure the chain returns infinitely often to **sets A with $\pi(A) > 0$**
 - Intuitively, we must not “wander off to infinity”
- Irreducible, aperiodic and **positive Harris recurrent** chains are called **ergodic** !
 - This again guarantees a unique stationary distribution and convergence to it

³Finite-state irreducible and aperiodic chains automatically fulfill this.

- **Metropolis-Hastings:** Proposal Q_{ij} becomes **density** $q(\mathbf{y} \mid \mathbf{x})$ and

$$\alpha(\mathbf{x} \rightarrow \mathbf{y}) = \min \left(1, \frac{q(\mathbf{x} \mid \mathbf{y}) \tilde{\pi}(\mathbf{y})}{q(\mathbf{y} \mid \mathbf{x}) \tilde{\pi}(\mathbf{x})} \right)$$

where $\tilde{\pi}$ is the **target's unnormalized density**

- For example, **Gaussian proposal**

$$q(\mathbf{y} \mid \mathbf{x}) := \mathcal{N}(\mathbf{y}; \mathbf{x}, \sigma^2 I)$$

- For this choice, we have **symmetry** $q(\mathbf{y} \mid \mathbf{x}) = q(\mathbf{x} \mid \mathbf{y})$ and thus,

$$\alpha(\mathbf{x} \rightarrow \mathbf{y}) = \min \left(1, \frac{\tilde{\pi}(\mathbf{y})}{\tilde{\pi}(\mathbf{x})} \right)$$

CONTINUOUS STATE SPACE MCMC DEMO

<https://chi-feng.github.io/mcmc-demo/app.html>

ESTIMATING EXPECTATIONS WITH MCMC SAMPLES

Recall Goal: Draw i.i.d. samples from π .

Are samples from our Markov Chain i.i.d. samples from π ? 🤔

No !

- If samples were i.i.d., X_t and X_{t+1} would be **independent**
- In the beginning, X_t is not even **marginally** distributed like π (not converged yet)
- Even when the chain has perfectly converged to π , and thus $X_t \sim \pi$, $X_{t+1} \sim \pi$ **marginally**, we still have **possibly large correlation** between X_t and X_{t+1}
- In the finite-state setting:

$$p(X_t = j, X_{t+1} = i) = p(X_t = j)p(X_{t+1} = i \mid X_t = j) = \pi_j P_{ij} \neq \pi_j \pi_i = p(X_t = j)p(X_{t+1} = i)$$

Are we doomed? Monte Carlo estimators need **i.i.d.** samples 🥵

Thankfully, no! 🥳

Ergodic Theorem

Assume X_1, X_2, \dots is an **ergodic Markov Chain** with stationary distribution π . Then,

$$\hat{\mathbb{E}}_T := \frac{1}{T} \sum_{t=1}^T f(X_t) \xrightarrow{\text{a.s.}} \mathbb{E}_{X \sim \pi}[f(X)]$$

as $T \rightarrow \infty$ for every initial state $X_1 = x$.

- **Almost sure (a.s.) convergence** means

$$\mathbb{P} \left(\lim_{T \rightarrow \infty} \hat{\mathbb{E}}_T = \mathbb{E}_{X \sim \pi}[f(X)] \right) = 1$$

- This is the **MCMC counterpart** to classic i.i.d. Monte Carlo

Recap: Properties of MC Estimate

Given **i.i.d.** samples $Y_1, \dots, Y_T \sim \pi$, the **Monte Carlo estimate** of $\mathbb{E}_{X \sim \pi} [f(X)]$ is $\hat{\mathbb{E}}_T^{(MC)} := \frac{1}{T} \sum_{t=1}^T f(Y_t)$.

- **Unbiased** for every T : $\mathbb{E} \left[\hat{\mathbb{E}}_T^{(MC)} \right] = \mathbb{E}_{X \sim \pi} [f(X)]$

Properties of MCMC Estimate

- MCMC estimate is generally⁴ **biased** for any finite T
 - We can reduce bias by **discarding the first B samples (burn-in phase)**
- Bias goes to 0 as $T \rightarrow \infty$, i.e., **asymptotically unbiased**

⁴It would be unbiased if the initial distribution was already π , but we can't do that in practice.

Recap: Properties of MC Estimate

- **Variance** decreases **linearly** with T : $\text{var} \left[\hat{\mathbb{E}}_T^{(\text{MC})} \right] = \frac{\text{var}[f(X)]}{T}$

Properties of MCMC Estimate

- As $T \rightarrow \infty$, the **variance** of the MCMC estimator is approximately

$$\text{var} \left[\hat{\mathbb{E}}_T \right] \approx \frac{\sigma_{\text{as}}^2}{T}$$

with

$$\sigma_{\text{as}}^2 = \text{var}_{\pi}[f(X)] + 2 \sum_{n=1}^{\infty} \text{cov}_{\pi}(f(X_0), f(X_n)).$$

where $X_0 \sim \pi, X_1 \sim \pi, \dots$ (i.e., chain in **stationary** regime)

- **Note:** σ_{as}^2 typically **larger** than $\text{var}_{\pi}[f(X)]$, but still decreases **linearly** with T

- **Goal:** Sample from **complicated, high-dimensional distribution** π
 - Where typically only $\tilde{\pi}$ is known
- **Idea:** Construct Markov Chain that
 - Has **unique stationary distribution** π
 - Marginal distribution **converges to it** when $T \rightarrow \infty$
 - The transition distribution is **easy to sample from**
- **Sampling** from the chain for “long enough” yields
 - Samples with **marginal distribution close to** π
 - That are still **correlated** (even as $T \rightarrow \infty$)
 - But that **can still be used to estimate expectations strongly consistently**

Next week, you will hear about...

- **More practical, widely used MCMC algorithms**
 - Gibbs's Sampling
 - Langevin Dynamics
 - Hamiltonian Monte Carlo
- **Convergence & Diagnostics**
 - Burn-in Phase
 - Autocorrelation
 - Effective Sample Size