

Advanced Mathematics

Valeri Makarov

Dept. of Mathematical Analysis and Applied Mathematics, U.C.M.

Grado en Ingeniería Informática (grupo I)

F.CC. Matemáticas, Desp. 420

<http://www.mat.ucm.es/~vmakarov>

e-mail: vmakarov@mat.ucm.es

Chapter 2


Fourier series and Fourier transform

Periodic functions

Let's consider a function (or as we shall call it a **signal**)

$$s(t) = A \cos(\omega t + \phi)$$

blue : $A = 1, \omega = 2\pi\frac{1}{4}, \phi = 0$ **red** : $A = 0.6, \omega = 2\pi\frac{1}{2}, \phi = 2$



Def.: A signal $s(t)$ is **periodic** iff $\exists T > 0$ s.t. $s(t + T) = s(t)$.

Note: If $s(t + T) = s(t)$ then $s(t + 2T) = s(t)$ etc.

Def.: $T_0 = \min\{T\}$ is called the period of $s(t)$

In the example above: for the blue curve $T = 4$ s; for red $T = 2$ s.
Thus, the blue function (signal) makes one cycle per each 4 s.

The function $s(t) = A \cos\left(2\pi \frac{t}{T} + \phi\right)$ has period T .

Def.: $f = \frac{1}{T}$ (1/s) is called the frequency of $s(t)$. It is measured in Hertz


$$1 \text{ Hz} = \frac{1}{1 \text{ s}}$$

For example 4 Hz means 4 cycles per second.

Def.: The magnitude $\varphi(t) = 2\pi ft + \phi$ measured in radians is called the **phase or angle of the signal**.

Def.: The value $\omega = 2\pi f$ measured in [rad/s] is called the **angular velocity of the signal**.

Given two signals $s_1(t) = \cos(\omega t)$ and $s_2(t) = \cos(\omega t + \phi)$, we can define the phase shift ϕ between them.



The **time delay** τ can be calculated by:

$$s_1(t) = s_2(t-\tau) = \cos(\omega t) = \cos(\omega(t-\tau) + \phi) \Rightarrow \omega\tau = \phi \Rightarrow \tau = \frac{\phi}{\omega}$$

Def.: Given a signal $s(t) = A \cos(\omega t + \phi)$ the magnitude A is called the **amplitude** of the signal.

Problem 2.1a Find the period of $\sin(ax)$

By definition:

$$\sin(a(x+T)) = \sin(ax+aT) = [\text{if } aT = 2\pi] = \sin(ax+2\pi) = \sin(ax)$$

Thus, $T = \frac{2\pi}{a}$. From the other side we know $\omega = a = 2\pi f = \frac{2\pi}{T}$, which gives the same result.

Problem 2.1c Find the period of e^{iax}

We have

$$e^{iax} = e^{iax+2\pi i} = e^{ia(x+2\pi/a)} \Rightarrow T = \frac{2\pi}{a}$$

Orthogonal functions

Def.: Two functions $f(x)$ and $g(x)$ are called orthogonal if their scalar product is equal to zero:

$$\langle f, g \rangle = \int_a^b f^*(x)g(x) dx = 0$$

Problem 2.2a Show that $\{\cos nx, \sin nx\}$ ($n \geq 0$) are orthogonal in $[-\pi, \pi]$

$$\int_{-\pi}^{\pi} \cos nx \sin nx dx = -\frac{1}{n} \int_{-\pi}^{\pi} \cos nx d \cos nx = -\frac{\cos^2 nx}{2n} \Big|_{-\pi}^{\pi} = 0$$

Problem 2.3 Prove

$$\int_{-T/2}^{T/2} f(t) dt = \int_0^T f(t) dt = \int_{\alpha}^{\alpha+T} f(t) dt$$

$$\int_{-T/2}^{T/2} f(t) dt = \int_{-T/2}^0 f(t) dt + \int_0^{T/2} f(t) dt = [u = t - T] =$$

$$= \int_{T/2}^T f(u + T) du + \int_0^{T/2} f(t) dt = \int_0^T f(t) dt$$

$$\int_{\alpha}^{T+\alpha} f(t) dt = \int_{\alpha}^0 f(t) dt + \int_0^T f(t) dt + \int_T^{T+\alpha} f(t) dt$$

The last integral ($u = t - T$):

$$\int_T^{T+\alpha} f(t) dt = \int_0^{\alpha} f(u + T) du = - \int_{\alpha}^0 f(t) dt$$

Fourier series

Let's consider a signal $s(t)$ such that $s(t + 2\pi) = s(t)$ (2π periodic). In 1807 Jean-Baptiste Joseph Fourier proposed that

$$s(t) = \sum_{n=0}^{\infty} (a_n \cos(nt) + b_n \sin(nt))$$

where a_n, b_n are some real numbers. Later it was called the Fourier series.

This series allows representing $s(t)$ as a composition of simple sin and cosine functions.

The series must converge. Thus, for $n \gg 1$, a_n and b_n should be small. Therefore, $s(t) \approx \sum_{n=0}^N (a_n \cos(nt) + b_n \sin(nt))$, if N is big enough.

Calculation of a_n and b_n

Let's start from the definition and integrate it (assuming that the series converges uniformly):

$$\int_{-\pi}^{\pi} s(t) dt = \sum_{n=0}^{\infty} \int_{-\pi}^{\pi} (a_n \cos(nt) + b_n \sin(nt)) dt$$

Using Problem 2.2 we get the first coefficient:

$$\int_{-\pi}^{\pi} s(t) dt = 2\pi a_0 \Rightarrow a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} s(t) dt$$

Now we can multiply by $\cos(kt)$ and integrate

$$\int_{-\pi}^{\pi} s(t) \cos(kt) dt = \sum_{n=0}^{\infty} \int_{-\pi}^{\pi} \cos(kt) (a_n \cos(nt) + b_n \sin(nt)) dt$$

Due to the function orthogonality, only for $k = n$ we have nonzero term:

$$\int_{-\pi}^{\pi} s(t) \cos(nt) dt = a_n \int_{-\pi}^{\pi} \cos^2(nt) dt = a_n \int_{-\pi}^{\pi} \frac{1 + \cos(2nt)}{2} dt = \pi$$

Thus,

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} s(t) \cos(nt) dt \quad n > 0$$

We can do the same with sin:

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} s(t) \sin(nt) dt \quad n \geq 0$$

Note, $b_0 = 0$. Then we have: Given a signal $s : [-\pi, \pi] \rightarrow \mathbb{R}$ (could be not periodic) its Fourier series is

$$s(t) = a_0 + \sum_{n=1}^{\infty} (a_n \cos(nt) + b_n \sin(nt))$$

where a_n and b_n are calculated by the equations given above.

Problem 2.5a Find Fourier series for $s(t) = |t|$ on $[-\pi, \pi]$.

We note that this is an even function. Thus, $b_n = 0$. Let's find a_n :

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} |t| dt = \frac{1}{\pi} \int_0^{\pi} t dt = \frac{1}{\pi} \frac{\pi^2}{2} = \frac{\pi}{2}$$

$$\begin{aligned} a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} |t| \cos(nt) dt = \frac{2}{\pi} \int_0^{\pi} t \cos(nt) dt = [\text{by parts}] = \\ &= \frac{2}{\pi n} \left[t \sin(nt) \Big|_0^{\pi} - \int_0^{\pi} \sin(nt) dt \right] = \frac{2}{\pi n^2} \cos(nt) \Big|_0^{\pi} = \frac{2[(-1)^n - 1]}{\pi n^2} = \\ &= -\frac{4}{\pi n^2} \begin{cases} 0, & \text{if } n = 2k \\ 1, & \text{if } n = 2k + 1 \end{cases} \end{aligned}$$

Matlab script for Problem 2.5a





```
%% Fourier Series for  $s(t) = |t|$  on  $[-\pi, \pi]$ 
t = linspace(-pi,pi,200); % time
s = abs(t);                % signal

a0 = pi/2; f0 = ones(1,200); % constant term

k = (1:5)';
a = -4./(pi*(2*k-1).^2); % values for  $a_n$ 
e = cos((2*k-1)*t);      %  $\cos(nt)$  functions

% Make drawing
figure('color','w','position',[100 100 800 500])
sp = a0*f0;
for k = 1:4
    subplot(2,2,k)
    plot(t,s,'m','LineWidth',2)
    hold on
    sp = sp + a(k)*f(k,:);
end
```

Graphic representation for Problem 2.5a



Fourier series: General case

Let $s(t)$ be defined on the interval $[0, T]$. **The Fourier series:**

$$s(t) = a_0 + \sum_{n=1}^{\infty} \left(a_n \cos \left(\frac{2\pi n}{T} t \right) + b_n \sin \left(\frac{2\pi n}{T} t \right) \right)$$

$$a_0 = \frac{1}{T} \int_0^T s(t) dt, \quad a_n = \frac{2}{T} \int_0^T s(t) \cos \left(\frac{2\pi n}{T} t \right) dt$$

$$b_n = \frac{2}{T} \int_0^T s(t) \sin \left(\frac{2\pi n}{T} t \right) dt$$

Theorem (Convergence). If a Fourier series of a function $s(t)$ with period T converges uniformly, then the series converges to $s(t)$.

Convergence of Fourier series

Theorem: Let $s : \mathbb{R} \rightarrow \mathbb{R}$, $s(t + T) = s(t)$ be a piecewise continuous (finite number of step-type discontinuities are allowed). If for $t \in [0, T]$ there exist lateral derivatives $s'(t^-)$ and $s'(t^+)$, then the Fourier series converges pointwise to

$$\frac{s(t^+) + s(t^-)}{2}$$

Note: If $s(t)$ is derivable, then its Fourier series converges pointwise to $s(t)$. Indeed, in this case we have

$$(s(t^+) + s(t^-))/2 = s(t).$$

Parseval's equality

If $s \in L_2$ is 2π periodic and continuous, then

$$\frac{1}{\pi} \int_{-\pi}^{\pi} s^2(t) dt = 2a_0^2 + \sum_{n=1}^{\infty} (a_n^2 + b_n^2)$$

(formally it can be obtained by squaring the formula and integrating using the orthogonality).

Problem 2.8a: Use the Fourier series of x^2 on $[-\pi, \pi]$ to find $\sum_{n=1}^{\infty} \frac{1}{n^2}$.

1. Fourier series. The function is even, therefore $b_n = 0$.

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} x^2 dx = \frac{x^3}{6\pi} \Big|_{-\pi}^{\pi} = \frac{\pi^2}{3}$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 \cos(nx) dx = [\text{two times by parts}] = \frac{4(-1)^n}{n^2}$$

Thus

$$x^2 = \frac{\pi^2}{3} + 4 \sum_{n=1}^{\infty} \frac{(-1)^n \cos(nx)}{n^2}$$

2. If $x = 0$ we have

$$0 = \frac{\pi^2}{3} + 4 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \Rightarrow \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} = -\frac{\pi^2}{12}$$

3. If $x = \pi$ we have $\cos(\pi n) = (-1)^n$ and then

$$\pi^2 = \frac{\pi^2}{3} + 4 \sum_{n=1}^{\infty} \frac{(-1)^n (-1)^n}{n^2} \Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$$

4. Let's apply the Parseval's equality:

$$\frac{1}{\pi} \int_{-\pi}^{\pi} x^4 dx = \frac{2\pi^4}{5} = 2a_0^2 + \sum_{n=1}^{\infty} a_n^2 = \frac{2\pi^4}{9} + 16 \sum_{n=1}^{\infty} \frac{1}{n^4}$$

Thus

$$\sum_{n=1}^{\infty} \frac{1}{n^4} = \frac{\pi^4}{90}$$

$$\sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2} = \sum_{k=0}^{\infty} \frac{(-1)^{2k}}{(2k+1)^2} - \sum_{k=1}^{\infty} \frac{(-1)^{2k}}{(2k)^2} \Rightarrow \sum_{k=0}^{\infty} \frac{1}{(2k+1)^2} = \frac{\pi^2}{8}$$