



CARINGBAH HIGH SCHOOL

2024 HIGHER SCHOOL CERTIFICATE TRIAL EXAMINATION

Mathematics Advanced

- Reading time – 10 minutes
- Working time – 3 hours
- Write using black pen.
- Calculators approved by NESA may be used.
- Reference sheet is provided.
- For questions in Section II, show relevant mathematical reasoning and/or calculations.

Section I – 10 marks (pages 3 - 7)

- Attempt Questions 1–10
- Allow about 15 minutes for this section.
- Answer the questions on page 40.

Section II – 90 marks (pages 10 – 34)

- Attempt Questions 11–32
- Allow about 2 hours and 45 minutes for this section.

Student Number: _____ **Class:** _____

Marker's Use Only								
Section I	Section II							Total
Q1-10	11 - 15	16 - 18	19 - 22	23 - 25	26- 28	29 - 30	31-32	
/10	/12	/13	/16	/13	/11	/10	/15	/100

Section I


10 marks

Attempt Question 1-10

Allow about 15 minutes for this section.

Use the multiple-choice answer sheet for Questions 1-10.

- 1 The graph of a relation is shown.



Which type of relationship does the graph represent?

- (A) one-to-one
 - (B) one-to-many
 - (C) many-to-one
 - (D) many-to-many
- 2 Consider the graph $y = 4x + 1$

What is the equation of the graph after it has been translated 2 units to the right?

- (A) $y = 2x - 7$
- (B) $y = 2x + 9$
- (C) $y = 4x - 7$
- (D) $y = 4x + 9$

3 Which of the following is the domain of the function $y = \frac{1}{\sqrt{x^2 - 1}}$?

(A) $(-\infty, -1] \cup [1, \infty)$

(B) $(-\infty, -1) \cup (1, \infty)$

(C) $(-1, 1)$

(D) $[-1, 1]$

4 Families in Caringbah were surveyed about the types of pets they own. The results of the survey are shown below.

- 40% of families own a dog.
- 36% of families own a cat.
- 18% of families own both a cat and a dog.

A family is chosen at random and found to own a cat.

What is the probability that the family also owns a dog?

(A) 18%

(B) 36%

(C) 40%

(D) 50%

5 A supermarket assistant sets up a display of lemonade cans. The display forms a triangle, where each row has one fewer can than the one below. There are 15 cans at the base of the display and one can at the top of the display.

What is the total number of cans used in the display?


(A) 30

(B) 120

(C) 160

(D) 180

- 6 The diagram shows a circle with radius r cm and a sector with an internal angle of $\frac{\pi}{6}$ radians.



NOT TO SCALE


If the area of the shaded segment is 1.7 cm^2 , which of the following is closest to the radius of the circle?

- (A) 0.12 cm
 - (B) 0.34 cm
 - (C) 12.00 cm
 - (D) 144.08 cm
- 7 Alicia and Steven are participating in a fitness program. The probability that Alicia will complete the program is 0.8, and the probability that Steven will complete the program is 0.7.

What is the probability that only one of them will successfully complete the program?

- (A) 0.14
- (B) 0.24
- (C) 0.38
- (D) 0.56

- 8 The graph of $y = x\ln(x^2)$ is shown.



How many solutions does the equation $|x\ln(x^2)| = \frac{1}{3}(x + 1)$ have?


- (A) 3
- (B) 4
- (C) 5
- (D) 6
- 9 The following cumulative frequency table shows heights of students at a school.

Height (cm)	Cumulative Frequency
131 – 140	70
141 – 150	210
151 – 160	420
161 – 170	680
171 – 180	800

What is the height of a student in the 7th decile?

- (A) 131 – 140 cm
- (B) 151 – 160 cm
- (C) 161 – 170 cm
- (D) 171 – 180 cm

- 10 The graph shows two cubic functions, $y = f(x)$ and $y = g(x)$.



It is given that $f(x)$ is an odd function and that $g(x) = f(-x)$.

Define $h(x) = f(x) - g(x)$.

How many stationary points does $y = h(x)$ have?

- (A) 0
- (B) 1
- (C) 2
- (D) 3

End of Section I

Question 11

Differentiate $y = xe^{7x}$, expressing your answer in factorised form.

2

Question 12

Find the exact gradient of the tangent to the curve $y = e^{2x+3}$ at the point where $x = 0$.

2

Question 13

Solve the equation $\frac{\operatorname{cosec} x}{5} - 2 = 0$ for $0^\circ \leq x \leq 360^\circ$, correct to the nearest minute.

3

Question 14

A bag contains six cards that are each printed with one of the following numbers.

111, 121, 122, 211, 221, 222

A student chooses one card at random.

Let N_1 be the event that the first digit of the number on the card is 1, and N_2 be the event that the second digit of the number on the card is 1.

Determine whether N_1 and N_2 are independent events.

3

Question 15


Evaluate $\int_0^{\frac{\pi}{4}} \cos(2x) \, dx$

2

Question 16

Darren and Sylvia visit Sydney Harbour on New Year's Eve to watch the fireworks.

Darren is standing outside the Sydney Opera House and Sylvia is on a yacht in the harbour. They can both see the fireworks at the top of the Sydney Harbour Bridge.




From Darren's position (D), the Sydney Harbour Bridge (B) is at a bearing of 78° , and the angle of elevation to the top of the bridge (T) is 35° .

From Sylvia's position (S), the Sydney Harbour Bridge is at a bearing of 140° , and the angle of elevation to the top of the bridge is 48° .

The top of the bridge is 50 m above the base of the bridge.

- a) Using appropriate diagrams and reasoning, show that $\angle SBD = 62^\circ$.

2



Question 16 continues on page 13

b) Hence, find the distance between Sylvia and Darren, correct to 2 decimal places.

3

This image shows a full page of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page, typical of notebook or legal stationery. There are no margins, text, or other markings on the page.

End of Question 16

3

[illegible]

The velocity of two racing cars, P and Q , are shown on the graph below.



The velocity of car P at time t seconds is given by the function $v_P = 3t^2 - 6t + 4$, where v_P is the velocity in metres per second.

The velocity of car Q is accelerating at a constant rate. Both cars start at the same point and have the same velocity at times where $t = 0$ and $t = 4$.

- a) Show that the equation for the velocity of car Q , v_Q , is given by $v_Q = 6t + 4$.

2

- b) Both cars start the race from the same point.

3

Find the earliest time when car P will pass car Q after the race starts.

End of Question 18

Question 19

a) Find the derivative of $y = \tan(x^2)$.

1

b) Hence, or otherwise, find $\int x \tan^2(x^2) dx$.

3

Question 20

Use two applications of the trapezoidal rule to find an approximate value of $\int_1^2 \ln x dx$.

2

Give your answer correct to 2 decimal places.

Question 21

a) Show that $\frac{1}{\sec\theta - \tan\theta} - \frac{1}{\sec\theta + \tan\theta} = 2\tan\theta$

2

b) Hence or otherwise, find $\int \frac{1}{\sec\theta - \tan\theta} - \frac{1}{\sec\theta + \tan\theta} d\theta$

2

Question 22

A continuous random variable has the probability density function $f(x)$ given by:

$$f(x) = \begin{cases} \frac{kx}{5-x^2} & 0 \leq x \leq 2 \\ 0 & \text{for all other values of } x \end{cases}$$

- a) Show that the value of k is $\frac{2}{\ln 5}$.

2

This image shows a full page of white paper with horizontal dashed lines, typical of primary school handwriting practice paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.


Question 22 continues on page 19

c) Hence, or otherwise, show that the median is $\sqrt{5 - \sqrt{5}}$.

2

Question 23

A factory produces boxes in the shape of an open rectangular prism (without a lid).



The length, height and width of the box are l cm, x cm and $2x$ cm respectively.

The factory uses $k \text{ cm}^2$ of sheet metal to make the box, where k is a constant.

a) Show that $l = \frac{k - 4x^2}{4x}$.

1

Question 23 continues on page 21

b) Hence, find the maximum volume of the box that can be produced from a sheet of metal with an area of 1200 cm^2 .

3

This image shows a full page of a worksheet designed for handwriting practice. It features 20 evenly spaced, horizontal dashed lines across the entire width of the page. The background is plain white, and there are no margins, text, or other markings present.

End of Question 23

APPROXIMATELY HALFWAY – 55 marks out of 100 complete at this point.

Question 24


Consider the series $\ln x - 3(\ln x)^2 + 9(\ln x)^3 - 27(\ln x)^4 + \dots$

- a) Show that this series is a geometric series.

2

- b) On the set of axes below, sketch the graphs of $y = \ln x$, $y = -\frac{1}{3}$ and $y = \frac{1}{3}$, showing all points of intersection and intercepts with the coordinate axes.

2



Question 24 continues on page 23

c) Hence, or otherwise, find the values of x for which the series has a limiting sum.

2

Question 25


The graph $y = \sin x$ is vertically dilated by a scale factor of 2 and horizontally dilated such that it has a period of $\frac{2\pi}{3}$.

Find the x -coordinates of the points of intersection of the transformed sine curve and the line $y = \sqrt{2}$ for $x \in [0, 2\pi]$.

3

Question 26

For a sample of fifteen high school students, their age in years, x , and the length of their right foot in centimetres, y , were recorded.



The graph shows the data as well as a regression line which passes through (12, 22.5) and (19, 27.5).

- a) Find the equation of the regression line in the form $y = mx + c$.

2

Question 26 continues on page 25

b) Layton has a five-year old brother whose right foot is 13 cm in length.

He extends the line of best fit and notes that it predicts that a five-year old should have a foot length of approximately 17.5 cm.

Give TWO reasons why Layton is incorrect to assume from the bivariate data that his brother has smaller than normal feet for a five-year old.

2

Question 27

The random variable X has this probability distribution.

X	16	17	18	19	20
$P(X = x)$	0.2	0.4	0.1	0.2	0.1

1


a) Find the expected value.

2

b) Find the standard deviation, correct to 2 decimal places. Show working to justify your answer.

Question 28

The diagram shows the function $y = x^3$ and the line $y = b$



NOT TO
SCALE

- a) By solving an appropriate equation, show that the coordinates of A are $(\sqrt[3]{b}, b)$.

1

- b) By finding the area between the curve $y = x^3$, the line $y = b$ and the y -axis, or otherwise, find the value of b such that the shaded area is $\frac{4}{27}$ unit².

3

Question 29

An astronomer is attempting to predict the height of a satellite above the ground as it orbits around the Earth. Using precise measurements, the height of the satellite can be modelled with the function:

$$h(t) = 6\sin\left(\frac{\pi}{6}(t-15)\right) + 8$$


where $h(t)$ is the height of the satellite in thousands of kilometres at time t hours since midnight on any day of the year.

- a) Find the minimum and maximum height reached by the satellite.

1

- b) On the grid below, sketch the graph of $h(t)$ for $0 \leq t \leq 24$.

2



Question 28 continue on page 29


- c) The satellite is out of range if it is higher than 11 000 km above the ground. Use algebraic techniques to find how many hours in the day the satellite is out of range.

[illegible]

End of Question 29

Question 30

The graph of a cumulative distribution function is shown below.



The curved section of the graph is part of the function $y = \frac{(x-1)^3}{6}$

- a) Show that the value of A is $\sqrt[3]{6} + 1$.

2

- b) Find the probability density function for the given cumulative distribution function, including any restrictions on the domain.

2

Question 31

Consider the gradient function $f'(x) = 3(x + 3)(x - 1)$.

- a) The graph of $y = f(x)$ passes through the point $(2, -8)$. Show that $f(x) = x^3 + 3x^2 - 9x - 10$.

3

- b) Find the coordinates of the minimum and maximum values in the interval $-4 \leq x \leq 4$.

2

You DO NOT need to determine the nature of the stationary points.


Question 31 continues on page 32

c) Show that a point of inflection exists and find its coordinates.

2

d) Sketch the graph of $y = f(x)$ in the interval $-4 \leq x \leq 4$, showing the locations of the endpoints, the stationary points and the inflection point.

2



End of Question 31

Question 32

An electric vehicle with an empty battery is being recharged. The capacity, $C\%$, of the battery while charging at time t minutes may be modelled by the equation:

$$C = 100(1 - 2^{-kt})$$

The battery is charged at 35% capacity after 50 minutes.

- a) Show that the value of k is 0.01243, correct to 4 significant figures.

2

- b) To prolong the life of the battery, the charger may be set to switch off when the battery reaches 90% capacity. For how long will the battery be on charge until it reaches 90% capacity?

2

Question 32 continue on page 34

- c) By considering the first and second derivative, describe the behaviour of C for $t \geq 0$ if the battery is left on charge indefinitely.

2

This image shows a full page of white paper with horizontal dashed lines, typical of primary school handwriting practice paper. The lines are evenly spaced and run across the entire width of the page. There are no margins, text, or other markings present.

End of Examination!!! 😊