

Moduli Stabilisation and the Statistics of SUSY Breaking in the Landscape

Igor Bröckel

Summer Series on String Phenomenology
15.09.2020



Istituto Nazionale di Fisica Nucleare

Moduli Stabilisation and the Statistics of SUSY Breaking in the Landscape

Igor Broeckel,^a Michele Cicoli,^a Anshuman Maharana,^b Kajal Singh^b and Kuver Sinha^c

^a*Dipartimento di Fisica e Astronomia, Università di Bologna, via Irnerio 46, 40126 Bologna, Italy
and INFN, Sezione di Bologna, viale Berti Pichat 6/2, 40127 Bologna, Italy*

^b*Harish-Chandra Research Institute, HBNI, Jhunsi, Allahabad, UP 211019, India*

^c*Department of Physics and Astronomy, University of Oklahoma, Norman, OK 73019, USA*

E-mail: igor.broeckel@bo.infn.it, michele.cicoli@unibo.it,
anshumanmaharana@hri.res.in, kajalsingh@hri.res.in,
kuver.sinha@ou.edu

arXiv:2007.04327

Content

- 1. Review of Statistical Approach**
- 2. Importance of the Kähler moduli**
- 3. Stabilisation mechanism**
 - 3.1 LVS models**
 - 3.2 KKLT models**
 - 3.3 Perturbatively stabilised models**
- 4. SUSY breaking statistics**
- 5. Phenomenological Implications**
- 6. Conclusion**

Review of Statistical Approach

- SUSY is a central idea in Pheno and Theory (Hierarchy probl., DM candidates, etc.)
- Can String Theory give guidance in the search for SUSY?
- Landscape is large, no vacuum is preferred (yet), many vacua at least roughly match SM → Statistical analysis
- First studies found a preference for high scale SUSY, due to a uniform distribution of SUSY breaking scale
[Douglas, 04], [Denef,Douglas, 04], [Denef,Douglas, 05]
- These studies focused on the dilaton and complex structure F-terms and neglected the Kähler moduli F-terms, since these fields are stabilized beyond tree-level → only sub-leading correction?
- Based on dynamical SUSY breaking arguments a logarithmic behavior of the SUSY breaking scale was also expected (BUT: for KKLT)
[Dine,Gorbatov,Thomas, 04],[Dine, 05],[Dine,O'Neil,Sun, 05],[Dine, 04]

→ **What is the origin for the power-law / logarithmic scaling?**

Importance of the Kähler moduli

- Short summary of the results of D.D.

$$K_{\text{tree}} = -2 \ln \mathcal{V} - \ln (S + \bar{S}) - \ln \left(-i \int_X \Omega(U) \wedge \bar{\Omega}(\bar{U}) \right), \quad W_{\text{tree}} = \int_X G_3 \wedge \Omega(U)$$

$$\rightarrow V_{\text{tree}} = |F^S|^2 + |F^U|^2 + |F^T|^2 - 3m_{3/2}^2 \approx |F^S|^2 + |F^U|^2 - 3m_{3/2}^2$$

- Where the gravitino mass is given by: $m_{3/2} = e^{K/2} |W|$
 - Kähler moduli not stabilised at tree-level \rightarrow only a small correction to leading order?
 - Distribution of SUSY breaking vacua was assumed to be:
- $$dN_{\Lambda=0}(F) = \prod d^2 F^S d^2 F^U d\hat{\Lambda} \rho(F, \hat{\Lambda}) \delta \left(|F^S|^2 + |F^U|^2 - \hat{\Lambda} \right)$$
- Assumptions: Several hidden sectors, vanishing cosmological constant, uniform distribution of axion-dilaton and complex structure

$$\rightarrow \boxed{dN_{\Lambda=0}(m_{3/2}) \sim \rho(m_{3/2}) m_{3/2} dm_{3/2}} \quad \boxed{\rho(m_{3/2}) \sim m_{3/2}^\beta, \quad \beta = 0} \quad [\text{Douglas, 04}]$$

Importance of the Kähler moduli

- **BUT:** Using the ‘no-scale’ relation we can rewrite the scalar potential as

$$V_{\text{tree}} = |F^S|^2 + |F^U|^2 + m_{3/2}^2 \left(K_{\bar{T}} K^{\bar{T}T} K_T - 3 \right) = \frac{e^{K_{cs}}}{\mathcal{V}^2 (S + \bar{S})} (|D_S W|^2 + |D_U W|^2)$$

- any vacuum with $D_i W \neq 0$ is unstable since it gives rise to a run-away for the volume mode. Hence a stable solution requires $F^S = F^U = 0$
- at tree-level the gravitino mass is set by the F-terms of the T-moduli since ‘no-scale’ implies $|F^T|^2 = 3m_{3/2}^2$
- soft terms are of order $m_{3/2}$ only for matter located on D7 branes, not for D3. For instance, gaugino masses for D3’s are set by F^S , which is non-zero due to sub-leading corrections beyond tree-level. In order to determine F^S one needs to stabilise the Kähler moduli

[Jockers, 05]

→ **SUSY statistics should be driven by the Kähler moduli**

Stabilisation mechanism - KKLT

- Purely non-perturbative stabilisation: $W = W_0 + Ae^{-aT}$
[Kachru,Kalosh,Linde,Trivedi, 03]
 - Here the Kähler modulus is $T = \tau + i\Theta$ and $a = 2\pi/\mathfrak{n}$ is a parameter that determines the nature of the non-perturbative effect.
 - Minimizing the scalar potential leads to: $e^{a\langle\tau\rangle} \sim \frac{2Aa\langle\tau\rangle}{3W_0} \Leftrightarrow \langle\tau\rangle \sim \frac{1}{a} |\ln W_0|$
 - The gravitino mass at the minimum is:
$$m_{3/2} = \frac{\pi g_s^{1/2}}{\mathfrak{n}^{3/2}} \frac{|W_0| M_p}{|\ln W_0|^{3/2}}$$
- In order to be able to neglect stringy corrections to the effective action and pert. corrections to K one needs: $W_0 \ll 1$

→ **the gravitino mass in KKLT is mainly driven by W_0**

Stabilisation mechanism - LVS

- Perturbative and non-perturbative stabilisation:
[Balasubramanian, Berglund, Conlon, Quevedo, 05]
[Cicoli, Conlon, Quevedo, 08]
 - perturbative: $K = -2 \ln \left(\mathcal{V} + \frac{\xi}{2} \left(\frac{S + \bar{S}}{2} \right)^{3/2} \right) + \dots$
 - non-perturbative: $W = W_0 + A_s e^{-a_s T_s}$
- Minimizing the scalar potential leads to: $\langle \mathcal{V} \rangle \sim \frac{3\sqrt{\langle \tau_s \rangle} |W_0|}{4a_s A_s} e^{a_s \langle \tau_s \rangle}, \quad \langle \tau_s \rangle \sim \frac{1}{g_s} \left(\frac{\xi}{2} \right)^{2/3}$
- The gravitino mass at the minimum is: $m_{3/2} \sim c_1 \frac{g_s M_p}{\mathfrak{n}} e^{-\frac{c_2}{g_s \mathfrak{n}}}$
- Where c_1 and c_2 are numerical coefficients
 - the gravitino mass in LVS is mainly driven by g_s

Stabilisation mechanism – perturbative

- Purely perturbative stabilisation:
[Berg, Haack, Kors, 06]

$$K_{g_s^0 \alpha'^3} = -\frac{\xi}{g_s^{3/2} \mathcal{V}}, \quad K_{g_s^2 \alpha'^2} = g_s \frac{b(U)}{\mathcal{V}^{2/3}}, \quad K_{g_s^2 \alpha'^4} = \frac{c(U)}{\mathcal{V}^{4/3}}.$$

- The functions $b(U), c(U)$ are known explicitly only for simple toroidal orientifolds but are expected to be $\mathcal{O}(1 - 10)$
- Minimizing the scalar potential leads to: $\langle \mathcal{V} \rangle \sim 26 g_s^{9/2} \left(\frac{c(U)}{|\xi|} \right)^3$
- The gravitino mass at the minimum is: $m_{3/2} \sim \lambda \frac{|W_0| M_p}{g_s^4 c(U)^3}$
- Consistency of the stabilisation requires $\langle \mathcal{V} \rangle \gg 1, g_s \ll 1$

→ the gravitino mass in pert. stabilisation is mainly driven by $c(U)$

SUSY breaking statistics

- Gravitino mass is mainly determined by W_0 , g_s , n , $c(U)$

→ The distribution of $|W_0|^2$ as a complex variable is assumed to be uniform:
[Douglas, 04]

$$dN \sim |W_0| d|W_0|$$

→ The distribution of g_s was checked to be uniform for rigid CY. And was shown to hold in more general cases:

[Shok,Douglas, 04][Denef,Douglas, 04]

[Blanco-Pillado,Sousa,Urkiola,Wachter, 20]

$$dN \sim dg_s$$

→ The distribution of the rank of the condensing gauge group is still poorly understood. We expect the number of states N to decrease when n increases, since D7-tadpole cancellation is more difficult to satisfy

$$dN \sim -n^{-r} dn$$

→ Since $c(U)$ is a function of the complex structure, large values are considered as fine tuned

$$dN \sim -c^{-k} dc$$

SUSY breaking statistics - LVS

- Using the scaling of the underlying parameters, we can compute the scaling behavior of the gravitino in LVS:

$$\rightarrow dm_{3/2} \sim n m_{3/2} \left[\ln \left(\frac{M_p}{m_{3/2}} \right) \right]^2 \left[1 - \frac{c_2 n^{r-2}}{\ln \left(\frac{M_p}{m_{3/2}} \right)} \right] dN$$

- For any value of the exponent r the leading order result is given by

$$\rightarrow \boxed{\rho_{\text{LVS}}(m_{3/2}) \sim \frac{1}{n m_{3/2}^2} \left[\ln \left(\frac{M_p}{m_{3/2}} \right) \right]^{-2}} \quad \boxed{N_{\text{LVS}} \sim \ln \left(\frac{m_{3/2}}{M_p} \right)}$$

- In LVS we have: $m_{3/2} \sim M_{\text{soft}}^{1/p}$, where the value of p depends on the specific model (D3, D7, sequestered)

→ LVS vacua feature a logarithmic distribution of soft terms

SUSY breaking statistics - KKLT

- Using the scaling of the underlying parameters, we can compute the scaling behavior of the gravitino in KKLT:

$$dm_{3/2} \sim \frac{M_p^2}{m_{3/2}} \left[\frac{g_s}{\mathbf{n}^3 |\ln W_0|^3} + \frac{m_{3/2}^2}{2M_p^2} \left(\frac{1}{g_s} + 3\mathbf{n}^{r-1} \right) \right] dN$$

- For any value of the exponent r the leading order result is given by

$$\rightarrow \rho_{\text{KKLT}}(m_{3/2}) \sim \frac{1}{M_p^2} \left(\frac{\mathbf{n}^3 |\ln W_0|^3}{g_s} \right) \sim \text{const.}$$

$$N_{\text{KKLT}} \sim \left(\frac{m_{3/2}}{M_p} \right)$$

- In KKLT we have: $m_{3/2} \sim M_{\text{soft}}$

→ KKLT vacua feature a power-law distribution of soft terms

SUSY breaking statistics - perturbative

- Using the scaling of the underlying parameters, we can compute the scaling behavior of the gravitino in pert. stabilisation:

$$dm_{3/2} \sim m_{3/2} \left(3c^{k-1} - \frac{4}{g_s} \right) dN$$

- Control over the effective field theory requires $k > 1$

$$\rightarrow \boxed{\rho_{\text{PERT}}(m_{3/2}) \sim \frac{1}{M_p^2} \left(\frac{m_{3/2}}{M_p} \right)^{\frac{k-7}{3}}} \quad \boxed{N_{\text{PERT}} \sim \left(\frac{m_{3/2}}{M_p} \right)^{\frac{k-1}{3}}}$$

- Qualitatively similar to KKLT (equal for $k=7$)
- Soft masses are expected to behave as in LVS

→ **pert. stabilised vacua feature a power-law distribution of soft terms**

Phenomenological Implications

- We have found a draw towards high scale SUSY → reason for no SUSY at LHC?
- Problem with high scale SUSY → fine tuning for the Higgs-mass
- However, in LVS models a logarithmic distribution makes low-energy SUSY appear less tuned
- Quantifying fine-tuning: Barbieri-Giudice measure $\Delta_{BG} \equiv \max_i \left| \frac{\partial \ln m_Z^2}{\partial \ln p_i} \right|$
[Barbieri, Giudice, 88]
→ 10% fine-tuning for most superpartners at TeV scale
- Introducing fine-tuning penalties like anthropic arguments would set a bound on the mass of the Z boson → bound on scale of superpartners
- Introducing cosmological constraints

Conclusion

- We have stressed that Kähler moduli stabilisation is a critical requirement for a proper treatment of the statistics of SUSY breaking
- Different no-scale breaking effects used to fix the Kähler moduli lead to a different dependence of $m_{3/2}$ on the flux dependent microscopic parameters
- In LVS models the distribution of the gravitino mass and soft terms are **logarithmic**
- In KKLT and perturbative stabilisation the distribution are **power-law**
- Determining which distribution is more representative of the structure of the flux landscape translates into the question of which vacua are more frequent, LVS or KKLT?
- LVS needs less tuning → larger parameter space → LVS models favoured?
- Definite answer requires more detailed studies