



Paul-Konstantin Oehlmann

Uppsala Universitet

Based on

- [arXiv:1910.04095](#) with: Nadir Hajouji
- [arXiv:2005.12929](#) with: Markus Dierigl and Fabian Ruehle

Summer Series on String Phenomenology
June 23rd 2020

What is a field theory with $\pi_1(G) \neq 1$?

Example: The group $SU(2)$ admits a \mathbb{Z}_2 center and reps carry charges

Matter	<i>Rep</i>	1	2	3	4	...
--------	------------	----------	----------	----------	----------	-----

What is a field theory with $\pi_1(G) \neq 1$?

Example: The group $SU(2)$ admits a \mathbb{Z}_2 center and reps carry charges

Matter	<i>Rep</i>	1	2	3	4	...
	q_{cent}	0	1	0	1	...

What is a field theory with $\pi_1(G) \neq 1$?

Example: The group $SU(2)$ admits a \mathbb{Z}_2 center and reps carry charges

Matter	Rep	1	2	3	4	...
	q_{cent}	0✓	1✗	0✓	1✗	...

- Matter with $q_{cent} = 0$ only: electric 1-form symmetry [Gaiotto, Kapustin, Seiberg, Willet '14]
- **Gauging** this symmetry one obtains $SU(2)/\mathbb{Z}_2 \sim SO(3)$
- Gauging impacts the **spectrum** of Wilson Line and t'Hooft surface operators [Aharony, Seiberg, Tachikawa '13]

What is a field theory with $\pi_1(G) \neq 1$?

Example: The group $SU(2)$ admits a \mathbb{Z}_2 center and reps carry charges

Matter	Rep	1	2	3	4	...
	q_{cent}	0✓	1✗	0✓	1✗	...

- Matter with $q_{cent} = 0$ only: electric 1-form symmetry [Gaiotto, Kapustin, Seiberg, Willet '14]
- **Gauging** this symmetry one obtains $SU(2)/\mathbb{Z}_2 \sim SO(3)$
- Gauging impacts the **spectrum** of Wilson Line and t'Hooft surface operators [Aharony, Seiberg, Tachikawa '13]

Lie Group	Center
$SU(p)$	\mathbb{Z}_p
$SO(4n)$	$\mathbb{Z}_2 \times \mathbb{Z}_2$
$SO(4n+2)$	\mathbb{Z}_4
E_6	\mathbb{Z}_3
E_7	\mathbb{Z}_2
E_8	\mathbb{Z}_1

What is a field theory with $\pi_1(G) \neq 1$?

Example: The group $SU(2)$ admits a \mathbb{Z}_2 center and reps carry charges

Matter	Rep	1	2	3	4	...
	q_{cent}	0✓	1✗	0✓	1✗	...

- Matter with $q_{cent} = 0$ only: electric 1-form symmetry [Gaiotto, Kapustin, Seiberg, Willet '14]
- **Gauging** this symmetry one obtains $SU(2)/\mathbb{Z}_2 \sim SO(3)$
- Gauging impacts the **spectrum** of Wilson Line and t'Hooft surface operators [Aharony, Seiberg, Tachikawa '13]

Lie Group	Center
$SU(p)$	\mathbb{Z}_p
$SO(4n)$	$\mathbb{Z}_2 \times \mathbb{Z}_2$
$SO(4n+2)$	\mathbb{Z}_4
E_6	\mathbb{Z}_3
E_7	\mathbb{Z}_2
E_8	\mathbb{Z}_1

← Can take arbitrary high quotient $SU(p)/\mathbb{Z}_p$?!


Landscape: There should be a cutoff!

- We want to **systematically** construct SQFTs and SUGRA theories with Gauge groups $G = G^*/T$
- **Flexible** (all sorts of G^*) and **consistent** using geometric arguments
- **Tool of choice is F-theory**: allows (especially in 6D) to construct a large patch of the string landscape and **relate swampland questions to geometry**

- We want to **systematically** construct SQFTs and SUGRA theories with Gauge groups $G = G^*/T$
- **Flexible** (all sorts of G^*) and **consistent** using geometric arguments
- **Tool of choice is F-theory**: allows (especially in 6D) to construct a large patch of the string landscape and **relate swampland questions to geometry**
- $G^* = ABCDE$ algebras geometrically realized by torus fiber singularities [Kodaira '68] ; $\pi_1(G)$ encoded in finite Mordell-Weil group [Aspinwall, Morrison '98; Mayrhofer, Morrison, Till, Weigand '14]
- **M-theory dual** for higher form symmetries in **lower dimensions** [Morrison, Schafer-Nameki, Willet, Albertini, del Zotto, Etxebarria, Hosseini '20]
- Other **fun geometric games**: (Fiberwise-mirror duality, smooth quotients. . .) [Klevers, Mayorga, O. Piragua, Reuter '14; O. Reuter, Schimannek; Cvetič, Klevers, Piragua, Poretschkin '16; Anderson, Gray, O. '19]


- We want to **systematically** construct SQFTs and SUGRA theories with Gauge groups $G = G^*/T$
- **Flexible** (all sorts of G^*) and **consistent** using geometric arguments
- **Tool of choice is F-theory**: allows (especially in 6D) to construct a large patch of the string landscape and **relate swampland questions to geometry**
- $G^* = ABCDE$ algebras geometrically realized by torus fiber singularities [Kodaira '68] ; $\pi_1(G)$ encoded in finite Mordell-Weil group [Aspinwall, Morrison '98; Mayrhofer, Morrison, Till, Weigand '14]
- **M-theory dual** for higher form symmetries in **lower dimensions** [Morrison, Schafer-Nameki, Willet, Albertini, del Zotto, Etxebarria, Hosseini '20]
- Other **fun geometric games**: (Fiberwise-mirror duality, smooth quotients. . .) [Klevers, Mayorga, O. Piragua, Reuter '14; O. Reuter, Schimannek; Cvetič, Klevers, Piragua, Poretschkin '16; Anderson, Gray, O. '19]
- **Question 1**: what $T = \pi_1(G)$ in a **SUGRA** are allowed?
- **Question 2**: What structure does an **SCFT** with $T = \pi_1(G_F)$ have?

① Geometrize via F-theory *(G always non-Abelian!)




② SUGRA: $G = G^*/T$
 (swampland) Question:
 which T are possible?

③ 6D SCFT: G_F/T
 SCFTs with modified
 global flavor structure




$[p,q]$ 7 Branes act as $SL(2, \mathbb{Z})$ defects on the IIB axio-dilaton τ




$[p,q]$ 7 Branes act as $SL(2, \mathbb{Z})$ defects on the IIB axio-dilaton τ

- A $[1,0]$ 7-brane acts on τ by a trafo $\mathcal{T} \in SL(2, \mathbb{Z})$
- A $[p,q]$ 7-branes engineered by combinations of $\mathcal{S}, \mathcal{T} \in SL(2, \mathbb{Z})$



Fix an additional point s_p of order p on the elliptic curve \mathcal{E}




Fix an additional point s_p of order p on the elliptic curve \mathcal{E}

- Second point s_p generates an addition law and a group on the elliptic curve
- If $s_p \oplus^{(p)} s_p = s_0$ the group is finite $MW(\mathcal{E}) = T \sim \mathbb{Z}_p$



MW Addition Law

Neutral Element




Fix an additional point s_p of order p on the elliptic curve \mathcal{E}

- Second point s_p generates an addition law and a group on the elliptic curve
- If $s_p \oplus^{(p)} s_p = s_0$ the group is finite $MW(\mathcal{E}) = T \sim \mathbb{Z}_p$
- If torsion s_p preserved over all B **construct projector** on states $\sigma(s_p)$


$$MW(\mathcal{E}) \hookrightarrow NS(Y_{res}), \quad [\text{Mayrhofer, Morrison, Till, Weigand '14}]$$

$\sigma(s_p)$ allows only compatible matter representations ✓




Fix an additional point s_p of order p on the elliptic curve \mathcal{E}

- Second point s_p generates an addition law and a group on the elliptic curve
- If $s_p \oplus^{(p)} s_p = s_0$ the group is finite $MW(\mathcal{E}) = T \sim \mathbb{Z}_p$
- **Different perspective:** which $SL(2, \mathbb{Z})$ elements **stabilize** the \mathbb{Z}_p torsion point




Fix an additional point s_p of order p on the elliptic curve \mathcal{E}

- Second point s_p generates an addition law and a group on the elliptic curve
- If $s_p \oplus^{(p)} s_p = s_0$ the group is finite $MW(\mathcal{E}) = T \sim \mathbb{Z}_p$
- **Different perspective:** which $SL(2, \mathbb{Z})$ elements **stabilize** the \mathbb{Z}_p torsion point



Fix an additional point s_p of order p on the elliptic curve \mathcal{E}

- Second point s_p generates an addition law and a group on the elliptic curve
- If $s_p \oplus^{(p)} s_p = s_0$ the group is finite $MW(\mathcal{E}) = T \sim \mathbb{Z}_p$
- **Different perspective:** which $SL(2, \mathbb{Z})$ elements **stabilize** the \mathbb{Z}_p torsion point



Fix an additional point s_p of order p on the elliptic curve \mathcal{E}

- Second point s_p generates an addition law and a group on the elliptic curve
- If $s_p \oplus^{(p)} s_p = s_0$ the group is finite $MW(\mathcal{E}) = T \sim \mathbb{Z}_p$
- **Different perspective:** which $SL(2, \mathbb{Z})$ elements **stabilize** the \mathbb{Z}_p torsion point
- This is the **congruence subgroup** $\Gamma_1(p) \in SL(2, \mathbb{Z})$

$$\Gamma_1(p) = \left\{ \gamma \in SL(2, \mathbb{Z}) : \gamma \equiv \begin{pmatrix} 1 & 0 \\ * & 1 \end{pmatrix} \pmod{p} \right\}. \quad (1)$$

Fiber	(f, g, Δ)	Monodromy	Subgroups	Group	center
II^*	$(\geq 1, 1, 2)$	$\begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$	-	-	-
I_n	$(0, 0, n)$	$\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$	$\Gamma_1(n), \Gamma(n)$	$SU(n)$	\mathbb{Z}_n
III	$(\geq 1, \geq 2, 3)$	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$	$\Gamma_1(2)$	$SU(2)$	\mathbb{Z}_2
IV	$(\geq 2, 2, 4)$	$\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$	$\Gamma_1(3)$	$SU(3)$	\mathbb{Z}_3
I_{2n}^*	$(\geq 2, 3, 6 + 2n)$	$\begin{pmatrix} -1 & -2n \\ 0 & -1 \end{pmatrix}$	$\Gamma_1(2), \Gamma(2)$	$SO(8 + 4n)$	$\mathbb{Z}_2 \times \mathbb{Z}_2$
I_{2n+1}^*	$(\geq 2, 3, 7 + 2n)$	$\begin{pmatrix} -1 & -2n-1 \\ 0 & -1 \end{pmatrix}$	$\Gamma_1(2), \Gamma_1(4)$	$SO(10 + 4n)$	\mathbb{Z}_4
IV^*	$(\geq 3, 4, 8)$	$\begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}$	$\Gamma_1(3)$	E_6	\mathbb{Z}_3
III^*	$(3, \geq 5, 9)$	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$	$\Gamma_1(2)$	E_7	\mathbb{Z}_2
II^*	$(4, \geq 5, 10)$	$\begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$	-	E_8	-



7 Brane monodromies consistent with center of the covering group ($\Gamma(p)$ fixes a $\mathbb{Z}_p \times \mathbb{Z}_p$ group)

Fiber	(f, g, Δ)	Monodromy	Subgroups	Group	center
II^*	$(\geq 1, 1, 2)$	$\begin{pmatrix} 1 & 1 \\ -1 & 0 \end{pmatrix}$	-	-	-
I_n	$(0, 0, n)$	$\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$	$\Gamma_1(n), \Gamma(n)$	$SU(n)$	\mathbb{Z}_n
III	$(\geq 1, \geq 2, 3)$	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$	$\Gamma_1(2)$	$SU(2)$	\mathbb{Z}_2
IV	$(\geq 2, 2, 4)$	$\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$	$\Gamma_1(3)$	$SU(3)$	\mathbb{Z}_3
I_{2n}^*	$(\geq 2, 3, 6 + 2n)$	$\begin{pmatrix} -1 & -2n \\ 0 & -1 \end{pmatrix}$	$\Gamma_1(2), \Gamma(2)$	$SO(8 + 4n)$	$\mathbb{Z}_2 \times \mathbb{Z}_2$
I_{2n+1}^*	$(\geq 2, 3, 7 + 2n)$	$\begin{pmatrix} -1 & -2n-1 \\ 0 & -1 \end{pmatrix}$	$\Gamma_1(2), \Gamma_1(4)$	$SO(10 + 4n)$	\mathbb{Z}_4
IV^*	$(\geq 3, 4, 8)$	$\begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}$	$\Gamma_1(3)$	E_6	\mathbb{Z}_3
III^*	$(3, \geq 5, 9)$	$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$	$\Gamma_1(2)$	E_7	\mathbb{Z}_2
II^*	$(4, \geq 5, 10)$	$\begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix}$	-	E_8	-

7 Brane monodromies consistent with center of the covering group ($\Gamma(p)$ fixes a $\mathbb{Z}_p \times \mathbb{Z}_p$ group)



Q: **What is the highest torsion point s_p a compact fibration can preserve?**

A: $T^2 : \mathbb{Z}_{12}$ $K3 : \mathbb{Z}_8$, $CY_{3/4} ??$ [Mazur'78; Oguiso, Shioda'91; Shimada'00]



What is the **moduli space** $X(1)$ of an elliptic curve \mathcal{E}_τ ?



- “The Key Hole” no structure: $X(1)^* = \tau \in \mathcal{H}/SL(2, \mathbb{Z})$
- There is a **cusp** at infinity where $\Delta = 4f^3 + 27g^2 = 0$
- Adding the point at infinity we obtain the compactified modular curve $X(1)$ topologically a \mathbb{P}^1 .



What is the **moduli space** $X(1)$ of an elliptic curve \mathcal{E}_τ ?



- “The Key Hole” no structure: $X(1)^* = \tau \in \mathcal{H}/SL(2, \mathbb{Z})$
- There is a **cusp** at infinity where $\Delta = 4f^3 + 27g^2 = 0$
- Adding the point at infinity we obtain the compactified modular curve $X(1)$ topologically a \mathbb{P}^1 .

Over a compact base space you **will always hit** the I_1 **cusp** somewhere

$X(1)$  $X_1(2)$ 

What is the moduli space $X_1(p)$ of an elliptic curve and a torsion point s_p ?



- **Tessellation** into triangular regions of modular curve $X_1(p) = \mathcal{H}/\Gamma_1(p)$

$X(1)$  $X_1(2)$ 

new I_2 cusp from 2 triangles meeting

What is the moduli space $X_1(p)$ of an elliptic curve and a torsion point s_p ?

- **Tessellation** into triangular regions of modular curve $X_1(p) = \mathcal{H}/\Gamma_1(p)$
- **Surprise: additional cusps of higher order** where triangles meet

$X(1)$  $X_1(2)$ 



new I_2 cusp from 2 triangles meeting

What is the moduli space $X_1(p)$ of an elliptic curve and a torsion point s_p ?

- **Tessellation** into triangular regions of modular curve $X_1(p) = \mathcal{H}/\Gamma_1(p)$
- **Surprise: additional cusps of higher order** where triangles meet

Equivalent construct the Weierstrass model explicitly [Aspinwall, Morrison '98]

$$s_2 @ y = x = 0 : \quad y^2 = x(x + a_2 x z^2 + a_4 z^4), \quad \Delta = \underbrace{a_4^2}_{I_2} \underbrace{(a^2 + a_4)}_{I_1} \quad a_i \in c_1(B)^i$$

$X(1)$  $X_1(2)$ 

new I_2 cusp from 2 triangles meeting


What is the moduli space $X_1(p)$ of an elliptic curve and a torsion point s_p ?

- **Tessellation** into triangular regions of modular curve $X_1(p) = \mathcal{H}/\Gamma_1(p)$
- **Surprise: additional cusps of higher order** where triangles meet

Equivalent construct the Weierstrass model explicitly [Aspinwall, Morrison '98]

$$s_2 @ y = x = 0 : \quad y^2 = x(x + a_2 x z^2 + a_4 z^4), \quad \Delta = \underbrace{a_4^2}_{I_2} \underbrace{(a^2 + a_4)}_{I_1} \quad a_i \in c_1(B)^i$$


Physics: $\pi_1(G^*/T) = \mathbb{Z}_2$ requires $G^* = SU(2)$ at least!



$\Gamma_1(p)$	T	G_{min}
$\Gamma_1(2)$	\mathbb{Z}_2	$SU(2)/\mathbb{Z}_2$
$\Gamma_1(3)$	\mathbb{Z}_3	$SU(3)/\mathbb{Z}_3$

Cusps in the modular curve **enforce minimal gauge groups** G with $\pi(G) = T$!


- How does this moduli space look like for large torsion?
- Larger torsion enforces **multiple** gauge algebra factors!



$\Gamma_1(p)$	T	G_{min}
$\Gamma_1(2)$	\mathbb{Z}_2	$SU(2)/\mathbb{Z}_2$
$\Gamma_1(3)$	\mathbb{Z}_3	$SU(3)/\mathbb{Z}_3$
$\Gamma_1(5)$	\mathbb{Z}_5	$SU(5)^2/\mathbb{Z}_5$

Cusps in the modular curve **enforce minimal gauge groups** G with $\pi(G) = T$!


- How does this moduli space look like for large torsion?
- Larger torsion enforces **multiple** gauge algebra factors!



$\Gamma_1(p)$	T	G_{min}
$\Gamma_1(2)$	\mathbb{Z}_2	$SU(2)/\mathbb{Z}_2$
$\Gamma_1(3)$	\mathbb{Z}_3	$SU(3)/\mathbb{Z}_3$
$\Gamma_1(5)$	\mathbb{Z}_5	$SU(5)^2/\mathbb{Z}_5$
$\Gamma_1(7)$	\mathbb{Z}_7	$SU(7)^3/\mathbb{Z}_7$

Cusps in the modular curve **enforce minimal gauge groups** G with $\pi(G) = T$!

- How does this moduli space look like for large torsion?
- Larger torsion enforces **multiple** gauge algebra factors!



$\Gamma_1(p)$	T	G_{min}
$\Gamma_1(2)$	\mathbb{Z}_2	$SU(2)/\mathbb{Z}_2$
$\Gamma_1(3)$	\mathbb{Z}_3	$SU(3)/\mathbb{Z}_3$
$\Gamma_1(5)$	\mathbb{Z}_5	$SU(5)^2/\mathbb{Z}_5$
$\Gamma_1(7)$	\mathbb{Z}_7	$SU(7)^3/\mathbb{Z}_7$
$\Gamma_1(9)$	\mathbb{Z}_9	$SU(9)^4/\mathbb{Z}_9$

Cusps in the modular curve **enforce minimal gauge groups** G with $\pi(G) = T$!

- How does this moduli space look like for large torsion?
- Larger torsion enforces **multiple** gauge algebra factors!

Torsion	$K(\mathbb{P}^1)$	K3	dP_9	$CY_{3/4}$
$\{0\}$	✓	✓	✓	✓
$\{\mathbb{Z}_2\}$	✓	✓	✓	✓
$\{\mathbb{Z}_3\}$	✓	✓	✓	✓
$\{\mathbb{Z}_4\}$	✓	✓	✓	✓
$\{\mathbb{Z}_2 \times \mathbb{Z}_2\}$	✓	✓	✓	✓
$\{\mathbb{Z}_5\}$	✓	✓	✓	✓
$\{\mathbb{Z}_6\}$	✓	✓	✓	✓
$\{\mathbb{Z}_2 \times \mathbb{Z}_4\}$	✓	✓	✓	✓
$\{\mathbb{Z}_3 \times \mathbb{Z}_3\}$	✓	✓	✓	✓
$\{\mathbb{Z}_7\}$	✓	✓	✗	✗
$\{\mathbb{Z}_8\}$	✓	✓	✗	✗
$\{\mathbb{Z}_2 \times \mathbb{Z}_6\}$	✓	✓	✗	✗
$\{\mathbb{Z}_2 \times \mathbb{Z}_8\}$	✓	✓	✗	✗
$\{\mathbb{Z}_4 \times \mathbb{Z}_4\}$	✓	✓	✗	✗
$\{\mathbb{Z}_9\}$	✓	✗	✗	✗
$\{\mathbb{Z}_{10}\}$	✓	✗	✗	✗
$\{\mathbb{Z}_{12}\}$	✓	✗	✗	✗
$\{\mathbb{Z}_3 \times \mathbb{Z}_6\}$	✓	✗	✗	✗
$\{\mathbb{Z}_5 \times \mathbb{Z}_5\}$	✓	✗	✗	✗

Constraints for surfaces:

- $\text{rank}(G_{\min}) > 8$ no dP_9
- $\text{rank}(G_{\min}) > 16$ no K3

[Miranda, Persson '89] ✓

Torsion	$K(\mathbb{P}^1)$	K3	dP_9	$CY_{3/4}$
$\{0\}$	✓	✓	✓	✓
$\{\mathbb{Z}_2\}$	✓	✓	✓	✓
$\{\mathbb{Z}_3\}$	✓	✓	✓	✓
$\{\mathbb{Z}_4\}$	✓	✓	✓	✓
$\{\mathbb{Z}_2 \times \mathbb{Z}_2\}$	✓	✓	✓	✓
$\{\mathbb{Z}_5\}$	✓	✓	✓	✓
$\{\mathbb{Z}_6\}$	✓	✓	✓	✓
$\{\mathbb{Z}_2 \times \mathbb{Z}_4\}$	✓	✓	✓	✓
$\{\mathbb{Z}_3 \times \mathbb{Z}_3\}$	✓	✓	✓	✓
$\{\mathbb{Z}_7\}$	✓	✓	✗	✗
$\{\mathbb{Z}_8\}$	✓	✓	✗	✗
$\{\mathbb{Z}_2 \times \mathbb{Z}_6\}$	✓	✓	✗	✗
$\{\mathbb{Z}_2 \times \mathbb{Z}_8\}$	✓	✓	✗	✗
$\{\mathbb{Z}_4 \times \mathbb{Z}_4\}$	✓	✓	✗	✗
$\{\mathbb{Z}_9\}$	✓	✗	✗	✗
$\{\mathbb{Z}_{10}\}$	✓	✗	✗	✗
$\{\mathbb{Z}_{12}\}$	✓	✗	✗	✗
$\{\mathbb{Z}_3 \times \mathbb{Z}_6\}$	✓	✗	✗	✗
$\{\mathbb{Z}_5 \times \mathbb{Z}_5\}$	✓	✗	✗	✗

Constraints for surfaces:

- $\text{rank}(G_{\min}) > 8$ no dP_9
- $\text{rank}(G_{\min}) > 16$ no K3

[Miranda, Persson '89] ✓


Constraints for $CY_{3/4}$

New non-minimal singularities:

$T = \mathbb{Z}_p$	Singularities
• $p < 4$	Regular
• $4 < p < 6$	(4,6,12) in codim 2
• $6 < p < 6$	(8,12,24) in codim 2
→ No crepant resolution ✗	

$CY_{3/4}$ folds: $T \in dP_9 \in E_8$
Aspinwall-Morrison '98 is complete!

SCFTs with restricted Monodromies



2 dim non-compact Base B_2


6D SCFT construction in IIB: Take B_2 to be **non-compact** and of the form of

- A generalized quiver of k **compact** \mathbb{P}^1 's wrapped by 7-branes giving rise to **gauge algebras** \mathfrak{g}_i and bifundamental matter
- **Non-compact Flavor branes** with flavor algebra \mathfrak{g}_{F_i}

[Witten'96; Seiberg'96; Harvey, Minasian,

Moore, Heckman, Vafa, Hanany, Zaffaroni, del Zotto, Rudelius, Morrison, Mekareeya, Tomasiello, Ohmori, Tachikawa,

Zafir, Park, Bertolini, Merks, Cordova, Dumitrescu, Intriligator, Bhardwaj...]



2 dim non-compact Base B_2

Flavor Brane

Flavor Brane


Compact \mathbb{P}^1 of selfintersection $-n_k$
and gauge algebra \mathfrak{g}_k

6D SCFT construction in IIB: Take B_2 to be **non-compact** and of the form of

- A generalized quiver of k **compact** \mathbb{P}^1 's wrapped by 7-branes giving rise to **gauge algebras** \mathfrak{g}_i and bifundamental matter
- **Non-compact Flavor branes** with flavor algebra \mathfrak{g}_{F_i}


[Witten '96; Seiberg '96; Harvey, Minasian,

Moore, Heckman, Vafa, Hanany, Zaffaroni, del Zotto, Rudelius, Morrison, Mekareeya, Tomasiello, Ohmori, Tachikawa, Zafrir, Park, Bertolini, Merks, Cordova, Dumitrescu, Intriligator, Bhardwaj...]



6D SCFT construction in IIB: Take B_2 to be **non-compact** and of the form of


- A generalized quiver of k **compact** \mathbb{P}^1 's wrapped by 7-branes giving rise to **gauge algebras** \mathfrak{g}_i and bifundamental matter
- **Non-compact Flavor branes** with flavor algebra \mathfrak{g}_{F_i}
- **Massive 6D strings** from D3 branes that wrap compact cycles
- **SCFT point** where strings become **tensionless** at the origin of the tensor branch by **contracting all curves** (if possible) [Witten'96; Seiberg'96; Harvey, Minasian, Moore, Heckman, Vafa, Hanany, Zaffaroni, del Zotto, Rudelius, Morrison, Mekareeya, Tomasiello, Ohmori, Tachikawa, Zafrir, Park, Bertolini, Merks, Cordova, Dumitrescu, Intriligator, Bhardwaj...]




6D SCFT construction in IIB: Take B_2 to be **non-compact** and of the form of

- A generalized quiver of k **compact** \mathbb{P}^1 's wrapped by 7-branes giving rise to **gauge algebras** \mathfrak{g}_i and bifundamental matter
- **Non-compact Flavor branes** with flavor algebra \mathfrak{g}_{F_i}
- **Massive 6D strings** from D3 branes that wrap compact cycles
- **SCFT point** where strings become **tensionless** at the origin of the tensor branch by **contracting all curves** (if possible) [Witten'96; Seiberg'96; Harvey, Minasian, Moore, Heckman, Vafa, Hanany, Zaffaroni, del Zotto, Rudelius, Morrison, Mekareeya, Tomasiello, Ohmori, Tachikawa, Zafrir, Park, Bertolini, Merks, Cordova, Dumitrescu, Intriligator, Bhardwaj...]

Construct SCFTs with global quotient structure in the Flavor group, by restricting to a congruence subgroup!




1. Basic building block: Non-Higgsable Clusters [Taylor Morrison'12]
 (chains) of \mathbb{P}^1 's with high negative self intersection $n > 2$



1. Basic building block: Non-Higgsable Clusters [Taylor Morrison'12]
 (chains) of \mathbb{P}^1 's with high negative self intersection $n > 2$

- Combine $n > 2$ curves with torsion models!



1. Basic building block: Non-Higgsable Clusters [Taylor Morrison'12]

(chains) of \mathbb{P}^1 's with high negative self intersection $n > 2$

- Combine $n > 2$ curves with torsion models!
- Torsion model **enhances** the NHC to a compatible gauge group

$$\mathbb{Z}_1 : \boxed{\mathfrak{su}_1} - \boxed{\mathfrak{e}_8}$$


2. Buidling Block: When bifundamental matter is not perturbatively allowed, it becomes superconformal matter, a 6D SCFT by itself

- **Simplest case**, E-string theory $\mathfrak{su}_1 \times \mathfrak{e}_8$ collision
- Changed to **discrete holonomy instanton** theories [Aspinwall, Morrison '98]

$$\mathbb{Z}_1 : \boxed{\mathfrak{su}_1} \circ \boxed{\mathfrak{e}_8}$$


2. Buidling Block: When bifundamental matter is not perturbatively allowed, it becomes superconformal matter, a 6D SCFT by itself

- **Simplest case**, E-string theory $\mathfrak{su}_1 \times \mathfrak{e}_8$ collision
- Changed to **discrete holonomy instanton** theories [Aspinwall, Morrison '98]



2. Buidling Block: When bifundamental matter is not perturbatively allowed, it becomes superconformal matter, a 6D SCFT by itself

- **Simplest case**, E-string theory $\mathfrak{su}_1 \times \mathfrak{e}_8$ collision
- Changed to **discrete holonomy instanton** theories [Aspinwall, Morrison'98]





2. Buidling Block: When bifundamental matter is not perturbatively allowed, it becomes superconformal matter, a 6D SCFT by itself

- **Simplest case**, E-string theory $\mathfrak{su}_1 \times \mathfrak{e}_8$ collision
- Changed to **discrete holonomy instanton** theories [Aspinwall, Morrison'98]

$$\mathbb{Z}_1 : \boxed{\mathfrak{e}_7} - \boxed{\mathfrak{su}_{2n}}$$


Higher rank collisions can be systematically constructed and their tensor branch can be analyzed

- Example: $[\mathfrak{e}_7] \times [\mathfrak{su}_{2n}]$ admits an diagonal \mathbb{Z}_2 center
Inconsistent with modding: \mathfrak{su}_{2k+1} factors, **56** matter, $(2k, \overline{2k-1})$ matter
- $([E_7] \times [SU(2n)])/\mathbb{Z}_2$ admits a **different tensor branch** gauge group and matter compatible with global modding [Ohmori, Tachikawa, Zafir'18]

$\mathbb{Z}_1 :$ 

Higher rank collisions can be systematically constructed and their tensor branch can be analyzed

- Example: $[e_7] \times [su_{2n}]$ admits an diagonal \mathbb{Z}_2 center
Inconsistent with modding: su_{2k+1} factors, **56** matter, $(2k, \overline{2k-1})$ matter
- $([E_7] \times [SU(2n)])/\mathbb{Z}_2$ admits a **different tensor branch** gauge group and matter compatible with global modding [Ohmori, Tachikawa, Zafir'18]



Higher rank collisions can be systematically constructed and their tensor branch can be analyzed

- Example: $[e_7] \times [su_{2n}]$ admits an diagonal \mathbb{Z}_2 center
Inconsistent with modding: su_{2k+1} factors, **56** matter, $(2k, \overline{2k-1})$ matter
- $([E_7] \times [SU(2n)])/\mathbb{Z}_2$ admits a **different tensor branch** gauge group and matter compatible with global modding [Ohmori, Tachikawa, Zafir'18]

$$\mathbb{Z}_3 : \boxed{\mathfrak{e}_6} - \boxed{\mathfrak{su}_{3n}} \quad \leftrightarrow \quad \boxed{\mathfrak{e}_6} - \overset{\mathfrak{su}_3}{\bigcirc 1} - \overset{\mathfrak{su}_6}{\bigcirc 2} - \dots - \overset{\mathfrak{su}_{3n-3}}{\bigcirc 2} - \boxed{\mathfrak{su}_{3n}}$$

A wide range of possibilities: exotic Matter, non-prime quotients, multiple generators.....

$$\begin{array}{l}
 \mathbb{Z}_3 : \quad \boxed{e_6} - \boxed{su_{3n}} \quad \leftrightarrow \quad \boxed{e_6} - \overset{su_3}{\bigcirc 1} - \overset{su_6}{\bigcirc 2} - \dots - \overset{su_{3n-3}}{\bigcirc 2} - \boxed{su_{3n}} \\
 \\
 \mathbb{Z}_3 : \quad \frac{1}{2} \mathbf{20} - \overset{su_6}{\bigcirc 1} - \overset{su_{15}}{\bigcirc 2} - \overset{su_{24}}{\bigcirc 2} - \dots - \overset{su_{9n-3}}{\bigcirc 2} - \boxed{\phantom{su_{9n+6}}} - su_{9n+6}
 \end{array}$$

A wide range of possibilities: exotic Matter, non-prime quotients, multiple generators.....

$$\mathbb{Z}_3 : \quad \boxed{\mathfrak{e}_6} - \boxed{\mathfrak{su}_{3n}} \quad \leftrightarrow \quad \boxed{\mathfrak{e}_6} \text{---} \overset{\mathfrak{su}_3}{(1)} \overset{\mathfrak{su}_6}{(2)} \overset{\mathfrak{su}_{3n-3}}{(2)} \cdots (2) \boxed{\mathfrak{su}_{3n}}$$

$$\mathbb{Z}_3 : \quad \frac{1}{2} \mathbf{20} \text{---} \overset{\mathfrak{su}_6}{(1)} \overset{\mathfrak{su}_{15}}{(2)} \overset{\mathfrak{su}_{24}}{(2)} \cdots \overset{\mathfrak{su}_{9n-3}}{(2)} \boxed{\phantom{\mathfrak{su}_{9n+6}}} \boxed{\mathfrak{su}_{9n+6}}$$

$$\mathbb{Z}_6 : \quad \boxed{\mathfrak{su}_6} \text{---} (1) \overset{\mathfrak{su}_3}{(2)} \overset{\mathfrak{su}_6}{(2)} \cdots \overset{\mathfrak{su}_{3n-3}}{(2)} \boxed{\mathfrak{su}_{3n}}$$

$\boxed{\mathfrak{su}_2}$

$$\mathbb{Z}_2 \times \mathbb{Z}_2 : \quad \boxed{\mathfrak{so}_{12}} \text{---} \overset{\mathfrak{su}_2}{(1)} \overset{\mathfrak{so}_8}{(3)} \overset{\mathfrak{sp}_1}{(1)} \overset{\mathfrak{so}_{12}}{(4)} \overset{\mathfrak{sp}_3}{(1)} \overset{\mathfrak{so}_{16}}{(4)} \cdots \overset{\mathfrak{sp}_{2n+1}}{(1)} \overset{\mathfrak{so}_{4n+4}}{(4)} \overset{\mathfrak{sp}_{2n+2}}{(1)} \boxed{\mathfrak{so}_{4n+8}}$$

$\boxed{\mathfrak{su}_2}$

A wide range of possibilities: exotic Matter, non-prime quotients, multiple generators.....

Elliptic fibrations with additional points s_p of order p $\pi_1(G) = T = \mathbb{Z}_p$.

Use the **same geometric approach** to construct **SCFTs** with non-simply connected flavor groups $\pi_1(G_F) = T$.

Elliptic fibrations with additional points s_p of order p $\pi_1(G) = T = \mathbb{Z}_p$.

- Fibrations have **restricted monodromies** that lie in a congruence subgroup $\Gamma_1(p)$ (or generalizations)
- **Minimal gauge groups** in compact geometries can be deduced from the modular curve $X_1(n) = \mathcal{H}/\Gamma_1(n)$
- **Swampland criterion** in compact geometries $\pi_1(G)$ must be finite Abelian subgroup of E_8 , otherwise non-crepant singularities (modulo 8D)

Use the **same geometric approach** to construct **SCFTs** with non-simply connected flavor groups $\pi_1(G_F) = T$.

Elliptic fibrations with additional points s_p of order p $\pi_1(G) = T = \mathbb{Z}_p$.

- Fibrations have **restricted monodromies** that lie in a congruence subgroup $\Gamma_1(p)$ (or generalizations)
- **Minimal gauge groups** in compact geometries can be deduced from the modular curve $X_1(n) = \mathcal{H}/\Gamma_1(n)$
- **Swampland criterion** in compact geometries $\pi_1(G)$ must be finite Abelian subgroup of E_8 , otherwise non-crepant singularities (modulo 8D)

Use the **same geometric approach** to construct **SCFTs** with non-simply connected flavor groups $\pi_1(G_F) = T$.

- **Enhanced** flavor and gauge algebras over **non-Higgsable clusters**
- **Quotient extends** to gauge group and matter at the tensor branch

Elliptic fibrations with additional points s_p of order p $\pi_1(G) = T = \mathbb{Z}_p$.

- Fibrations have **restricted monodromies** that lie in a congruence subgroup $\Gamma_1(p)$ (or generalizations)
- **Minimal gauge groups** in compact geometries can be deduced from the modular curve $X_1(n) = \mathcal{H}/\Gamma_1(n)$
- **Swampland criterion** in compact geometries $\pi_1(G)$ must be finite Abelian subgroup of E_8 , otherwise non-crepant singularities (modulo 8D)

Use the **same geometric approach** to construct **SCFTs** with non-simply connected flavor groups $\pi_1(G_F) = T$.

- **Enhanced** flavor and gauge algebras over **non-Higgsable clusters**
- **Quotient extends** to gauge group and matter at the tensor branch

Thank you very much!