

# 8D String Universality and Non-Simply-Connected Gauge Groups

Based on 2008.10605

with Mirjam Cvetič, Markus Dierigl and Ling Lin

Hao Y. Zhang

University of Pennsylvania

Sep. 29, 2020

# Section I: String Universality

*I*

Goal: String Universality

Focusing on 8D

Non-simply-connected Gauge Groups

*II*

F-theory on elliptic K3: 8D Vacua

Mordell-Weil Group and Arithmetics

*III*

Higher Form Center Symmetry

$B_4$  Periodicity and Mixed Anomaly

Anomaly-free 8D Theories

*IV*

Comparison: Strong Evidence for

8D String Universality

With  $G$  Non-simply-connected

*V*

Future Directions

# String Universality in 10D

# String Universality in 10D

String universality conjecture: All consistent theories of quantum gravity should come as low energy limits of string theory.

# String Universality in 10D

String universality conjecture: All consistent theories of quantum gravity should come as low energy limits of string theory.

10D  $\mathcal{N} = 1$  supergravities [Green, Schwarz, Witten, Vol.2]:

$$E_8 \times E_8, \quad SO(32), \quad E_8 \times U(1)^{248}, \quad U(1)^{496} \quad (1)$$

The first two are realized in heterotic string theories. Long time puzzle: the last two theories are neither realized in string theory nor ruled out.

# String Universality in 10D

String universality conjecture: All consistent theories of quantum gravity should come as low energy limits of string theory.

10D  $\mathcal{N} = 1$  supergravities [Green, Schwarz, Witten, Vol.2]:

$$E_8 \times E_8, \quad SO(32), \quad E_8 \times U(1)^{248}, \quad U(1)^{496} \quad (1)$$

The first two are realized in heterotic string theories. Long time puzzle: the last two theories are neither realized in string theory nor ruled out.

Solution: they are found inconsistent via the following non-stringy approaches:

- ▶ Via SUSY and Abelian gauge invariance [Adams, DeWolfe, Taylor '10]
- ▶ Unitarity bound:  $\text{rank} \leq 26-d = 16$  [Kim, Shiu, Vafa '19]

# String Universality in 10D

String universality conjecture: All consistent theories of quantum gravity should come as low energy limits of string theory.

10D  $\mathcal{N} = 1$  supergravities [Green, Schwarz, Witten, Vol.2]:

$$E_8 \times E_8, \quad SO(32), \quad E_8 \times U(1)^{248}, \quad U(1)^{496} \quad (1)$$

The first two are realized in heterotic string theories. Long time puzzle: the last two theories are neither realized in string theory nor ruled out.

Solution: they are found inconsistent via the following non-stringy approaches:

- ▶ Via SUSY and Abelian gauge invariance [Adams, DeWolfe, Taylor '10]
- ▶ Unitarity bound:  $\text{rank} \leq 26-d = 16$  [Kim, Shiu, Vafa '19]

Recent progress [Montero, Vafa '20]: e.g., 8D supergravity theories need to have rank 18, 10 or 2. This explains the modulo-8 rank pattern in 8D string vacua without referring to string theory.

# Global Structure of Gauge Groups



# Global Structure of Gauge Groups

Global structure of standard model gauge group - candidates:

$$(SU(3) \times SU(2) \times U(1))/\mathbb{Z}_n, \quad n = 1, 2, 3, 6 \quad (2)$$

# Global Structure of Gauge Groups

Global structure of standard model gauge group - candidates:

$$(SU(3) \times SU(2) \times U(1))/\mathbb{Z}_n, \quad n = 1, 2, 3, 6 \quad (2)$$

$SU(2)$  vs  $SO(3) \simeq SU(2)/\mathbb{Z}_2$ : e.g., **2** rep. allowed/forbidden.

# Global Structure of Gauge Groups

Global structure of standard model gauge group - candidates:

$$(SU(3) \times SU(2) \times U(1))/\mathbb{Z}_n, \quad n = 1, 2, 3, 6 \quad (2)$$

$SU(2)$  vs  $SO(3) \simeq SU(2)/\mathbb{Z}_2$ : e.g., **2** rep. allowed/forbidden.

Visible at the level of non-local objects: Wilson/'t Hooft lines, etc.

# Global Structure of Gauge Groups

Global structure of standard model gauge group - candidates:

$$(SU(3) \times SU(2) \times U(1))/\mathbb{Z}_n, \quad n = 1, 2, 3, 6 \quad (2)$$

$SU(2)$  vs  $SO(3) \simeq SU(2)/\mathbb{Z}_2$ : e.g., **2** rep. allowed/forbidden.

Visible at the level of non-local objects: Wilson/'t Hooft lines, etc.

Focus on 8D theories: 8D string vacua exhibits intricate patterns in gauge groups global structure. Can also be understood in F-theory via arithmetic structure of elliptic K3s.

# Global Structure of Gauge Groups

Global structure of standard model gauge group - candidates:

$$(SU(3) \times SU(2) \times U(1))/\mathbb{Z}_n, \quad n = 1, 2, 3, 6 \quad (2)$$


$SU(2)$  vs  $SO(3) \simeq SU(2)/\mathbb{Z}_2$ : e.g., **2** rep. allowed/forbidden.

Visible at the level of non-local objects: Wilson/'t Hooft lines, etc.

Focus on 8D theories: 8D string vacua exhibits intricate patterns in gauge groups global structure. Can also be understood in F-theory via arithmetic structure of elliptic K3s.

- ▶ Part II: Connection to F-theory arithmetics
- ▶ Part III: Connection to higher form symmetries

# Section II: 8D Stringy Vacua and Arithmetics



# F-theory on Elliptic K3s

# F-theory on Elliptic K3s

F-theory: Strong coupling version of type IIB,  $SL(2, \mathbb{Z})$  geometrized. [Vafa '96; Morrison Vafa '96], **review:** [Weigand '18]



# F-theory on Elliptic K3s

F-theory: Strong coupling version of type IIB,  $SL(2, \mathbb{Z})$  geometrized. [Vafa '96; Morrison Vafa '96], **review:** [Weigand '18]

F-theory 8D SUGRA vacua corresponds to 24 7-branes of specific  $(p, q)$  types locating on the base  $\mathbb{P}^1$ , whose overall monodromy is identity.

# F-theory on Elliptic K3s

F-theory: Strong coupling version of type IIB,  $SL(2, \mathbb{Z})$  geometrized. [Vafa '96; Morrison Vafa '96], **review:** [Weigand '18]


F-theory 8D SUGRA vacua corresponds to 24 7-branes of specific  $(p, q)$  types locating on the base  $\mathbb{P}^1$ , whose overall monodromy is identity.

Elliptically fibered K3 perspective: Gauge group can be read off via Kodaira's classification. Generically 24  $I_1$  fibers, collision gives enhancement.

# Mordell-Weil Torsion in F-theory and Global Structure of Gauge Group


# Mordell-Weil Torsion in F-theory and Global Structure of Gauge Group

Mordell-Weil(MW) group of an elliptic fibration consists of all of its global sections. The group law is the fiberwise addition law of cubic curves. See [Cvetič, Lin '18] for a review in the F-theory context.



# Mordell-Weil Torsion in F-theory and Global Structure of Gauge Group

Mordell-Weil(MW) group of an elliptic fibration consists of all of its global sections. The group law is the fiberwise addition law of cubic curves. See [Cvetič, Lin '18] for a review in the F-theory context.




As a finitely generated Abelian group, one has:

$$MW(Y) = \mathbb{Z}^r \oplus \bigoplus_i \mathbb{Z}_{n_i} \quad (3)$$

# Mordell-Weil Torsion in F-theory and Global Structure of Gauge Group

Mordell-Weil(MW) group of an elliptic fibration consists of all of its global sections. The group law is the fiberwise addition law of cubic curves. See [Cvetič, Lin '18] for a review in the F-theory context.



As a finitely generated Abelian group, one has:

$$MW(Y) = \mathbb{Z}^r \oplus \bigoplus_i \mathbb{Z}_{n_i} \quad (3)$$

- ▶ Each free section gives a  $U(1)$  gauge symmetry [Morrison, Park '12], and a quotient  $\frac{G_{nA} \times U(1)}{Z(G)}$  at the presence of  $G_{nA}$  [Cvetič, Lin '17].
- ▶ Each  $\ell$ -torsional section gives a  $\mathbb{Z}_\ell$  global quotient: **Main**

**focus.** [Aspinwall, Morrison '98][Mayrhofer, Morrison, Till, Weigand '14][Hajouji, Oehlmann '19].

# Geometric Properties in Elliptic K3s

# Geometric Properties in Elliptic K3s

$G$	[1]	[2]	[3]	[4]	[5]	[6]	[7]	[8]	[2, 2]	[4, 2]	[6, 2]	[3, 3]	[4, 4]	Total
$ \mathcal{P}^G $	2746	732	85	41	6	10	1	1	61	5	1	3	1	3693


**Figure:** A complete list of ADE fiber types in elliptic K3s [Shimada '00].  $[n]$ : a  $\mathbb{Z}_n$  MW torsion,  $[n, m]$ : a  $\mathbb{Z}_n \times \mathbb{Z}_m$  MW torsion.



# Geometric Properties in Elliptic K3s

$G$	[1]	[2]	[3]	[4]	[5]	[6]	[7]	[8]	[2, 2]	[4, 2]	[6, 2]	[3, 3]	[4, 4]	Total
$ \mathcal{P}^G $	2746	732	85	41	6	10	1	1	61	5	1	3	1	3693

**Figure:** A complete list of ADE fiber types in elliptic K3s [Shimada '00].  $[n]$ : a  $\mathbb{Z}_n$  MW torsion,  $[n, m]$ : a  $\mathbb{Z}_n \times \mathbb{Z}_m$  MW torsion.



**Figure:**  $k_i$ : # of component in  $I_{n_i}$  fiber intersected by the  $\ell$ -torsional section.

# Arithmetic Constraint on Global Structure of Gauge Groups

# Arithmetic Constraint on Global Structure of Gauge Groups

For a K3 with only  $I_n$  fibers and a single  $\ell$ -torsional section, the global gauge group is  $(\bigotimes_i SU(n_i))/\mathbb{Z}_\ell$ .

# Arithmetic Constraint on Global Structure of Gauge Groups

For a K3 with only  $I_n$  fibers and a single  $\ell$ -torsional section, the global gauge group is  $(\bigotimes_i SU(n_i))/\mathbb{Z}_\ell$ .

But what is the embedding? Now  $G = \bigotimes_i SU(n_i)$ , and this  $(k_1, \dots, k_s)$  is precisely the generator of the embedded  $\mathbb{Z}_\ell$  quotient into the overall center  $Z(G) = \bigotimes_i Z(G_i)$ .

# Arithmetic Constraint on Global Structure of Gauge Groups

For a K3 with only  $I_n$  fibers and a single  $\ell$ -torsional section, the global gauge group is  $(\bigotimes_i SU(n_i))/\mathbb{Z}_\ell$ .

But what is the embedding? Now  $G = \bigotimes_i SU(n_i)$ , and this  $(k_1, \dots, k_s)$  is precisely the generator of the embedded  $\mathbb{Z}_\ell$  quotient into the overall center  $Z(G) = \bigotimes_i Z(G_i)$ .

Not all embedding are allowed. [Miranda, Persson '88 '89 '91]:

$$q(k_1, \dots, k_s) = \sum_{i=1}^s \frac{n_i - 1}{2n_i} k_i^2 \equiv 0 \pmod{\mathbb{Z}} \quad (4)$$

# Arithmetic Constraint on Global Structure of Gauge Groups

For a K3 with only  $I_n$  fibers and a single  $\ell$ -torsional section, the global gauge group is  $(\bigotimes_i SU(n_i))/\mathbb{Z}_\ell$ .

But what is the embedding? Now  $G = \bigotimes_i SU(n_i)$ , and this  $(k_1, \dots, k_s)$  is precisely the generator of the embedded  $\mathbb{Z}_\ell$  quotient into the overall center  $Z(G) = \bigotimes_i Z(G_i)$ .

Not all embedding are allowed. [Miranda, Persson '88 '89 '91]:

$$q(k_1, \dots, k_s) = \sum_{i=1}^s \frac{n_i - 1}{2n_i} k_i^2 \equiv 0 \pmod{\mathbb{Z}} \quad (4)$$

e.g.  $G = SU(7)^3/\mathbb{Z}_7$  with a  $\mathbb{Z}_7$  torsional section:  
 $(k_1, k_2, k_3) = (1, 2, 3)$  is an allowed intersection pattern, corresponding to an allowed  $\mathbb{Z}_7$  center embedding.  
However,  $(k_1, k_2, k_3) = (1, 1, 1)$  is not allowed.

# Arithmetic Constraint on Global Structure of Gauge Groups

For a K3 with only  $I_n$  fibers and a single  $\ell$ -torsional section, the global gauge group is  $(\bigotimes_i SU(n_i))/\mathbb{Z}_\ell$ .

But what is the embedding? Now  $G = \bigotimes_i SU(n_i)$ , and this  $(k_1, \dots, k_s)$  is precisely the generator of the embedded  $\mathbb{Z}_\ell$  quotient into the overall center  $Z(G) = \bigotimes_i Z(G_i)$ .


Not all embedding are allowed. [Miranda, Persson '88 '89 '91]:

$$q(k_1, \dots, k_s) = \sum_{i=1}^s \frac{n_i - 1}{2n_i} k_i^2 \equiv 0 \pmod{\mathbb{Z}} \quad (4)$$

e.g.  $G = SU(7)^3/\mathbb{Z}_7$  with a  $\mathbb{Z}_7$  torsional section:  
 $(k_1, k_2, k_3) = (1, 2, 3)$  is an allowed intersection pattern, corresponding to an allowed  $\mathbb{Z}_7$  center embedding.  
However,  $(k_1, k_2, k_3) = (1, 1, 1)$  is not allowed.

**Any field-theoretic reasons behind this?**

# Section III: Higher Form Symmetry and Mixed Anomalies





# Higher-form Global Symmetry

# Higher-form Global Symmetry

The operators charged under a  $q$ -form global symmetry are of dimension  $q$ . They have to be Abelian for  $q \geq 1$ . [Gaiotto, Kapustin, Seiberg, Willett '14] Here we focus on the class of center higher form symmetries.

# Higher-form Global Symmetry

The operators charged under a  $q$ -form global symmetry are of dimension  $q$ . They have to be Abelian for  $q \geq 1$ . [Gaiotto, Kapustin, Seiberg, Willett '14] Here we focus on the class of center higher form symmetries.

- ▶ A continuous example: 4D  $U(1)$  gauge theory: an *electric* 1-form global  $U(1)$  whose charged objects are the Wilson lines. (We ignore the magnetic symmetry for our purpose.)

# Higher-form Global Symmetry

The operators charged under a  $q$ -form global symmetry are of dimension  $q$ . They have to be Abelian for  $q \geq 1$ . [Gaiotto, Kapustin, Seiberg, Willett '14] Here we focus on the class of center higher form symmetries.

- ▶ A continuous example: 4D  $U(1)$  gauge theory: an *electric* 1-form global  $U(1)$  whose charged objects are the Wilson lines. (We ignore the magnetic symmetry for our purpose.)
- ▶ A discrete example: 4D  $SU(N)$  theory has a center  $Z(SU(N)) = \mathbb{Z}_N$ -valued 1-form electric global symmetry charging the discrete Wilson lines. **Main focus.**

# Gauging 1-form Global Symmetry

# Gauging 1-form Global Symmetry

For  $G$  gauge theories without any matter charged under a subgroup  $\Gamma \subset Z(G)$  of the center, it has a  $\Gamma$ -valued 1-form global symmetry.

# Gauging 1-form Global Symmetry

For  $G$  gauge theories without any matter charged under a subgroup  $\Gamma \subset Z(G)$  of the center, it has a  $\Gamma$ -valued 1-form global symmetry.

To gauge this 1-form global symmetry, one first couple it to a 2-form  $\Gamma$ -valued background gauge field  $C$ , reducing the electric global symmetry:  $Z(G) \rightarrow Z(G)/\Gamma$ .

# Gauging 1-form Global Symmetry

For  $G$  gauge theories without any matter charged under a subgroup  $\Gamma \subset Z(G)$  of the center, it has a  $\Gamma$ -valued 1-form global symmetry.

To gauge this 1-form global symmetry, one first couple it to a 2-form  $\Gamma$ -valued background gauge field  $C$ , reducing the electric global symmetry:  $Z(G) \rightarrow Z(G)/\Gamma$ .

One then sum over all possible  $C$ -configurations in the partition function, giving rise to a  $G/\Gamma$  partition function in the 0-form sector.



# Gauging 1-form Global Symmetry

For  $G$  gauge theories without any matter charged under a subgroup  $\Gamma \subset Z(G)$  of the center, it has a  $\Gamma$ -valued 1-form global symmetry.

To gauge this 1-form global symmetry, one first couple it to a 2-form  $\Gamma$ -valued background gauge field  $C$ , reducing the electric global symmetry:  $Z(G) \rightarrow Z(G)/\Gamma$ .

One then sum over all possible  $C$ -configurations in the partition function, giving rise to a  $G/\Gamma$  partition function in the 0-form sector.

In the end, the  $G$  bundle is twisted into a  $G/\Gamma$  bundle with a non-trivial  $\Gamma$ -valued second Stiefel-Whitney class.

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 4D Field Theories

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 4D Field Theories

In 4d Yang-Mills theories, there is a mixed anomaly between the 1-form global symmetry and the  $\theta$ -periodicity [Córdova, Freed, Lam, Seiberg '19].

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 4D Field Theories

In 4d Yang-Mills theories, there is a mixed anomaly between the 1-form global symmetry and the  $\theta$ -periodicity [Córdova, Freed, Lam, Seiberg '19].

$\frac{i\theta}{8\pi^2} \int \text{Tr}(F \wedge F)$  under a 1-form symmetry transformation and  $\theta \rightarrow \theta + 2\pi$  gains:

$$\mathcal{A}(\theta, C) = 2\pi i \alpha_G \int_{M_4} \mathcal{P}(C) \mod \mathbb{Z} \quad (5)$$

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 4D Field Theories

In 4d Yang-Mills theories, there is a mixed anomaly between the 1-form global symmetry and the  $\theta$ -periodicity [Córdova, Freed, Lam, Seiberg '19].

$\frac{i\theta}{8\pi^2} \int \text{Tr}(F \wedge F)$  under a 1-form symmetry transformation and  $\theta \rightarrow \theta + 2\pi$  gains:

$$\mathcal{A}(\theta, C) = 2\pi i \alpha_G \int_{M_4} \mathcal{P}(C) \mod \mathbb{Z} \quad (5)$$

$\alpha_G$ :  $G$ -dependent fractional group theory constant. e.g.,

$$\alpha_G = \frac{N-1}{2N} \text{ for } G = SU(N)$$

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 4D Field Theories

In 4d Yang-Mills theories, there is a mixed anomaly between the 1-form global symmetry and the  $\theta$ -periodicity [Córdova, Freed, Lam, Seiberg '19].

$\frac{i\theta}{8\pi^2} \int \text{Tr}(F \wedge F)$  under a 1-form symmetry transformation and  $\theta \rightarrow \theta + 2\pi$  gains:

$$\mathcal{A}(\theta, C) = 2\pi i \alpha_G \int_{M_4} \mathcal{P}(C) \mod \mathbb{Z} \quad (5)$$

$\alpha_G$ :  $G$ -dependent fractional group theory constant. e.g.,

$$\alpha_G = \frac{N-1}{2N} \text{ for } G = SU(N)$$

$\mathcal{P}(C)$ : Pontryagin square of  $C \in H^2(X, \mathbb{Z}_N)$ , an integral element of the 4-th cohomology. ( $\mathcal{P}(C)$  can be roughly understood as  $C \wedge C$  when promoted to differential forms.)

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 8D Supergravity

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 8D Supergravity

In 8D  $\mathcal{N} = 1$ , gravity multiplet contains  $B_2$  [Salam, Sezgin '85]. It can be dualized to  $B_4$  [Awada, Townsend '85] with coupling  $(G = \bigotimes_i G_i)$ :

$$\sum_i \int_{M_8} B_4 \wedge \frac{1}{2} \text{Tr}(F_i \wedge F_i) \quad (6)$$



# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 8D Supergravity

In 8D  $\mathcal{N} = 1$ , gravity multiplet contains  $B_2$  [Salam, Sezgin '85]. It can be dualized to  $B_4$  [Awada, Townsend '85] with coupling ( $G = \bigotimes_i G_i$ ):

$$\sum_i \int_{M_8} B_4 \wedge \frac{1}{2} \text{Tr}(F_i \wedge F_i) \quad (6)$$

A gauged  $\Gamma \subset Z(G) = \bigotimes_i Z(G_i)$  should be coupled to  $C^{(i)}$ 's each valued in  $Z(G_i)$ , giving an instanton contribution:

$$I_4(G/\Gamma) = \sum_i \alpha_{G_i} \mathcal{P}(C^{(i)}) \mod \mathbb{Z} \quad (7)$$

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 8D Supergravity

In 8D  $\mathcal{N} = 1$ , gravity multiplet contains  $B_2$  [Salam, Sezgin '85]. It can be dualized to  $B_4$  [Awada, Townsend '85] with coupling ( $G = \bigotimes_i G_i$ ):

$$\sum_i \int_{M_8} B_4 \wedge \frac{1}{2} \text{Tr}(F_i \wedge F_i) \quad (6)$$

A gauged  $\Gamma \subset Z(G) = \bigotimes_i Z(G_i)$  should be coupled to  $C^{(i)}$ 's each valued in  $Z(G_i)$ , giving an instanton contribution:

$$I_4(G/\Gamma) = \sum_i \alpha_{G_i} \mathcal{P}(C^{(i)}) \mod \mathbb{Z} \quad (7)$$

Together with the integral period shift  $B_4 \rightarrow B_4 + b_4$ , we obtain:

$$\mathcal{A}(b_4, C^{(i)}) = 2\pi i \sum_i \alpha_{G_i} \int_{M_8} b_4 \cup \mathcal{P}(C^{(i)}) \mod \mathbb{Z} \quad (8)$$

# $\mathbb{Z}_N$ 1-form Symmetry and Mixed Anomaly in 8D Supergravity

In 8D  $\mathcal{N} = 1$ , gravity multiplet contains  $B_2$  [Salam, Sezgin '85]. It can be dualized to  $B_4$  [Awada, Townsend '85] with coupling ( $G = \bigotimes_i G_i$ ):

$$\sum_i \int_{M_8} B_4 \wedge \frac{1}{2} \text{Tr}(F_i \wedge F_i) \quad (6)$$

A gauged  $\Gamma \subset Z(G) = \bigotimes_i Z(G_i)$  should be coupled to  $C^{(i)}$ 's each valued in  $Z(G_i)$ , giving an instanton contribution:

$$I_4(G/\Gamma) = \sum_i \alpha_{G_i} \mathcal{P}(C^{(i)}) \mod \mathbb{Z} \quad (7)$$

Together with the integral period shift  $B_4 \rightarrow B_4 + b_4$ , we obtain:

$$\mathcal{A}(b_4, C^{(i)}) = 2\pi i \sum_i \alpha_{G_i} \int_{M_8} b_4 \cup \mathcal{P}(C^{(i)}) \mod \mathbb{Z} \quad (8)$$

In 6D theories, an analogous anomaly comes from a GS term

$$\int_{M_6} \Omega_{ij} B^i \wedge \text{Tr}(F^j \wedge F^j) \text{ [Apruzzi, Dierigl, Lin '20].}$$

# Anomaly-free 8D Supergravity Theories

# Anomaly-free 8D Supergravity Theories

$$\mathcal{A}(b_4, C^{(i)}) = 2\pi i \sum_i \alpha_{G_i} \int_{M_8} b_4 \cup \mathcal{P}(C^{(i)}) \mod \mathbb{Z} \quad (9)$$

According to [Cordova, Ohmori '19], such mixed anomaly cannot be cancelled. It also gives a fractional electric  $B_4$  charge, which is not consistent [Apruzzi, Dierigl, Lin '20]. **So it must vanish modulo integer.**

# Anomaly-free 8D Supergravity Theories

$$\mathcal{A}(b_4, C^{(i)}) = 2\pi i \sum_i \alpha_{G_i} \int_{M_8} b_4 \cup \mathcal{P}(C^{(i)}) \mod \mathbb{Z} \quad (9)$$

According to [Cordova, Ohmori '19], such mixed anomaly cannot be cancelled. It also gives a fractional electric  $B_4$  charge, which is not consistent [Apruzzi, Dierigl, Lin '20]. **So it must vanish modulo integer.**

$\bigoplus_i SU(N_i)$  case: assuming we have a gauged  $\mathbb{Z}_\ell$  factor with a  $\mathbb{Z}_\ell$ -valued 2-form  $C$  field. We further set  $C^{(i)} = k^i C$ , namely the  $\mathbb{Z}_\ell$  quotient of  $G/\mathbb{Z}_\ell$  is generated by  $(k_1, \dots, k_s)$  in  $Z(G)$ . Then one has a constraint:

$$\sum_i \frac{N_i - 1}{2N_i} k_i^2 \in \mathbb{Z} \quad (10)$$

# Anomaly-free 8D Supergravity Theories

$$\mathcal{A}(b_4, C^{(i)}) = 2\pi i \sum_i \alpha_{G_i} \int_{M_8} b_4 \cup \mathcal{P}(C^{(i)}) \mod \mathbb{Z} \quad (9)$$


According to [Cordova, Ohmori '19], such mixed anomaly cannot be cancelled. It also gives a fractional electric  $B_4$  charge, which is not consistent [Apruzzi, Dierigl, Lin '20]. **So it must vanish modulo integer.**

$\bigoplus_i SU(N_i)$  case: assuming we have a gauged  $\mathbb{Z}_\ell$  factor with a  $\mathbb{Z}_\ell$ -valued 2-form  $C$  field. We further set  $C^{(i)} = k^i C$ , namely the  $\mathbb{Z}_\ell$  quotient of  $G/\mathbb{Z}_\ell$  is generated by  $(k_1, \dots, k_s)$  in  $Z(G)$ . Then one has a constraint:

$$\sum_i \frac{N_i - 1}{2N_i} k_i^2 \in \mathbb{Z} \quad (10)$$

**This is exactly the same as the arithmetic condition for elliptic K3s with only  $I_n$  singularities!**

## IV: Results and Conclusions





# Constraints on $G$ with Non-trivial $\pi_1(G)$

## Constraints on $G$ with Non-trivial $\pi_1(G)$

(a) Product-of- $SU$  gauge groups. Assuming there is a non-trivial  $\mathbb{Z}_N$  gauged 1-form symmetry, also taking into account the rank  $\leq 26 - d = 18$  unitarity bound [Kim, Shiu, Vafa '19]:

# Constraints on $G$ with Non-trivial $\pi_1(G)$

(a) Product-of- $SU$  gauge groups. Assuming there is a non-trivial  $\mathbb{Z}_N$  gauged 1-form symmetry, also taking into account the rank  $\leq 26 - d = 18$  unitarity bound [Kim, Shiu, Vafa '19]:

- ▶ For  $N = p^k \geq 7$  power of prime, all anomaly free theories are realized via string theory.

# Constraints on $G$ with Non-trivial $\pi_1(G)$

(a) Product-of- $SU$  gauge groups. Assuming there is a non-trivial  $\mathbb{Z}_N$  gauged 1-form symmetry, also taking into account the rank  $\leq 26 - d = 18$  unitarity bound [Kim, Shiu, Vafa '19]:

- ▶ For  $N = p^k \geq 7$  power of prime, all anomaly free theories are realized via string theory.
  - ▶ Exactly one anomaly free theory for  $p^k = 7, 8$  resp.
  - ▶ No anomaly free theories for  $9 \leq p^k \leq 19$ .

# Constraints on $G$ with Non-trivial $\pi_1(G)$

(a) Product-of- $SU$  gauge groups. Assuming there is a non-trivial  $\mathbb{Z}_N$  gauged 1-form symmetry, also taking into account the rank  $\leq 26 - d = 18$  unitarity bound [Kim, Shiu, Vafa '19]:

- ▶ For  $N = p^k \geq 7$  power of prime, all anomaly free theories are realized via string theory.
  - ▶ Exactly one anomaly free theory for  $p^k = 7, 8$  resp.
  - ▶ No anomaly free theories for  $9 \leq p^k \leq 19$ .
- ▶  $N$  not a power of prime. If one assume that  $\mathbb{Z}_N$  acts faithfully on one simple  $G_i$  factor, then all such  $10 \leq N \leq 18$  give no anomaly-free theories.

# Constraints on $G$ with Non-trivial $\pi_1(G)$

(a) Product-of- $SU$  gauge groups. Assuming there is a non-trivial  $\mathbb{Z}_N$  gauged 1-form symmetry, also taking into account the rank  $\leq 26 - d = 18$  unitarity bound [Kim, Shiu, Vafa '19]:

- ▶ For  $N = p^k \geq 7$  power of prime, all anomaly free theories are realized via string theory.
  - ▶ Exactly one anomaly free theory for  $p^k = 7, 8$  resp.
  - ▶ No anomaly free theories for  $9 \leq p^k \leq 19$ .
- ▶  $N$  not a power of prime. If one assume that  $\mathbb{Z}_N$  acts faithfully on one simple  $G_i$  factor, then all such  $10 \leq N \leq 18$  give no anomaly-free theories.

(b) Single-factor gauge groups.

# Constraints on $G$ with Non-trivial $\pi_1(G)$

(a) Product-of- $SU$  gauge groups. Assuming there is a non-trivial  $\mathbb{Z}_N$  gauged 1-form symmetry, also taking into account the rank  $\leq 26 - d = 18$  unitarity bound [Kim, Shiu, Vafa '19]:

- ▶ For  $N = p^k \geq 7$  power of prime, all anomaly free theories are realized via string theory.
  - ▶ Exactly one anomaly free theory for  $p^k = 7, 8$  resp.
  - ▶ No anomaly free theories for  $9 \leq p^k \leq 19$ .
- ▶  $N$  not a power of prime. If one assume that  $\mathbb{Z}_N$  acts faithfully on one simple  $G_i$  factor, then all such  $10 \leq N \leq 18$  give no anomaly-free theories.

(b) Single-factor gauge groups.

- ▶  $\frac{SU(16)}{\mathbb{Z}_2}, \quad \frac{SU(18)}{\mathbb{Z}_3}, \quad \frac{Spin(32)}{\mathbb{Z}_2} : \text{F-theory on K3}$

## Constraints on $G$ with Non-trivial $\pi_1(G)$

(a) Product-of- $SU$  gauge groups. Assuming there is a non-trivial  $\mathbb{Z}_N$  gauged 1-form symmetry, also taking into account the rank  $\leq 26 - d = 18$  unitarity bound [Kim, Shiu, Vafa '19]:

- ▶ For  $N = p^k \geq 7$  power of prime, all anomaly free theories are realized via string theory.
  - ▶ Exactly one anomaly free theory for  $p^k = 7, 8$  resp.
  - ▶ No anomaly free theories for  $9 \leq p^k \leq 19$ .
- ▶  $N$  not a power of prime. If one assume that  $\mathbb{Z}_N$  acts faithfully on one simple  $G_i$  factor, then all such  $10 \leq N \leq 18$  give no anomaly-free theories.

(b) Single-factor gauge groups.

- ▶  $\frac{SU(16)}{\mathbb{Z}_2}$ ,  $\frac{SU(18)}{\mathbb{Z}_3}$ ,  $\frac{Spin(32)}{\mathbb{Z}_2}$  : F-theory on K3
- ▶  $\frac{Sp(4)}{\mathbb{Z}_2}$ ,  $\frac{Sp(8)}{\mathbb{Z}_2}$ ,  $\frac{SU(8)}{\mathbb{Z}_2}$ ,  $\frac{SU(9)}{\mathbb{Z}_3}$ ,  $\frac{Spin(16)}{\mathbb{Z}_2}$  : 8D CHL



# Constraints on $G$ with Non-trivial $\pi_1(G)$

(a) Product-of- $SU$  gauge groups. Assuming there is a non-trivial  $\mathbb{Z}_N$  gauged 1-form symmetry, also taking into account the rank  $\leq 26 - d = 18$  unitarity bound [Kim, Shiu, Vafa '19]:

- ▶ For  $N = p^k \geq 7$  power of prime, all anomaly free theories are realized via string theory.
  - ▶ Exactly one anomaly free theory for  $p^k = 7, 8$  resp.
  - ▶ No anomaly free theories for  $9 \leq p^k \leq 19$ .
- ▶  $N$  not a power of prime. If one assume that  $\mathbb{Z}_N$  acts faithfully on one simple  $G_i$  factor, then all such  $10 \leq N \leq 18$  give no anomaly-free theories.

(b) Single-factor gauge groups.

- ▶  $\frac{SU(16)}{\mathbb{Z}_2}$ ,  $\frac{SU(18)}{\mathbb{Z}_3}$ ,  $\frac{Spin(32)}{\mathbb{Z}_2}$  : F-theory on K3
- ▶  $\frac{Sp(4)}{\mathbb{Z}_2}$ ,  $\frac{Sp(8)}{\mathbb{Z}_2}$ ,  $\frac{SU(8)}{\mathbb{Z}_2}$ ,  $\frac{SU(9)}{\mathbb{Z}_3}$ ,  $\frac{Spin(16)}{\mathbb{Z}_2}$  : 8D CHL
- ▶  $\frac{Sp(12)}{\mathbb{Z}_2}$ ,  $\frac{Sp(16)}{\mathbb{Z}_2}$  : **Unknown**

# Conclusions

# Conclusions

We have provided strong evidence for string universality for non-simply-connect non-Abelian gauge groups  $G/Z$  in eight dimensions.

# Conclusions

We have provided strong evidence for string universality for non-simply-connect non-Abelian gauge groups  $G/Z$  in eight dimensions.

- ▶ We observe intricate patterns and restrictions regarding global structure of gauge groups in 8D string vacua.

# Conclusions

We have provided strong evidence for string universality for non-simply-connect non-Abelian gauge groups  $G/Z$  in eight dimensions.

- ▶ We observe intricate patterns and restrictions regarding global structure of gauge groups in 8D string vacua.
  - ▶ This corresponds to patterns in arithmetic structures of elliptic fibrations in F-theory.

# Conclusions

We have provided strong evidence for string universality for non-simply-connect non-Abelian gauge groups  $G/Z$  in eight dimensions.

- ▶ We observe intricate patterns and restrictions regarding global structure of gauge groups in 8D string vacua.
  - ▶ This corresponds to patterns in arithmetic structures of elliptic fibrations in F-theory.
- ▶ Key supergravity constraint: the mixed anomaly between center 1-form symmetry and large gauge transformation of the  $B_4$  supergravity field should vanish.

# Conclusions

We have provided strong evidence for string universality for non-simply-connect non-Abelian gauge groups  $G/Z$  in eight dimensions.

- ▶ We observe intricate patterns and restrictions regarding global structure of gauge groups in 8D string vacua.
  - ▶ This corresponds to patterns in arithmetic structures of elliptic fibrations in F-theory.
- ▶ Key supergravity constraint: the mixed anomaly between center 1-form symmetry and large gauge transformation of the  $B_4$  supergravity field should vanish.
  - ▶ Such a constraint nicely matches with arithmetic properties of elliptic K3 surfaces in the pure  $I_n$  singularity case.

# Conclusions

We have provided strong evidence for string universality for non-simply-connect non-Abelian gauge groups  $G/Z$  in eight dimensions.

- ▶ We observe intricate patterns and restrictions regarding global structure of gauge groups in 8D string vacua.
  - ▶ This corresponds to patterns in arithmetic structures of elliptic fibrations in F-theory.
- ▶ Key supergravity constraint: the mixed anomaly between center 1-form symmetry and large gauge transformation of the  $B_4$  supergravity field should vanish.
  - ▶ Such a constraint nicely matches with arithmetic properties of elliptic K3 surfaces in the pure  $I_n$  singularity case.
- ▶ Via this constraint we can rule out a large family of apparently consistent 8D  $\mathcal{N} = 1$  theories (with  $G$  non-simply-connected) that cannot be realized via string theory.




# Conclusions

We have provided strong evidence for string universality for non-simply-connect non-Abelian gauge groups  $G/Z$  in eight dimensions.

- ▶ We observe intricate patterns and restrictions regarding global structure of gauge groups in 8D string vacua.
  - ▶ This corresponds to patterns in arithmetic structures of elliptic fibrations in F-theory.
- ▶ Key supergravity constraint: the mixed anomaly between center 1-form symmetry and large gauge transformation of the  $B_4$  supergravity field should vanish.
  - ▶ Such a constraint nicely matches with arithmetic properties of elliptic K3 surfaces in the pure  $I_n$  singularity case.
- ▶ Via this constraint we can rule out a large family of apparently consistent 8D  $\mathcal{N} = 1$  theories (with  $G$  non-simply-connected) that cannot be realized via string theory. → Towards string universality in 8D.

# V: Future Directions



# Future Directions

# Future Directions

- ▶ One can seek for new insights into the realization of 8D CHL string gauge groups with global quotients. (See also [Font, Fraiman, Graña, Núñez, Parra '20] )

# Future Directions

- ▶ One can seek for new insights into the realization of 8D CHL string gauge groups with global quotients. (See also [Font, Fraiman, Graña, Núñez, Parra '20] )
- ▶ Incorporating the  $U(1)$ s will introduce additional global quotients  $(G \times U(1))/Z(G)$  in F-theory [Cvetič, Lin '17]. One could seek for a field-theoretic explanation for this, possibly via unHiggsing.

# Future Directions

- ▶ One can seek for new insights into the realization of 8D CHL string gauge groups with global quotients. (See also [Font, Fraiman, Graña, Núñez, Parra '20] )
- ▶ Incorporating the  $U(1)$ s will introduce additional global quotients  $(G \times U(1))/Z(G)$  in F-theory [Cvetič, Lin '17]. One could seek for a field-theoretic explanation for this, possibly via unHiggsing.
- ▶ Finally, we can compare this analysis with other swampland constraints. [McNamara, Vafa '19][Montero, Vafa '20]. In particular, the results obtained in [Montero, Vafa '20] regarding non-simply-connected gauge groups in 8D is consistent with ours.

Thanks!

terima kasih

આભાર

*salamat*

drink u

*Danke*

спасибо

감사합니다 merci

*Tak*

शुक्रिया

σας ευχαριστώ

 $a\check{c}i\bar{u}$ 

Thank you

謝謝

شکرا

ありがとう

*aitäh*

**ขอบเขต**

*obrigado*