
COBORDISMS, ANOMALIES, AND SWAMPLAND IN NINE DIMENSIONS

Miguel Montero

Harvard

StringPheno Summer series

July 21 2020



-
- In QG, we expect there are **no global symmetries.**
[Wheeler in the 50's??]

-
- In QG, we expect there are **no global symmetries.**
[Wheeler in the 50's??]
 - Lots of **evidence**, proven in worldsheet & AdS/CFT.
[Banks '88, Polchinski textbook, Ooguri-Harlow '18]

-
- In QG, we expect there are **no global symmetries.**
[Wheeler in the 50's??]
 - Lots of **evidence**, proven in worldsheet & AdS/CFT.
[Banks '88, Polchinski textbook, Ooguri-Harlow '18]
 - **But** when it comes to constraining low-energy physics...

- In QG, we expect there are **no global symmetries.**
[Wheeler in the 50's??]
- Lots of **evidence**, proven in worldsheet & AdS/CFT.
[Banks '88, Polchinski textbook, Ooguri-Harlow '18]
- **But** when it comes to constraining low-energy physics...

...its more
like this...



- In QG, we expect there are **no global symmetries.**
[Wheeler in the 50's??]
- Lots of **evidence**, proven in worldsheet & AdS/CFT.
[Banks '88, Polchinski textbook, Ooguri-Harlow '18]
- **But** when it comes to constraining low-energy physics...

...its more
like this...



...than
like this



- In QG, we expect there are **no global symmetries.**
[Wheeler in the 50's??]
- Lots of **evidence**, proven in worldsheet & AdS/CFT.
[Banks '88, Polchinski textbook, Ooguri-Harlow '18]
- **But** when it comes to constraining low-energy physics...

...its more
like this...



...than
like this



Reason: Doesn't tell us how strongly the
symmetry is broken

[See however Fichet-Saraswat, Daus-Hebecker-Leondhart-MarchRussell '20]

THE GIST OF THIS TALK

[MM, Cumrun Vafa, to appear]

THE GIST OF THIS TALK

[MM, Cumrun Vafa, to appear]



No global symmetries

THE GIST OF THIS TALK

[MM, Cumrun Vafa, to appear]



[McNamara & Vafa '19]

$$\Omega_{\text{QG}} = 0$$

THE GIST OF THIS TALK

[MM, Cumrun Vafa, to appear]



[McNamara & Vafa '19]

$$\Omega_{\text{QG}} = 0$$

+

+

THE GIST OF THIS TALK

[MM, Cumrun Vafa, to appear]



[McNamara & Vafa '19]

$$\Omega_{\text{QG}} = 0$$

+

+



Gauge anomaly

THE GIST OF THIS TALK


[MM, Cumrun Vafa, to appear]



[McNamara & Vafa '19]

$$\Omega_{\text{QG}} = 0$$

+



+

Gauge anomaly

=

=



Swampland (for the low-energy EFT!)

THE GIST OF THIS TALK

[MM, Cumrun Vafa, to appear]




[McNamara & Vafa '19]

$$\Omega_{\text{QG}} = 0$$

+

+



Gauge anomaly

=

=



Swampland (for the low-energy EFT!)

[Similar ideas: Kim-Shiu-Vafa '19, Kim-Tarazi-Vafa '19, Katz-Kim-Tarazi-Vafa '20]

THE GIST OF THIS TALK

So the **plan** is:

THE GIST OF THIS TALK

So the **plan** is:

- Review of cobordism conjecture
-

THE GIST OF THIS TALK

So the **plan** is:

- Review of cobordism conjecture
- Application: **Ifolds in 8d, 9d, and constraints on gauge group**

THE GIST OF THIS TALK

So the **plan** is:

- Review of cobordism conjecture
 - Application: **Ifolds in 8d, 9d, and constraints on gauge group**
 - Constraints in 8d, 7d, conclusions
-

THE GIST OF THIS TALK

So the **plan** is:

- Review of cobordism conjecture
- Application: **Ifolds in 8d, 9d, and constraints on gauge group**
- Constraints in 8d, 7d, conclusions

Please interrupt me!

-
- We are used to **topological symmetries** in field theory.

$$Q_m = \int F$$


(these must also be broken/gauged in gravity!)

- In a theory with dynamical gravity, there can also be **topological symmetries involving geometry**
-

For instance, take 5d gravity and on a $t=0$ slice




For instance, take 5d gravity and on a $t=0$ slice



and glue a K3!

For instance, take 5d gravity and on a $t=0$ slice




and glue a K3!

Then

$$\int p_1 = \int_{K3} \text{tr}(R^2)$$

is a conserved
topological **global**
charge in supergravity

- p_1 is actually a **bordism invariant**



$$p_1(K3) \sim p_1(M_4)$$

- More generally, these new charges are classified by **bordism groups**

e.g.


$$\Omega_4^{\text{Spin}} = \mathbb{Z}$$

or


$$\Omega_2^{\text{Pin}^-} = \mathbb{Z}_8$$

-
- In order to **kill** this charge, we need to be able to “shrink K3 to a point” in the full theory:

- In order to **kill** this charge, we need to be able to “shrink K3 to a point” in the full theory:



- In order to **kill** this charge, we need to be able to “shrink K3 to a point” in the full theory:



- In order to **kill** this charge, we need to be able to “shrink K3 to a point” in the full theory:



- In order to **kill** this charge, we need to be able to “shrink K3 to a point” in the full theory:



Killing the global symmetry forces the introduction of
defects

- In order to **kill** this charge, we need to be able to “shrink K3 to a point” in the full theory:



Killing the global symmetry forces the introduction of
defects

- In order to **kill** this charge, we need to be able to “shrink K3 to a point” in the full theory:



Killing the global symmetry forces the introduction of
defects

These defects can be potentially **anomalous**

If cobordisms predict an

If cobordisms predict an
anomalous object

If cobordisms predict an

anomalous object

in some low-energy EFT, and the anomaly
cannot be cancelled somehow

If cobordisms predict an

anomalous object

in some low-energy EFT, and the anomaly
cannot be cancelled somehow

then the theory is in the **Swamp**

If cobordisms predict an

anomalous object

in some low-energy EFT, and the anomaly
cannot be cancelled somehow

then the theory is in the **Swamp**

We will use this to get **new constraints** on theories with 16
supercharges, in 9,8,7d.

-
- In 10d, 9d, 8d, and 7d, minimal susy is 16 supercharges.

-
- In 10d, 9d, 8d, and 7d, minimal susy is 16 supercharges.
 - There are just **two multiplets**
 - Gravity: g, A_i, B, ϕ
 - Gauge: A, ϕ_i

- In 10d, 9d, 8d, and 7d, minimal susy is 16 supercharges.
- There are just **two multiplets**
 - Gravity: g, A_i, B, ϕ
 - Gauge: A, ϕ_i
- Low-energy interactions are **completely fixed** by susy.
- There is a **Narain** moduli space parametrized by the scalars

$$\frac{SO(10-d, r)}{SO(10-d) \times SO(r)}$$

-
- On a **generic point** in moduli space, the group is $U(l)^{r+d}$

-
- On a **generic point** in moduli space, the group is $U(1)^{r+d}$
 - At some points in moduli space, **symmetry enhancement** can occur.

-
- On a **generic point** in moduli space, the group is $U(1)^{r+d}$
 - At some points in moduli space, **symmetry enhancement** can occur.
 - So a sugra is **fully specified** by the **rank** of the gauge group...

- On a **generic point** in moduli space, the group is $U(1)^{r+d}$
- At some points in moduli space, **symmetry enhancement** can occur.
- So a sugra is **fully specified** by the **rank** of the gauge group...

...and a choice of **Lie algebra** (at special points in moduli space)

- On a **generic point** in moduli space, the group is $U(1)^{r+d}$
- At some points in moduli space, **symmetry enhancement** can occur.
- So a sugra is **fully specified** by the **rank** of the gauge group...
...and a choice of **Lie algebra** (at special points in moduli space)
- The **Swampland** is just figuring out which combinations have a consistent UV embedding!

- On a **generic point** in moduli space, the group is $U(1)^{r+d}$
- At some points in moduli space, **symmetry enhancement** can occur.
- So a sugra is **fully specified** by the **rank** of the gauge group...
...and a choice of **Lie algebra** (at special points in moduli space)
- The **Swampland** is just figuring out which combinations have a consistent UV embedding!
- In 10d, only 4 possibilities:

$$E_8 \times E_8, \text{Spin}(32)/\mathbb{Z}_2, E_8 \times U(1)^{248}, U(1)^{496}$$

- On a **generic point** in moduli space, the group is $U(1)^{r+d}$
- At some points in moduli space, **symmetry enhancement** can occur.
- So a sugra is **fully specified** by the **rank** of the gauge group...
...and a choice of **Lie algebra** (at special points in moduli space)
- The **Swampland** is just figuring out which combinations have a consistent UV embedding!
- In 10d, only 4 possibilities:

$$E_8 \times E_8, \text{Spin}(32)/\mathbb{Z}_2, E_8 \times U(1)^{248}, U(1)^{496}$$

We get **string universality**

[Adams-De Wolfe- Taylor '10, Kim-Shiu-Vafa '19]

-
- Let us look at a **generic point** in the Coulomb branch. Only data is the **rank**

-
- Let us look at a **generic point** in the Coulomb branch. Only data is the **rank**

$$\text{rank} \leq 26 - d \quad [\text{Kim-Tarazi-Vafa '19}]$$

(from a strong version of SDC)

-
- Let us look at a **generic point** in the Coulomb branch. Only data is the **rank**

$$\text{rank} \leq 26 - d \quad [\text{Kim-Tarazi-Vafa '19}]$$

(from a strong version of SDC)

9d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17

8d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18

7d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

- Let us look at a **generic point** in the Coulomb branch. Only data is the **rank**

$$\text{rank} \leq 26 - d \quad [\text{Kim-Tarazi-Vafa '19}]$$

(from a strong version of SDC)

9d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17

8d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18

7d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

- Let us look at a **generic point** in the Coulomb branch. Only data is the **rank**

$$\text{rank} \leq 26 - d \quad [\text{Kim-Tarazi-Vafa '19}]$$

(from a strong version of SDC)

9d 0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17
--------	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----

8d 0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
--------	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----

7d 0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
--------	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----

- Let us look at a **generic point** in the Coulomb branch. Only data is the **rank**

$$\text{rank} \leq 26 - d \quad [\text{Kim-Tarazi-Vafa '19}]$$

(from a strong version of SDC)

9d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17

8d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18

7d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

- Let us look at a **generic point** in the Coulomb branch. Only data is the **rank**

$$\text{rank} \leq 26 - d \quad [\text{Kim-Tarazi-Vafa '19}]$$

(from a strong version of SDC)

9d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17

8d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18

7d | 0 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19

We will explain the **modulo 8** periodicity in 8d and 9d

-
- Fact: The 9d, 8d supergravity lagrangians have a **parity symmetry** anywhere on the Coulomb branch

$$A \rightarrow -A, \quad B \rightarrow -B, \quad \phi \rightarrow \phi$$

-
- Fact: The 9d, 8d supergravity lagrangians have a **parity symmetry** anywhere on the Coulomb branch

$$A \rightarrow -A, \quad B \rightarrow -B, \quad \phi \rightarrow \phi$$

- Fact: This symmetry squares to (-1) on fermions. It provides a **Pin-structure**.

-
- Fact: The 9d, 8d supergravity lagrangians have a **parity symmetry** anywhere on the Coulomb branch

$$A \rightarrow -A, \quad B \rightarrow -B, \quad \phi \rightarrow \phi$$

- Fact: This symmetry squares to (-1) on fermions. It provides a **Pin-structure**.
 - Fact: In all known stringy compactifications to 8d and 9d this is an **exact symmetry**
-

-
- Fact: The 9d, 8d supergravity lagrangians have a **parity symmetry** anywhere on the Coulomb branch

$$A \rightarrow -A, \quad B \rightarrow -B, \quad \phi \rightarrow \phi$$

- Fact: This symmetry squares to (-1) on fermions. It provides a **Pin-structure**.
 - Fact: In all known stringy compactifications to 8d and 9d this is an **exact symmetry**
 - Fact: This means the theory can be consistently put on certain **non-orientable manifolds**
-

-
- Fact: The 9d, 8d supergravity lagrangians have a **parity symmetry** anywhere on the Coulomb branch

$$A \rightarrow -A, \quad B \rightarrow -B, \quad \phi \rightarrow \phi$$

- Fact: This symmetry squares to (-1) on fermions. It provides a **Pin-structure**.
- Fact: In all known stringy compactifications to 8d and 9d this is an **exact symmetry**
- Fact: This means the theory can be consistently put on certain **non-orientable manifolds**

Assumption: Every $N=1$ theory in 8d, 9d makes sense on non-orientable manifolds

-
- If the theory makes sense on Pin- manifolds, we can put it on \mathbb{RP}^2

-
- If the theory makes sense on Pin- manifolds, we can put it on RP²
 - But RP² carries a global discrete charge valued in

$$\Omega_2^{\text{Pin}^-} = \mathbb{Z}_8$$

-
- If the theory makes sense on Pin- manifolds, we can put it on RP²
 - But RP² carries a global discrete charge valued in

$$\Omega_2^{\text{Pin}^-} = \mathbb{Z}_8$$

- Cancelling this charge requires the existence of **defects** that look like

$$\mathbb{R}^3/\mathbb{Z}_2$$

- If the theory makes sense on Pin- manifolds, we can put it on RP²
- But RP² carries a global discrete charge valued in

$$\Omega_2^{\text{Pin}^-} = \mathbb{Z}_8$$

- Cancelling this charge requires the existence of **defects** that look like

$$\mathbb{R}^3/\mathbb{Z}_2$$

These are **orbifolds** or **orientifolds** in examples

We call them **I-folds**

-
- Existence of Ifolds means that **it is consistent to compactify the 9d theory on**

$$T^3/\mathbb{Z}_2$$

-
- Existence of Ifolds means that **it is consistent to compactify the 9d theory on**

$$T^3/\mathbb{Z}_2$$

- Each fixed point contributes the same (Subtlety involving tadpole cancellation)
-

-
- Existence of Ifolds means that **it is consistent to compactify the 9d theory on**

$$T^3/\mathbb{Z}_2$$

- Each fixed point contributes the same (Subtlety involving tadpole cancellation)
- This is a 6d (1,0) model. Gravitational anomaly cancellation demands

$$273 = n_T + n_H - n_V$$

$$n_T = 8n_I^T + 1, \quad n_H = 1 + r + 8n_V^I, \quad n_V = 8n_V^I$$

$$r \equiv 1 \pmod{8}$$

-
- Similar story in 8d: Compactify on

$$T^3/\mathbb{Z}_2$$

to get a 6d theory with a **modulo 16** parity anomaly.

-
- Similar story in 8d: Compactify on

$$T^3/\mathbb{Z}_2$$

to get a 6d theory with a **modulo 16** parity anomaly.

- I-folds contribute multiple of 8 to the anomaly
-

-
- Similar story in 8d: Compactify on

$$T^3/\mathbb{Z}_2$$

to get a 6d theory with a **modulo 16** parity anomaly.

- I-folds contribute multiple of 8 to the anomaly
- Sugra fields contribute $2+r$, so

$$r \equiv 2 \pmod{8}$$

- Similar story in 8d: Compactify on

$$T^3/\mathbb{Z}_2$$

to get a 6d theory with a **modulo 16** parity anomaly.

- I-folds contribute multiple of 8 to the anomaly
- Sugra fields contribute $2+r$, so

$$r \equiv 2 \pmod{8}$$

- Possible subtlety involving a **topological Green-Schwartz mechanism** [Garcia-Etxebarria-Hayashi-Ohmori-Tachikawa-Yonekura '17]

-
- In 7d, compactification on S1/Z2 yields

$$r \equiv 1 \pmod{2}$$

so we don't explain the pattern in this case.

-
- What about the gauge algebra/group?

$9d$	$ $	\mathfrak{a}_k	\mathfrak{b}_k	\mathfrak{c}_k	\mathfrak{d}_k	\mathfrak{e}_6	\mathfrak{e}_7	\mathfrak{e}_8	\mathfrak{f}_4	\mathfrak{g}_2
------	-----	------------------	------------------	------------------	------------------	------------------	------------------	------------------	------------------	------------------

$8d$	$ $	\mathfrak{a}_k	\mathfrak{b}_k	\mathfrak{c}_k	\mathfrak{d}_k	\mathfrak{e}_6	\mathfrak{e}_7	\mathfrak{e}_8	\mathfrak{f}_4	\mathfrak{g}_2
------	-----	------------------	------------------	------------------	------------------	------------------	------------------	------------------	------------------	------------------

- What about the gauge algebra/group?

$9d$ | \mathfrak{a}_k \mathfrak{b}_k \mathfrak{c}_k \mathfrak{d}_k \mathfrak{e}_6 \mathfrak{e}_7 \mathfrak{e}_8 \mathfrak{f}_4 \mathfrak{g}_2

$8d$ | \mathfrak{a}_k \mathfrak{b}_k \mathfrak{c}_k \mathfrak{d}_k \mathfrak{e}_6 \mathfrak{e}_7 \mathfrak{e}_8 \mathfrak{f}_4 \mathfrak{g}_2

- What about the gauge algebra/group?

$$9d \mid \textcircled{a}_k \quad \cancel{\textcircled{b}_k} \quad \textcircled{c}_k \quad \textcircled{d}_k \quad \textcircled{e}_6 \quad \textcircled{e}_7 \quad \textcircled{e}_8 \quad \cancel{\textcircled{f}_4} \quad \mathfrak{g}_2$$

$$8d \mid \textcircled{a}_k \quad \cancel{\textcircled{b}_k} \quad \textcircled{c}_k \quad \textcircled{d}_k \quad \textcircled{e}_6 \quad \textcircled{e}_7 \quad \textcircled{e}_8 \quad \cancel{\textcircled{f}_4} \quad \mathfrak{g}_2$$

[Garcia-Etxebarria-Hayashi-Ohmori-Tachikawa-Yonekura '17]

- What about the gauge algebra/group?

$9d \mid \textcircled{a}_k \times \textcircled{b}_k \textcircled{c}_k \textcircled{d}_k \textcircled{e}_6 \textcircled{e}_7 \textcircled{e}_8 \times \textcircled{f}_4 \textcircled{g}_2$

$8d \mid \textcircled{a}_k \times \textcircled{b}_k \textcircled{c}_k \textcircled{d}_k \textcircled{e}_6 \textcircled{e}_7 \textcircled{e}_8 \times \textcircled{f}_4 \textcircled{g}_2$

[Garcia-Etxebarria-Hayashi-Ohmori-Tachikawa-Yonekura '17]

- Just need to explain G2 and Sp(n) in 9d and G2 in 8d to claim **string universality**

- What about the gauge algebra/group?

$$9d \mid \textcircled{a}_k \quad \cancel{\textcircled{b}_k} \quad \textcircled{c}_k \quad \textcircled{d}_k \quad \textcircled{e}_6 \quad \textcircled{e}_7 \quad \textcircled{e}_8 \quad \cancel{\textcircled{f}_4} \quad \mathfrak{g}_2$$

$$8d \mid \textcircled{a}_k \quad \cancel{\textcircled{b}_k} \quad \textcircled{c}_k \quad \textcircled{d}_k \quad \textcircled{e}_6 \quad \textcircled{e}_7 \quad \textcircled{e}_8 \quad \cancel{\textcircled{f}_4} \quad \mathfrak{g}_2$$

[Garcia-Etxebarria-Hayashi-Ohmori-Tachikawa-Yonekura '17]

- Just need to explain G2 and Sp(n) in 9d and G2 in 8d to claim **string universality**
- Still, there are constraints on the global form of the gauge group.

- These come from applying cobordism conjecture to (gauge transformation+parity)

Algebra	$\dim(\mathfrak{g}) + \text{rank}(\mathfrak{g})$	$\neq 0 \bmod 8?$	Group	Real reps?
A_k	$k^2 + 3k$	$k \neq 0, 5 \bmod 8$	$SU(k+1)$	\times
			$PSU(k+1)$	\checkmark
B_k	$2k(2k+1)$	$k \equiv 1, 2 \bmod 4$	$Spin(2k+1)$	\times
			$SO(2k+1)$	\checkmark
C_k	$2k(2k+1)$	$k \equiv 1, 2 \bmod 4$	$Sp(k)$	\times
			$Sp(k)/\mathbb{Z}_2$	\checkmark
D_k	$2k^2$	$k \text{ odd}$	$Spin(2k)$	\times
			$SO(2k)$	\checkmark
			$Spin(2k)/\mathbb{Z}_4$	\checkmark
E_6	84	\checkmark	E_6	\times
			E_6/\mathbb{Z}_3	\checkmark
E_7	140	\checkmark	E_7	\times
			E_7/\mathbb{Z}_2	\checkmark
E_8	256	\times	E_8	\checkmark
F_4	56	\times	F_4	\checkmark
G_2	16	\times	G_2	\checkmark

TO WRAP UP

- **Cobordism + anomalies = Swampland**
 - Established **Coulomb branch string universality** in 8d and 9d.
 - Apply these ideas to **models with less susy.**
 - 6d (1,0)
 - SM?
-

Thank you!

-
- Bonus track: Tadpoles in T3/Z2
 - The orbifold T3/Z2 can carry monopole charge under 9d gauge fields. **If non-integer, it will spoil the argument**
 - Integrality can be determined from Dirac quantization and anomaly of dual particle (BPS with 8d supercharges).

$$\eta(\mathbb{R}\mathbb{P}^2) = \frac{1}{8} \cdot 8$$

- Argument fails for higher codimension l-folds
-