

# Non-flat elliptic fourfolds, three-form cohomology and 4D strongly coupled theories

Paul-Konstantin Oehlmann

Uppsala University

Based on **arXiv:2102.10722**

String Pheno Seminar Series 2021, Harvard, April 6th. 2021

# Hodge diamonds of Calabi-Yau Geometries

$$\begin{array}{ccccccc}
 & & & & 1 & & \\
 & & & 0 & & 0 & \\
 & & 0 & & h^{1,1} & & 0 \\
 & 0 & & & & & 0 \\
 1 & & h^{3,1} & & & & h^{3,1} & 1 \\
 & 0 & & & & & 0 & \\
 & & 0 & & h^{1,1} & & 0 & \\
 & & 0 & & 0 & & 0 & \\
 & & & & 1 & & & 
 \end{array}$$

## Hodge numbers in fourfolds

Fourfolds admit **Kähler** and **complex structure moduli** counted by  $h^{1,1}$  and  $h^{3,1}$


# Hodge diamonds of Calabi-Yau Geometries

$$\begin{array}{ccccccc}
 & & & & 1 & & \\
 & & & & 0 & & \\
 & & 0 & & h^{1,1} & & 0 \\
 & 0 & 0 & h^{2,1} & h^{2,1} & 0 & \\
 1 & & h^{3,1} & h^{2,2} & h^{3,1} & & 1 \\
 & 0 & h^{2,1} & h^{2,1} & 0 & & \\
 & & 0 & h^{1,1} & 0 & & \\
 & & 0 & & 0 & & \\
 & & & & 1 & & 
 \end{array}$$

## Hodge numbers in fourfolds

Fourfolds admit **Kähler** and **complex structure moduli** counted by  $h^{1,1}$  and  $h^{3,1}$   
**Fourfolds** admit one additional **independent** Hodge number  $h^{2,1}$


# Hodge diamonds of Calabi-Yau Geometries



## Hodge numbers in fourfolds

Fourfolds admit **Kähler** and **complex structure moduli** counted by  $h^{1,1}$  and  $h^{3,1}$   
**Fourfolds** admit one additional **independent** Hodge number  $h^{2,1}$

# Hodge diamonds of Calabi-Yau Geometries




## Hodge numbers in fourfolds

Fourfolds admit **Kähler** and **complex structure moduli** counted by  $h^{1,1}$  and  $h^{3,1}$   
**Fourfolds** admit one additional **independent** Hodge number  $h^{2,1}$

- Kähler and complex structure **interchanged** by **mirror symmetry**

# Hodge diamonds of Calabi-Yau Geometries



## Hodge numbers in fourfolds

Fourfolds admit **Kähler** and **complex structure moduli** counted by  $h^{1,1}$  and  $h^{3,1}$

**Fourfolds** admit one additional **independent** Hodge number  $h^{2,1}$

- Kähler and complex structure **interchanged** by **mirror symmetry**
- $h^{2,1}$  is **self-mirror** → **deserves some attention** [Greiner, Grimm'15]

## $h^{2,1}$ in M/F-theory

**How do**  $h^{2,1}$  threeforms contribute in **M/F-theory** on such fourfolds  $X_4$ ?

## $h^{2,1}$ in M/F-theory

**How do  $h^{2,1}$  threeforms contribute in M/F-theory on such fourfolds  $X_4$ ?**

- **3D, M-theory on  $X_4$ :** expansion of  $C_3$  **threeform** yields **additional massless singlets**  $N^I C_3 \rightarrow N^I \eta_I^{(3)} + \dots$ ,  $I = 1 \dots h^{2,1}(X_4)$



# $h^{2,1}$ in M/F-theory

**How do  $h^{2,1}$  threeforms contribute in M/F-theory on such fourfolds  $X_4$ ?**

- **3D, M-theory on  $X_4$ :** expansion of  $C_3$  **threeform** yields **additional massless singlets**  $N^I C_3 \rightarrow N^I \eta_I^{(3)} + \dots$ ,  $I = 1 \dots h^{2,1}(X_4)$
- **4D F-theory lift**, if fourfold is **elliptic**, **splits up**  $h^{2,1}(X_4) = h^{2,1}(B_3) + h_{\text{fiber}}^{2,1}$ 
  - **Base contribution:**  $h^{2,1}(B_3)$  dualize to 4D RR **U(1)**'s
  - **Fiber contribution:**  $h_{\text{fiber}}^{2,1}$  stay 4D chiral **singlets**

# $h^{2,1}$ in M/F-theory

**How do  $h^{2,1}$  threeforms contribute in M/F-theory on such fourfolds  $X_4$ ?**

- **3D, M-theory on  $X_4$ :** expansion of  $C_3$  **threeform** yields **additional massless singlets**  $N^I C_3 \rightarrow N^I \eta_I^{(3)} + \dots$ ,  $I = 1 \dots h^{2,1}(X_4)$
- **4D F-theory lift, if fourfold is elliptic, splits up**  $h^{2,1}(X_4) = h^{2,1}(B_3) + h_{\text{fiber}}^{2,1}$ 
  - **Base contribution:**  $h^{2,1}(B_3)$  dualize to 4D RR  $U(1)$ 's
  - **Fiber contribution:**  $h_{\text{fiber}}^{2,1}$  stay 4D chiral singlets

The  $h_{\text{fiber}}^{2,1}$  singlets **mix with Kahler moduli and admit special shift symmetries** → **deserves some attention** [Grimm; Grimm, Taylor'12]

# $h^{2,1}$ in M/F-theory

**How do  $h^{2,1}$  threeforms contribute in M/F-theory on such fourfolds  $X_4$ ?**

- **3D, M-theory on  $X_4$ :** expansion of  $C_3$  **threeform** yields **additional massless singlets**  $N^I C_3 \rightarrow N^I \eta_I^{(3)} + \dots$ ,  $I = 1 \dots h^{2,1}(X_4)$
- **4D F-theory lift**, if fourfold is **elliptic**, **splits up**  $h^{2,1}(X_4) = h^{2,1}(B_3) + h_{\text{fiber}}^{2,1}$ 
  - **Base contribution:**  $h^{2,1}(B_3)$  dualize to 4D RR  $U(1)$ 's
  - **Fiber contribution:**  $h_{\text{fiber}}^{2,1}$  stay 4D chiral **singlets**

The  $h_{\text{fiber}}^{2,1}$  singlets **mix with Kahler moduli** and **admit special shift symmetries** → **deserves some attention** [Grimm; Grimm, Taylor'12]

**Q:** How to engineer  $h_{\text{fiber}}^{2,1}$  ?

# $h^{2,1}$ in M/F-theory

**How do  $h^{2,1}$  threeforms contribute in M/F-theory on such fourfolds  $X_4$ ?**


- **3D, M-theory on  $X_4$ :** expansion of  $C_3$  **threeform** yields **additional massless singlets**  $N^I C_3 \rightarrow N^I \eta_I^{(3)} + \dots$ ,  $I = 1 \dots h^{2,1}(X_4)$
- **4D F-theory lift**, if fourfold is **elliptic**, **splits up**  $h^{2,1}(X_4) = h^{2,1}(B_3) + h_{\text{fiber}}^{2,1}$ 
  - **Base contribution:**  $h^{2,1}(B_3)$  dualize to 4D RR  $U(1)$ 's
  - **Fiber contribution:**  $h_{\text{fiber}}^{2,1}$  stay 4D chiral **singlets**

The  $h_{\text{fiber}}^{2,1}$  singlets **mix with Kahler moduli** and **admit special shift symmetries** → **deserves some attention** [Grimm; Grimm, Taylor'12]

**Q:** How to engineer  $h_{\text{fiber}}^{2,1}$  ?

**A:** Compactifications of certain 6D SCFT sectors


# Resolved elliptic threefolds



Consider an **elliptic threefold** with  $G_1$  and  $G_2$  group type of **singularities**,

- $G_i$  **fibers** can be **resolved** in a **crepant** way


# Resolved elliptic threefolds



Consider an **elliptic threefold** with  $G_1$  and  $G_2$  group type of **singularities**,

- $G_i$  **fibers** can be **resolved** in a **crepant** way


# Resolved elliptic threefolds



Consider an **elliptic threefold** with  $G_1$  and  $G_2$  group type of **singularities**,

- $G_i$  **fibers** can be **resolved** in a **crepant** way
- The  $[G_1-G_2]$  **intersection** might be **non-minimal**  $\rightarrow$  **6D SCFT** !

# Resolved elliptic threefolds




Consider an **elliptic threefold** with  $G_1$  and  $G_2$  group type of **singularities**,

- $G_i$  **fibers** can be **resolved** in a **crepant** way
- The  $[G_1-G_2]$  **intersection** might be **non-minimal**  $\rightarrow$  **6D SCFT** !

**Two (crepant) ways to resolve the threefold**




# Resolution via base blow-up



Consider an **elliptic threefold** with  $[G_1-G_2]$  **collision**

- **T Blow-ups of the base** until the fiber is regular


# Resolution via base blow-up



Consider an **elliptic threefold** with  $[G_1-G_2]$  **collision**

- **T Blow-ups of the base** until the fiber is regular
- **Resolve all fibers** requires  $\dim(\sum_i \text{rank}(\mathfrak{g}_i)) = R$
- **Associated 6D SCFT** admits **T dimensional tensor branch** [del Zotto, Heckman, Tomasiello, Vafa '14]


# Resolution via base blow-up



Consider an **elliptic threefold** with  $[G_1-G_2]$  **collision**

- **T Blow-ups of the base** until the fiber is regular
- **Resolve all fibers** requires  $\dim(\sum_i \text{rank}(\mathfrak{g}_i)) = R$
- **Associated 6D SCFT** admits **T dimensional tensor branch** [del Zotto, Heckman, Tomasiello, Vafa '14]
- **CB** =  $T + R$  dimensional 5D **Coulomb branch**

# Resolution via base blow-up




Consider an **elliptic threefold** with  $[G_1-G_2]$  **collision**

- **T Blow-ups of the base** until the fiber is regular
- **Resolve all fibers** requires  $\dim(\sum_i \text{rank}(\mathfrak{g}_i)) = R$
- **Associated 6D SCFT** admits **T dimensional tensor branch** [del Zotto, Heckman, Tomasiello, Vafa '14]
- **CB** =  $T + R$  dimensional 5D **Coulomb branch**


There exists an **alternative resolution**

# Non-flat resolutions




Non-minimal singularities admits a **non-flat resolution** [Schafer-Nameki, Lawrie '12; Dierigl, Oehlmann, Ruehle '18]

# Non-flat resolutions



Non-minimal singularities admits a **non-flat resolution** [Schafer-Nameki, Lawrie '12; Dierigl, Oehlmann, Ruehle '18]


# Non-flat resolutions



Non-minimal singularities admits a **non-flat resolution** [Schafer-Nameki, Lawrie '12; Dierigl, Oehlmann, Ruehle '18]

- Replace fiber with **several surfaces**  $\dim_{\mathbb{C}}(E_i) = 2$

# Non-flat resolutions




Non-minimal singularities admits a **non-flat resolution** [Schafer-Nameki, Lawrie '12; Dierigl, Oehlmann, Ruehle '18]

- Replace fiber with **several surfaces**  $\dim_{\mathbb{C}}(E_i) = 2$




# Non-flat resolutions



Non-minimal singularities admits a **non-flat resolution** [Schafer-Nameki, Lawrie '12; Dierigl, Oehlmann, Ruehle '18]

- Replace fiber with **several surfaces**  $\dim_{\mathbb{C}}(E_i) = 2$


# Non-flat resolutions



Non-minimal singularities admits a **non-flat resolution** [Schafer-Nameki, Lawrie '12; Dierigl, Oehlmann, Ruehle '18]

- Replace fiber with **several surfaces**  $\dim_{\mathbb{C}}(E_i) = 2$


# Non-flat resolutions



Non-minimal singularities admits a **non-flat resolution** [Schafer-Nameki, Lawrie'12; Dierigl, Oehlmann, Ruehle'18]


- Replace fiber with **several surfaces**  $\dim_{\mathbb{C}}(E_i) = 2$
- Also needs **CB non-flat surfaces!**
- **Consistent** with 5D **M-theory duality** of **6D SCFT** compactifications [Aruzzi, Lin, Mayrhofer; Aruzzi, Lin, Schafer-Nameki, Wang, Lawrie, Hubner'18,19]

# Phases of non-minimal singularities



**Both** resolutions part of **extended Kahler cone** of the threefold


# Phases of non-minimal singularities



**Both** resolutions part of **extended Kahler cone** of the threefold

**Non-flatness** describes **resolved** non-minimal singularities **without changing the base**


# From three-to-fourfolds



**Extend the base** from  $B_2 \rightarrow B_3$ : threefold  $\rightarrow$  fourfold  $X_4$

- The fiber **singularity** appears over **Riemann surfaces**  $C_{\alpha} \in B_3$  of genus  $g_{\alpha}$


# From three-to-fourfolds



**Extend the base** from  $B_2 \rightarrow B_3$ : threefold  $\rightarrow$  fourfold  $X_4$

- The fiber **singularity** appears **over Riemann surfaces**  $C_\alpha \in B_3$  of genus  $g_\alpha$
- Fiber **resolution** via surfaces  $E_i$  **base independent**

# From three-to-fourfolds




**Extend the base** from  $B_2 \rightarrow B_3$ : threefold  $\rightarrow$  fourfold  $X_4$

- The fiber **singularity** appears **over Riemann surfaces**  $C_\alpha \in B_3$  of genus  $g_\alpha$
- Fiber **resolution** via surfaces  $E_i$  **base independent**
- Gives rise to divisor  $D_{i,\alpha}$  that itself admits a **fibration structure**



## From three-to-fourfolds




**Extend the base** from  $B_2 \rightarrow B_3$ : threefold  $\rightarrow$  fourfold  $X_4$

- The fiber **singularity** appears **over Riemann surfaces**  $C_\alpha \in B_3$  of genus  $g_\alpha$
- Fiber **resolution** via surfaces  $E_i$  **base independent**
- Gives rise to divisor  $D_{i,\alpha}$  that itself admits a **fibration structure**
- **Cohomology** of  $D_{i,\alpha}$  from **Leray-Hirsch spectral sequence**:

$$h^{0,0}(D_{i,\alpha}) = h^{0,0}(C_\alpha) \cdot h^{0,0}(E_{i,\alpha}) = 1$$

$$h^{1,0}(D_{i,\alpha}) = h^{1,0}(C_\alpha) \cdot h^{0,0}(E_{i,\alpha}) = g_\alpha$$


# Non-flat fourfold cohomology



$$\begin{array}{c}
 E_{i,\alpha} \rightarrow D_{i,\alpha} \\
 \downarrow \pi \\
 \mathcal{C}_\alpha
 \end{array}$$

- $D_{i,\alpha}$  **embedded** into fourfold  $X_4$  via **Gysin map**  $\iota$  [Greiner, Grimm'17]

# Non-flat fourfold cohomology




$$\begin{array}{ccc}
 E_{i,\alpha} & \rightarrow & D_{i,\alpha} \\
 & & \downarrow \pi \\
 & & \mathcal{C}_\alpha
 \end{array}$$

- $D_{i,\alpha}$  **embedded** into fourfold  $X_4$  via **Gysin map**  $\iota$  [Greiner, Grimm'17]
- Compute **cohomology** contribution from **Gysin complex** in  $X_4$  [Mavlyutov'00]

$$\oplus_{i,\alpha} H^{0,0}(D_{i,\alpha}) \xrightarrow{\iota} H^{1,1}(X_4)$$

$$\oplus_{i,\alpha} H^{1,0}(D_{i,\alpha}) \xrightarrow{\iota} H^{2,1}(X_4)$$

# Non-flat fourfold cohomology



$$E_{i,\alpha} \rightarrow D_{i,\alpha} \\ \downarrow \pi \\ C_\alpha$$

- $D_{i,\alpha}$  **embedded** into fourfold  $X_4$  via **Gysin map**  $\iota$  [Greiner, Grimm'17]
- Compute **cohomology** contribution from **Gysin complex** in  $X_4$  [Mavlyutov'00]

$$\oplus_{i,\alpha} H^{0,0}(D_{i,\alpha}) \xrightarrow{\iota} H^{1,1}(X_4)$$


$$\oplus_{i,\alpha} H^{1,0}(D_{i,\alpha}) \xrightarrow{\iota} H^{2,1}(X_4)$$

- **Sum** over all  $CB_\alpha$  surfaces  $E_i$  over each base curve  $C_\alpha$

$$h^{1,1}(X_4)_{\text{non-flat}} = \sum_{\alpha} CB_\alpha$$

$$h^{2,1}(X_4)_{\text{non-flat}} = \sum_{\alpha} g_\alpha \cdot CB_\alpha$$

# Non-flat fourfold cohomology



$$E_{i,\alpha} \rightarrow D_{i,\alpha} \downarrow \pi C_\alpha$$

- $D_{i,\alpha}$  **embedded** into fourfold  $X_4$  via **Gysin map**  $\iota$  [Greiner, Grimm'17]
- Compute **cohomology** contribution from **Gysin complex** in  $X_4$  [Mavlyutov'00]

$$\oplus_{i,\alpha} H^{0,0}(D_{i,\alpha}) \xrightarrow{\iota} H^{1,1}(X_4)$$

$$\oplus_{i,\alpha} H^{1,0}(D_{i,\alpha}) \xrightarrow{\iota} H^{2,1}(X_4)$$

- **Sum** over all  $CB_\alpha$  surfaces  $E_i$  over each base curve  $C_\alpha$

$$h^{1,1}(X_4)_{\text{non-flat}} = \sum_{\alpha} CB_\alpha$$

$$h_{\text{fiber}}^{2,1} \ni h^{2,1}(X_4)_{\text{non-flat}} = \sum_{\alpha} g_\alpha \cdot CB_\alpha$$

# Summary

**Double check** in toric examples via **Batyrev construction** [Klemm, Lian, Roan, Yau '97]

- **E-string theories** on genus  $g = 153$  curves ✓
- **Higher rank theories** e.g.  $[E_8 \times SU(n)]$  superconformal matter over  $g \gg 1$  curves ✓

# Summary

**Double check** in toric examples via **Batyrev construction** [Klemm, Lian, Roan, Yau '97]

- **E-string theories** on genus  $g = 153$  curves ✓
- **Higher rank theories** e.g.  $[E_8 \times SU(n)]$  superconformal matter over  $g \gg 1$  curves ✓

Consider **conifold type** transitions in fourfolds that **remove such non-flat fibers**

- 1 Removal via **Higgs branch type** of transitions
- 2 Removal via **Coulomb branch type** of transitions

# Summary

**Double check** in toric examples via **Batyrev construction** [Klemm, Lian, Roan, Yau '97]

- **E-string theories** on genus  $g = 153$  curves ✓
- **Higher rank theories** e.g.  $[E_8 \times SU(n)]$  superconformal matter over  $g \gg 1$  curves ✓


Consider **conifold type** transitions in fourfolds that **remove such non-flat fibers**

- 1 Removal via **Higgs branch type** of transitions
- 2 Removal via **Coulomb branch type** of transitions

**Disclaimer:** Fourfold examples  $X_4$  is **compact** and **no  $G_4$  flux!**




# Example: $E_6 \times U(1)$ in 6D



**Start in 6D:** Construct threefold with  $E_6 \times U(1)$  gauge group and an **E-string**


# Example: $E_6 \times U(1)$ in 6D



**Start in 6D:** Construct threefold with  $E_6 \times U(1)$  gauge group and an **E-string**

- **Phase 1: Base blow-up** does not change hodge numbers  $\Delta h^{i,j}(X_3) = 0$


# Example: $E_6 \times U(1)$ in 6D



**Start in 6D:** Construct threefold with  $E_6 \times U(1)$  gauge group and an **E-string**

- **Phase 1: Base blow-up** does not change hodge numbers  $\Delta h^{i,j}(X_3) = 0$
- **Phase 2: Deform** the E-string into  $(27_{-\frac{1}{3}} + 1_1)$  hypermultiplets
- **Enforced by 6D anomalies** [Dierigl, Oehlmann, Ruehle'18]

# Example: $E_6 \times U(1)$ in 6D




**Start in 6D:** Construct threefold with  $E_6 \times U(1)$  gauge group and an **E-string**

- **Phase 1: Base blow-up** does not change hodge numbers  $\Delta h^{i,j}(X_3) = 0$
- **Phase 2: Deform** the E-string into  $(27_{-\frac{1}{3}} + 1_1)$  hypermultiplets
- **Enforced by 6D anomalies** [Dierigl, Oehlmann, Ruehle'18]

**Perform the same transition in a fourfold**


# Example: $E_6 \times U(1)$ in 4D



Place **E-string type** fiber is over Riemann surface of e.g.  $g = 5$


- **Phase 1: Deform** the E-string curve into  $(27_{-\frac{1}{3}} + 1_1)$  curves
- Example: Cohomology change:  $\Delta h^{1,1} = -1, \Delta h^{2,1} = -5$  ✓

# The $E_6 \times U(1)$ deformed fourfold




The  $27_{-\frac{1}{3}}$  and  $1_1$  curves **re-intersect** in **codimension 3** non-flat point

# The $E_6 \times U(1)$ deformed fourfold



The  $27_{-\frac{1}{3}}$  and  $1_1$  curves **re-intersect** in **codimension 3** non-flat point

- This is a **non-perturbative superpotential** coupling term
- **How to see that?** further **break** the  $U(1)$  group


Further break to  $E_6$ 

**Same locus: flat fiber of  $E_8$  bouquet type**

- This is a  $27^3$  triple intersection point
- **Adding  $U(1)$  over enhances the  $E_8$  point**




Adding to  $E_6 \times U(1)$  again



Adding the  $U(1)$  renders  $27_{-\frac{1}{3}}^3$  not gauge invariant anymore


# Adding to $E_6 \times U(1)$ again



Adding the  $U(1)$  renders  $27_{-\frac{1}{3}}$  **not gauge invariant** anymore

- Inserting the  $1_{-1}$  singlet, it becomes **gauge invariant 4point coupling**
- **D1 string instanton** induced coupling [Achmed-Zade, Garcia-Etxebarria, Mayrhofer'18]

Adding to  $E_6 \times U(1)$  again




Adding the  $U(1)$  renders  $27_{-\frac{1}{3}}^3$  **not gauge invariant** anymore

- Inserting the  $1_{-1}$  singlet, it becomes **gauge invariant 4point coupling**
- **D1 string instanton** induced coupling [Achmed-Zade, Garcia-Etxebarria, Mayrhofer'18]

**A remnant of the deformed E-string curve**


## Back to E-string curve



### Phase 2: Blow-ups of $B_3$ to remove E-string curve

- Successive blow-ups **split up the non-flat** curve into **several**  $\mathbb{P}^1$ 's


# Back to E-string curve



## Phase 2: Blow-ups of $B_3$ to remove E-string curve

- Successive blow-ups **split up the non-flat** curve into **several  $\mathbb{P}^1$ 's**


# Back to E-string curve



## Phase 2: Blow-ups of $B_3$ to remove E-string curve

- Successive blow-ups **split up the non-flat** curve into **several  $\mathbb{P}^1$ 's**  
Example: cohomology change:  $\Delta(h^{1,1}, h^{2,1}, h^{3,1}) = (7, -5, -13)\checkmark$
- **6 more blow-ups** of  $B_3$  required with  $\Delta h^{i,1}(X_4) = 0$

# Back to E-string curve



## Phase 2: Blow-ups of $B_3$ to remove E-string curve

- Successive blow-ups **split up the non-flat** curve into **several  $\mathbb{P}^1$ 's**  
Example: cohomology change:  $\Delta(h^{1,1}, h^{2,1}, h^{3,1}) = (7, -5, -13)\checkmark$
- **6 more blow-ups** of  $B_3$  required with  $\Delta h^{i,1}(X_4) = 0$
- **Euler number invariant:**  $\Delta\chi = 6(8 + \Delta h^{1,1} + \Delta h^{3,1} - \Delta h^{2,1}) = 0$
- **D3 tadpole** unchanged:  $\Delta \int G_4^2 = \Delta n_{d3}/12$

## Non-flat fibers in codim 2

- Resolve non-minimal **fiber singularities**, sacrificing equ-dimensionality
- **Compactified 6D conformal** matter on **on Riemann surface**
- Contributes 4D **singlets** from  $h_{\text{fiber}}^{2,1} = g \cdot CB$



## Non-flat fibers in codim 2

- **Resolve** non-minimal **fiber singularities**, **sacrificing equi-dimensionality**
- **Compactified 6D conformal** matter on **on Riemann surface**
- Contributes 4D **singlets** from  $h_{\text{fiber}}^{2,1} = g \cdot \text{CB}$

## These fibers can be removed by

- **Higgs branch** type **deformation**, push non-flatness to **points in  $B_3$** 
  - **non-perturbative 4point couplings** among e.g. E-string left-overs
- **Coulomb branch** type transitions by **blow-ups** of  $B_3$  base
  - **changes all hodge numbers** but **keeps Euler number  $\chi(X_4)$  invariant**

## Non-flat fibers in codim 2

- **Resolve** non-minimal **fiber singularities**, **sacrificing** equi-dimensionality
- **Compactified 6D conformal** matter on **on Riemann surface**
- Contributes 4D **singlets** from  $h_{\text{fiber}}^{2,1} = g \cdot CB$

## These fibers can be removed by

- **Higgs branch** type **deformation**, push non-flatness to **points in  $B_3$** 
  - **non-perturbative 4point couplings** among e.g. E-string left-overs
- **Coulomb branch** type transitions by **blow-ups** of  $B_3$  base
  - **changes all hodge numbers** but **keeps Euler number  $\chi(X_4)$  invariant**

## Outlook

- **Need** to add  $G_4$ -**flux**! Use invariance of  $\chi$  under transitions
- Non-flat fibers with **monodromy**, transitions with **higher rank theories**

## Non-flat fibers in codim 2

- **Resolve** non-minimal **fiber singularities**, **sacrificing equi-dimensionality**
- **Compactified 6D conformal** matter on **on Riemann surface**
- Contributes 4D **singlets** from  $h_{\text{fiber}}^{2,1} = g \cdot CB$

## These fibers can be removed by

- **Higgs branch** type **deformation**, push non-flatness to **points in  $B_3$** 
  - **non-perturbative 4point couplings** among e.g. E-string left-overs
- **Coulomb branch** type transitions by **blow-ups** of  $B_3$  base
  - **changes all hodge numbers** but **keeps Euler number  $\chi(X_4)$  invariant**

Thank You Very Much

## Outlook

- **Need** to add  $G_4$ -**flux**! Use invariance of  $\chi$  under transitions
- Non-flat fibers with **monodromy**, transitions with **higher rank theories**