


# Insights from Non-Ambient Flops



Callum Brodie

Institut de Physique Théorique, Université Paris Saclay

Based on 2010.06597 and 2011.xxxxx  
with Andrei Constantin and Andre Lukas

String Phenomenology Seminar Series  
November 24th 2020

# Overview

---

# Overview

---

Flop: example of topological transition between CY three-folds

# Overview

---

Flop: example of topological transition between CY three-folds  
Typically deduced from flops of ambient space

# Overview

---

Flop: example of topological transition between CY three-folds

Typically deduced from flops of ambient space

'Non-ambient flop': flop not induced by ambient space

# Overview

---

Flop: example of topological transition between CY three-folds  
Typically deduced from flops of ambient space  
'Non-ambient flop': flop not induced by ambient space

In this talk:

- How to discern the presence of non-ambient flops

# Overview

---

Flop: example of topological transition between CY three-folds

Typically deduced from flops of ambient space

'Non-ambient flop': flop not induced by ambient space

In this talk:

- How to discern the presence of non-ambient flops
- How to perform them

# Overview

---

Flop: example of topological transition between CY three-folds

Typically deduced from flops of ambient space

'Non-ambient flop': flop not induced by ambient space

In this talk:

- How to discern the presence of non-ambient flops
- How to perform them
- What one can learn from them



# Motivation

---

# Motivation

---

Why study complete intersection Calabi-Yaus (CICYs)?

# Motivation

---

Why study complete intersection Calabi-Yaus (CICYs)?

- Below techniques generalise to other important constructions: toric complete intersections, gCICYs

# Motivation

---

Why study complete intersection Calabi-Yaus (CICYs)?

- Below techniques generalise to other important constructions: toric complete intersections, gCICYs
- These three provide a tractable corner of the landscape

# Motivation

---

## Why study complete intersection Calabi-Yaus (CICYs)?

- Below techniques generalise to other important constructions: toric complete intersections, gCICYs
- These three provide a tractable corner of the landscape
- They also provide a fruitful arena for model-building

# Motivation

---

Why study complete intersection Calabi-Yaus (CICYs)?

- Below techniques generalise to other important constructions: toric complete intersections, gCICYs
- These three provide a tractable corner of the landscape
- They also provide a fruitful arena for model-building

Why analyse flops?

# Motivation

---

## Why study complete intersection Calabi-Yaus (CICYs)?

- Below techniques generalise to other important constructions: toric complete intersections, gCICYs
- These three provide a tractable corner of the landscape
- They also provide a fruitful arena for model-building

## Why analyse flops?

- Physics through transition can be smooth, so flops connect phases of the theory into a larger moduli space

# Motivation

---

## Why study complete intersection Calabi-Yaus (CICYs)?

- Below techniques generalise to other important constructions: toric complete intersections, gCICYs
- These three provide a tractable corner of the landscape
- They also provide a fruitful arena for model-building

## Why analyse flops?

- Physics through transition can be smooth, so flops connect phases of the theory into a larger moduli space
- To analyse limits of the physical moduli space, important to understand extended Kähler cone



# Motivation

---

## Why study complete intersection Calabi-Yaus (CICYs)?

- Below techniques generalise to other important constructions: toric complete intersections, gCICYs
- These three provide a tractable corner of the landscape
- They also provide a fruitful arena for model-building

## Why analyse flops?

- Physics through transition can be smooth, so flops connect phases of the theory into a larger moduli space
- To analyse limits of the physical moduli space, important to understand extended Kähler cone
- Understanding flops can allow one to immediately write down data required for model-building

# Talk goals

---

# Talk goals

---

New twist on old story (CICYs), giving:

# Talk goals

---

New twist on old story (CICYs), giving:

- **New tool:** Read off non-ambient flops from CICY data

# Talk goals

---

New twist on old story (CICYs), giving:

- **New tool:** Read off non-ambient flops from CICY data
- **Characterisation:** Almost all CICYs admit flops, almost every Kähler direction is a flop, many CICYs flop to themselves

# Talk goals

---

New twist on old story (CICYs), giving:

- **New tool**: Read off non-ambient flops from CICY data
- **Characterisation**: Almost all CICYs admit flops, almost every Kähler direction is a flop, many CICYs flop to themselves
- **New tool**: How to perform flops explicitly

# Talk goals

---

New twist on old story (CICYs), giving:

- **New tool:** Read off non-ambient flops from CICY data
- **Characterisation:** Almost all CICYs admit flops, almost every Kähler direction is a flop, many CICYs flop to themselves
- **New tool:** How to perform flops explicitly
- **Novel situation:** Many CYs with infinitely many flops, extended Kähler cones has fractal structure

# Talk goals

---

New twist on old story (CICYs), giving:

- **New tool**: Read off non-ambient flops from CICY data
- **Characterisation**: Almost all CICYs admit flops, almost every Kähler direction is a flop, many CICYs flop to themselves
- **New tool**: How to perform flops explicitly
- **Novel situation**: Many CYs with infinitely many flops, extended Kähler cones has fractal structure
- **New tool**: All line bundle cohomology from CICY data



# Flops

---

# Flops

---

Codimension-two operation, replaces curve with another


# Flops

---

Codimension-two operation, replaces curve with another

Classic example: change in triangulation of toric variety

Often descends to CY, but  
not present for CICYs



# Flops

---

Codimension-two operation, replaces curve with another

Classic example: change in triangulation of toric variety

Often descends to CY, but  
not present for CICYs



Achieved continuously by shrinking curve to zero volume giving singular points, then growing new curve ('small resolution')

# Flops

Codimension-two operation, replaces curve with another

Classic example: change in triangulation of toric variety

Often descends to CY, but not present for CICYs




Achieved continuously by shrinking curve to zero volume giving singular points, then growing new curve ('small resolution')

Isomorphism in codimension 1, so

$$H^{1,1}(X) \cong H^{1,1}(X')$$

Manifold moves to new Kähler cone,

$$\mathcal{K}(X) \rightarrow \mathcal{K}(X')$$



$$\mathcal{K}(X) \rightarrow \mathcal{K}(X') \text{ in } H^{1,1}(X) \cong H^{1,1}(X')$$

# The Atiyah flop

---

# The Atiyah flop

---

Simplest example of a flop

# The Atiyah flop

---

Simplest example of a flop

Begin with singular three-fold with local description

$$\{yz = uv\} \subset \mathbb{C}^4[y, z, u, v]$$



# The Atiyah flop

---

Simplest example of a flop

Begin with singular three-fold with local description

$$\{yz = uv\} \subset \mathbb{C}^4[y, z, u, v]$$

Singular when  $y = z = u = v = 0$

# The Atiyah flop

---

Simplest example of a flop

Begin with singular three-fold with local description

$$\{yz = uv\} \subset \mathbb{C}^4[y, z, u, v]$$

Singular when  $y = z = u = v = 0$

Resolution of singularity into  $\mathbb{P}^1$  achieved as

$$\{yx_0 = ux_1, zx_0 = vx_1\} \subset \mathbb{C}^4[y, z, u, v] \times \mathbb{P}^1[x]$$

# The Atiyah flop

---

Simplest example of a flop

Begin with singular three-fold with local description

$$\{yz = uv\} \subset \mathbb{C}^4[y, z, u, v]$$

Singular when  $y = z = u = v = 0$

Resolution of singularity into  $\mathbb{P}^1$  achieved as

$$\{yx_0 = ux_1, zx_0 = vx_1\} \subset \mathbb{C}^4[y, z, u, v] \times \mathbb{P}^1[x]$$

Single solution in  $\mathbb{P}^1$  only when  $yz = uv$

# The Atiyah flop

---

Simplest example of a flop

Begin with singular three-fold with local description

$$\{yz = uv\} \subset \mathbb{C}^4[y, z, u, v]$$

Singular when  $y = z = u = v = 0$

Resolution of singularity into  $\mathbb{P}^1$  achieved as

$$\{yx_0 = ux_1, zx_0 = vx_1\} \subset \mathbb{C}^4[y, z, u, v] \times \mathbb{P}^1[x]$$

Single solution in  $\mathbb{P}^1$  only when  $yz = uv$

Entire  $\mathbb{P}^1$  precisely when  $y = z = u = v = 0$

# The Atiyah flop

---

Simplest example of a flop

Begin with singular three-fold with local description

$$\{yz = uv\} \subset \mathbb{C}^4[y, z, u, v]$$

Singular when  $y = z = u = v = 0$

Resolution of singularity into  $\mathbb{P}^1$  achieved as

$$\{yx_0 = ux_1, zx_0 = vx_1\} \subset \mathbb{C}^4[y, z, u, v] \times \mathbb{P}^1[x]$$

Single solution in  $\mathbb{P}^1$  only when  $yz = uv$

Entire  $\mathbb{P}^1$  precisely when  $y = z = u = v = 0$

But could have chosen to pair as  $\{yx_0 = vx_1, zx_0 = ux_1\}$

These two choices give spaces related by a flop



# CICYs

---

Idea: Ambient space  $\mathcal{A}$  is simply product of projective spaces,  
CY  $X$  given by intersection of zero locus of set of polynomials

# CICYs

---

Idea: Ambient space  $\mathcal{A}$  is simply product of projective spaces, CY  $X$  given by intersection of zero locus of set of polynomials

Example:



Idea: Ambient space  $\mathcal{A}$  is simply product of projective spaces, CY  $X$  given by intersection of zero locus of set of polynomials

Example:

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{array}{l} F_1^{(4)}(y) x_0 + F_2^{(4)}(y) x_1 = 0 \\ G_1^{(1)}(y) x_0 + G_2^{(1)}(y) x_1 = 0 \end{array} \right\} \subset \mathbb{P}^1[x] \times \mathbb{P}^4[y]$$

# CICYs

Idea: Ambient space  $\mathcal{A}$  is simply product of projective spaces, CY  $X$  given by intersection of zero locus of set of polynomials

Example:

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{array}{l} F_1^{(4)}(y) x_0 + F_2^{(4)}(y) x_1 = 0 \\ G_1^{(1)}(y) x_0 + G_2^{(1)}(y) x_1 = 0 \end{array} \right\} \subset \mathbb{P}^1[x] \times \mathbb{P}^4[y]$$

(CY  $\Leftrightarrow$  rows add to projective space dimension + 1)

# CICYs

Idea: Ambient space  $\mathcal{A}$  is simply product of projective spaces, CY  $X$  given by intersection of zero locus of set of polynomials

Example:

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{array}{l} F_1^{(4)}(y) x_0 + F_2^{(4)}(y) x_1 = 0 \\ G_1^{(1)}(y) x_0 + G_2^{(1)}(y) x_1 = 0 \end{array} \right\} \subset \mathbb{P}^1[x] \times \mathbb{P}^4[y]$$

(CY  $\Leftrightarrow$  rows add to projective space dimension + 1)

‘Favourable’ if  $h^{1,1}(X) = h^{1,1}(\mathcal{A})$

‘Kähler-favourable’ if  $\mathcal{K}(X) = \mathcal{K}(\mathcal{A})$  ( $\Rightarrow$  favourable)

For simplicity, below will only discuss Kähler-favourable CICYs

## Non-ambient flops (1/3)

---

# Non-ambient flops (1/3)

---

Consider example from previous slide

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \left( \begin{array}{cc} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

# Non-ambient flops (1/3)

---

Consider example from previous slide

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{pmatrix} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Single solution in  $\mathbb{P}^1$  only when  $F_1^{(4)}G_2^{(1)} = F_2^{(4)}G_1^{(1)}$

# Non-ambient flops (1/3)

---

Consider example from previous slide

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{pmatrix} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Single solution in  $\mathbb{P}^1$  only when  $F_1^{(4)}G_2^{(1)} = F_2^{(4)}G_1^{(1)}$

Entire  $\mathbb{P}^1$  precisely when  $F_1^{(4)} = G_2^{(1)} = F_2^{(4)} = G_1^{(1)} = 0$  (16 pts)

# Non-ambient flops (1/3)

Consider example from previous slide

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{pmatrix} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Single solution in  $\mathbb{P}^1$  only when  $F_1^{(4)} G_2^{(1)} = F_2^{(4)} G_1^{(1)}$

Entire  $\mathbb{P}^1$  precisely when  $F_1^{(4)} = G_2^{(1)} = F_2^{(4)} = G_1^{(1)} = 0$  (16 pts)

Singular manifold upon shrinking the  $\mathbb{P}^1$ s is

$$\{F_1^{(4)} G_2^{(1)} = F_2^{(4)} G_1^{(1)}\} \subset \mathbb{P}^4[y]$$



# Non-ambient flops (1/3)

Consider example from previous slide

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{pmatrix} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Single solution in  $\mathbb{P}^1$  only when  $F_1^{(4)} G_2^{(1)} = F_2^{(4)} G_1^{(1)}$

Entire  $\mathbb{P}^1$  precisely when  $F_1^{(4)} = G_2^{(1)} = F_2^{(4)} = G_1^{(1)} = 0$  (16 pts)

Singular manifold upon shrinking the  $\mathbb{P}^1$ s is

$$\{F_1^{(4)} G_2^{(1)} = F_2^{(4)} G_1^{(1)}\} \subset \mathbb{P}^4[y]$$

This is a tuned quintic, i.e. specific polynomial choice in  $[\mathbb{P}^4 | 5]$

# Non-ambient flops (1/3)

Consider example from previous slide

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{pmatrix} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Single solution in  $\mathbb{P}^1$  only when  $F_1^{(4)} G_2^{(1)} = F_2^{(4)} G_1^{(1)}$

Entire  $\mathbb{P}^1$  precisely when  $F_1^{(4)} = G_2^{(1)} = F_2^{(4)} = G_1^{(1)} = 0$  (16 pts)

Singular manifold upon shrinking the  $\mathbb{P}^1$ s is

$$\{F_1^{(4)} G_2^{(1)} = F_2^{(4)} G_1^{(1)}\} \subset \mathbb{P}^4[y]$$

This is a tuned quintic, i.e. specific polynomial choice in  $[\mathbb{P}^4 | 5]$

Known old story: CY reached from quintic via conifold transition

# Non-ambient flops (1/3)

Consider example from previous slide

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{pmatrix} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Single solution in  $\mathbb{P}^1$  only when  $F_1^{(4)} G_2^{(1)} = F_2^{(4)} G_1^{(1)}$

Entire  $\mathbb{P}^1$  precisely when  $F_1^{(4)} = G_2^{(1)} = F_2^{(4)} = G_1^{(1)} = 0$  (16 pts)

Singular manifold upon shrinking the  $\mathbb{P}^1$ s is

$$\{F_1^{(4)} G_2^{(1)} = F_2^{(4)} G_1^{(1)}\} \subset \mathbb{P}^4[y]$$

This is a tuned quintic, i.e. specific polynomial choice in  $[\mathbb{P}^4 | 5]$

Known old story: CY reached from quintic via conifold transition

(Allows to count shrinking curves: compare Euler characteristic)

## Non-ambient flops (2/3)

---

## Non-ambient flops (2/3)

---

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \left( \begin{array}{cc} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

## Non-ambient flops (2/3)

---

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \left( \begin{array}{cc} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Analogous to Atiyah flop: choose other pairing

## Non-ambient flops (2/3)

---

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{pmatrix} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Analogous to Atiyah flop: choose other pairing

$$\left\{ \begin{array}{l} F_1^{(4)}(y) x_0 + G_1^{(1)}(y) x_1 = 0 \\ F_2^{(4)}(y) x_0 + G_2^{(1)}(y) x_1 = 0 \end{array} \right\} \subset \mathcal{A}$$

## Non-ambient flops (2/3)

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \hline \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \begin{pmatrix} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{pmatrix} \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Analogous to Atiyah flop: choose other pairing

$$\left\{ \begin{array}{l} F_1^{(4)}(y) x_0 + G_1^{(1)}(y) x_1 = 0 \\ F_2^{(4)}(y) x_0 + G_2^{(1)}(y) x_1 = 0 \end{array} \right\} \subset \mathcal{A}$$

Now degrees of monomials don't seem to match



## Non-ambient flops (2/3)

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \hline \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \left( \begin{array}{cc} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Analogous to Atiyah flop: choose other pairing

$$\left\{ \begin{array}{l} F_1^{(4)}(y) x_0 + G_1^{(1)}(y) x_1 = 0 \\ F_2^{(4)}(y) x_0 + G_2^{(1)}(y) x_1 = 0 \end{array} \right\} \subset \mathcal{A}$$

Now degrees of monomials don't seem to match

$\Rightarrow$  Flopped space is complete intersection in toric variety  $\mathcal{A}$

## Non-ambient flops (2/3)

$$\left[ \begin{array}{c|cc} \mathbb{P}^1[x] & 1 & 1 \\ \hline \mathbb{P}^4[y] & 4 & 1 \end{array} \right] \equiv \left\{ \left( \begin{array}{cc} F_1^{(4)}(y) & F_2^{(4)}(y) \\ G_1^{(1)}(y) & G_2^{(1)}(y) \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \end{pmatrix} = \vec{0} \right\} \subset \mathbb{P}^1 \times \mathbb{P}^4$$

Analogous to Atiyah flop: choose other pairing

$$\left\{ \begin{array}{l} F_1^{(4)}(y) x_0 + G_1^{(1)}(y) x_1 = 0 \\ F_2^{(4)}(y) x_0 + G_2^{(1)}(y) x_1 = 0 \end{array} \right\} \subset \mathcal{A}$$

Now degrees of monomials don't seem to match

$\Rightarrow$  Flopped space is complete intersection in toric variety  $\mathcal{A}$

$x_0$	$x_1$	$y_0$	$y_1$	$y_2$	$y_3$	$y_4$	(Eq. 1)	(Eq. 2)
1	1	0	0	0	0	0	1	1
0	3	1	1	1	1	1	4	4

## Non-ambient flops (3/3)

---

## Non-ambient flops (3/3)

---

Above is special case of row of the form  $(\mathbb{P}^n \mid 1 \dots 1 \ 0 \dots 0)$

## Non-ambient flops (3/3)

---

Above is special case of row of the form  $(\mathbb{P}^n \mid 1 \dots 1 \, 0 \dots 0)$

Example:

# Non-ambient flops (3/3)

---

Above is special case of row of the form  $(\mathbb{P}^n \mid 1 \dots 1 0 \dots 0)$

Example:

$$\left[ \begin{array}{c|cccc} \mathbb{P}^2[x] & 1 & 1 & 1 & 0 & \dots \\ \vdots & \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vdots & \ddots \end{array} \right]$$

## Non-ambient flops (3/3)

---

Above is special case of row of the form  $(\mathbb{P}^n \mid 1 \dots 1 \ 0 \dots 0)$

Example:

$$\left[ \begin{array}{c|ccccc} \mathbb{P}^2[x] & 1 & 1 & 1 & 0 & \dots \\ \vdots & \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vdots & \ddots \end{array} \right] \equiv \left\{ \left( \begin{array}{ccc} F_1 & F_2 & F_3 \\ G_1 & G_2 & G_3 \\ H_1 & H_2 & H_3 \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \\ x_2 \end{pmatrix} = \vec{0} \right\}$$

## Non-ambient flops (3/3)

Above is special case of row of the form  $(\mathbb{P}^n \mid 1 \dots 1 \ 0 \dots 0)$

Example:

$$\left[ \begin{array}{c|cccc} \mathbb{P}^2[x] & 1 & 1 & 1 & 0 & \dots \\ \vdots & \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vdots & \ddots \end{array} \right] \equiv \left\{ \left( \begin{array}{ccc} F_1 & F_2 & F_3 \\ G_1 & G_2 & G_3 \\ H_1 & H_2 & H_3 \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \\ x_2 \end{pmatrix} = \vec{0} \right\}$$

Flopped space given by taking transpose of matrix of polynomials

$$X \equiv \{M\vec{x} = 0\} \subset \mathbb{P}^2 \times \dots \qquad X_{\text{flop}} \equiv \{M^T \vec{x} = 0\} \subset \mathcal{A}$$

New ambient space is toric with appropriate weight system



## Non-ambient flops (3/3)

Above is special case of row of the form  $(\mathbb{P}^n \mid 1 \dots 1 \ 0 \dots 0)$

Example:

$$\left[ \begin{array}{c|cccc} \mathbb{P}^2[x] & 1 & 1 & 1 & 0 & \dots \\ \vdots & \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vdots & \ddots \end{array} \right] \equiv \left\{ \left( \begin{array}{ccc} F_1 & F_2 & F_3 \\ G_1 & G_2 & G_3 \\ H_1 & H_2 & H_3 \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \\ x_2 \end{pmatrix} = \vec{0} \right\}$$

Flopped space given by taking transpose of matrix of polynomials

$$X \equiv \{M\vec{x} = 0\} \subset \mathbb{P}^2 \times \dots \qquad X_{\text{flop}} \equiv \{M^T \vec{x} = 0\} \subset \mathcal{A}$$

New ambient space is toric with appropriate weight system

New CY is isomorphic to original if  $\vec{v}_i$  are equal

## Non-ambient flops (3/3)

Above is special case of row of the form  $(\mathbb{P}^n | 1 \dots 1 0 \dots 0)$

Example:

$$\left[ \begin{array}{c|cccc} \mathbb{P}^2[x] & 1 & 1 & 1 & 0 & \dots \\ \vdots & \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vdots & \ddots \end{array} \right] \equiv \left\{ \left( \begin{array}{ccc} F_1 & F_2 & F_3 \\ G_1 & G_2 & G_3 \\ H_1 & H_2 & H_3 \end{array} \right) \begin{pmatrix} x_0 \\ x_1 \\ x_2 \end{pmatrix} = \vec{0} \right\}$$

Flopped space given by taking transpose of matrix of polynomials

$$X \equiv \{M\vec{x} = 0\} \subset \mathbb{P}^2 \times \dots \quad X_{\text{flop}} \equiv \{M^T \vec{x} = 0\} \subset \mathcal{A}$$

New ambient space is toric with appropriate weight system

New CY is isomorphic to original if  $\vec{v}_i$  are equal

Other rows giving flops:  $(\mathbb{P}^1 | 2 0 \dots 0)$ ,  $(\mathbb{P}^n | 2 1 \dots 1 0 \dots 0)$

## Application 1: Characterisation of the CICY list

---

## Application 1: Characterisation of the CICY list

---

There are 4,874 Kähler-favourable CICYs

## Application 1: Characterisation of the CICY list

---

There are 4,874 Kähler-favourable CICYs

There are 30,924 directions out of the Kähler cones

## Application 1: Characterisation of the CICY list

---

There are 4,874 Kähler-favourable CICYs

There are 30,924 directions out of the Kähler cones

Check for flops by checking for rows of the above forms

## Application 1: Characterisation of the CICY list

---

There are 4,874 Kähler-favourable CICYs

There are 30,924 directions out of the Kähler cones

Check for flops by checking for rows of the above forms

True number slightly smaller, some are 'ineffective splittings'

Check by comparing to Euler characteristic of singular space

## Application 1: Characterisation of the CICY list

---

There are 4,874 Kähler-favourable CICYs

There are 30,924 directions out of the Kähler cones

Check for flops by checking for rows of the above forms

True number slightly smaller, some are 'ineffective splittings'

Check by comparing to Euler characteristic of singular space

Characterisation:



## Application 1: Characterisation of the CICY list

---

There are 4,874 Kähler-favourable CICYs

There are 30,924 directions out of the Kähler cones

Check for flops by checking for rows of the above forms

True number slightly smaller, some are 'ineffective splittings'

Check by comparing to Euler characteristic of singular space

Characterisation:

More than 4,800 of the 4,874 CICYs admit flops

## Application 1: Characterisation of the CICY list

---

There are 4,874 Kähler-favourable CICYs

There are 30,924 directions out of the Kähler cones

Check for flops by checking for rows of the above forms

True number slightly smaller, some are 'ineffective splittings'

Check by comparing to Euler characteristic of singular space

### Characterisation:

More than 4,800 of the 4,874 CICYs admit flops

More than 29,000 of the 30,924 directions are flops

# Application 1: Characterisation of the CICY list

---

There are 4,874 Kähler-favourable CICYs

There are 30,924 directions out of the Kähler cones

Check for flops by checking for rows of the above forms

True number slightly smaller, some are 'ineffective splittings'

Check by comparing to Euler characteristic of singular space

## Characterisation:

More than 4,800 of the 4,874 CICYs admit flops

More than 29,000 of the 30,924 directions are flops

So non-ambient flops not only possible, but generic

## Application 2: Infinitely many Kähler cones (1/2)

---

## Application 2: Infinitely many Kähler cones (1/2)

---

In many cases, easy to see presence of infinitely many flops

## Application 2: Infinitely many Kähler cones (1/2)

---

In many cases, easy to see presence of infinitely many flops

Example:

$$\left[ \begin{array}{c|ccccc} \mathbb{P}^4 & 1 & 1 & 1 & 1 & 1 \\ \mathbb{P}^4 & 1 & 1 & 1 & 1 & 1 \end{array} \right] \quad (\text{CICY \#7761})$$

## Application 2: Infinitely many Kähler cones (1/2)

---

In many cases, easy to see presence of infinitely many flops

Example:

$$\left[ \begin{array}{c|ccccc} \mathbb{P}^4 & 1 & 1 & 1 & 1 & 1 \\ \mathbb{P}^4 & 1 & 1 & 1 & 1 & 1 \end{array} \right] \quad (\text{CICY \#7761})$$

See both Kähler cone boundaries are flops to isomorphic CY

## Application 2: Infinitely many Kähler cones (1/2)

In many cases, easy to see presence of infinitely many flops


Example:

$$\left[ \begin{array}{c|ccccc} \mathbb{P}^4 & 1 & 1 & 1 & 1 & 1 \\ \hline \mathbb{P}^4 & 1 & 1 & 1 & 1 & 1 \end{array} \right] \quad (\text{CICY \#7761})$$

See both Kähler cone boundaries are flops to isomorphic CY

But then those Kähler cones should admit further flops, etc

$\Rightarrow$  Infinitely many flops





## Application 2: Infinitely many Kähler cones (1/2)

In many cases, easy to see presence of infinitely many flops


Example:

$$\left[ \begin{array}{c|ccccc} \mathbb{P}^4 & 1 & 1 & 1 & 1 & 1 \\ \hline \mathbb{P}^4 & 1 & 1 & 1 & 1 & 1 \end{array} \right] \quad (\text{CICY \#7761})$$

See both Kähler cone boundaries are flops to isomorphic CY

But then those Kähler cones should admit further flops, etc

$\Rightarrow$  Infinitely many flops



Note: novel

Can't occur from ambient flops

(Num. triangulations always finite)

## Application 2: Infinitely many Kähler cones (2/2)

---

## Application 2: Infinitely many Kähler cones (2/2)

---

Occurs for higher  $h^{1,1}$ , e.g.

$$\left[ \begin{array}{c|cc} \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \end{array} \right]$$

CICY #7447

$$\left[ \begin{array}{c|c} \mathbb{P}^1 & 2 \\ \mathbb{P}^1 & 2 \\ \mathbb{P}^1 & 2 \\ \mathbb{P}^1 & 2 \end{array} \right]$$

CICY #7862

## Application 2: Infinitely many Kähler cones (2/2)

Occurs for higher  $h^{1,1}$ , e.g.

$$\left[ \begin{array}{c|cc} \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \end{array} \right]$$

CICY #7447

$$\left[ \begin{array}{c|c} \mathbb{P}^1 & 2 \\ \mathbb{P}^1 & 2 \\ \mathbb{P}^1 & 2 \\ \mathbb{P}^1 & 2 \end{array} \right]$$

CICY #7862

Admit flops to isomorphic  
CYs in every direction

⇒ Extended Kähler cones  
are fractals

## Application 2: Infinitely many Kähler cones (2/2)

Occurs for higher  $h^{1,1}$ , e.g.

$$\left[ \begin{array}{c|cc} \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \end{array} \right]$$


CICY #7447

$$\left[ \begin{array}{c|cc} \mathbb{P}^1 & 2 & \\ \mathbb{P}^1 & 2 & \\ \mathbb{P}^1 & 2 & \\ \mathbb{P}^1 & 2 & \end{array} \right]$$

CICY #7862

Admit flops to isomorphic  
CYs in every direction

⇒ Extended Kähler cones  
are fractals



Depiction of extended Kähler cones  
of CICYs #7447 and #7862

## Application 3: Cohomologies from configuration matrix

## Application 3: Cohomologies from configuration matrix

---

In many cases, flops allow one to write all zeroth cohomologies

## Application 3: Cohomologies from configuration matrix

---

In many cases, flops allow one to write all zeroth cohomologies

Simple example: flop between two isomorphic CYs



## Application 3: Cohomologies from configuration matrix

---

In many cases, flops allow one to write all zeroth cohomologies

Simple example: flop between two isomorphic CYs

Determining new Kähler cone  $K(X')$ :

## Application 3: Cohomologies from configuration matrix

---

In many cases, flops allow one to write all zeroth cohomologies

Simple example: flop between two isomorphic CYs

Determining new Kähler cone  $K(X')$ :

Know how  $c_2^i$ ,  $d_{ijk}$  change under flop

## Application 3: Cohomologies from configuration matrix

In many cases, flops allow one to write all zeroth cohomologies

Simple example: flop between two isomorphic CYs

Determining new Kähler cone  $K(X')$ :

Know how  $c_2^i, d_{ijk}$  change under flop

Isomorphic after  $\Rightarrow$  change of  $c_2^i, d_{ijk}$  undone by basis change

## Application 3: Cohomologies from configuration matrix

In many cases, flops allow one to write all zeroth cohomologies

Simple example: flop between two isomorphic CYs

Determining new Kähler cone  $K(X')$ :

Know how  $c_2^i, d_{ijk}$  change under flop

Isomorphic after  $\Rightarrow$  change of  $c_2^i, d_{ijk}$  undone by basis change

Getting cohomologies:

Zeroth cohomology in  $K(X')$  given by index using new  $c_2^i, d_{ijk}$

# Application 3: Cohomologies from configuration matrix

In many cases, flops allow one to write all zeroth cohomologies

Simple example: flop between two isomorphic CYs

Determining new Kähler cone  $K(X')$ :

Know how  $c_2^i, d_{ijk}$  change under flop

Isomorphic after  $\Rightarrow$  change of  $c_2^i, d_{ijk}$  undone by basis change

Getting cohomologies:

Zeroth cohomology in  $K(X')$  given by index using new  $c_2^i, d_{ijk}$

$$\left[ \begin{array}{c|c} \mathbb{P}^1 & 2 \\ \mathbb{P}^3 & 4 \end{array} \right]$$

# Application 3: Cohomologies from configuration matrix

In many cases, flops allow one to write all zeroth cohomologies

Simple example: flop between two isomorphic CYs


Determining new Kähler cone  $K(X')$ :

Know how  $c_2^i$ ,  $d_{ijk}$  change under flop

Isomorphic after  $\Rightarrow$  change of  $c_2^i$ ,  $d_{ijk}$  undone by basis change

Getting cohomologies:

Zeroth cohomology in  $K(X')$  given by index using new  $c_2^i$ ,  $d_{ijk}$



## Application 3: Cohomologies from configuration matrix

In many cases, flops allow one to write all zeroth cohomologies


## Determining new Kähler cone $K(X')$ :

## Know how $c_2^i$ , $d_{ijk}$ change under flop

Isomorphic after  $\Rightarrow$  change of  $c^i, d_{ijk}$  undone by basis change

## Getting cohomologies:

Zeroth cohomology in  $K(X')$  given by index using new  $c_2^i, d_{ijk}$



## Application 4: Non-ambient flops for toric CIs, gCICYs



## Application 4: Non-ambient flops for toric CIs, gCICYs

---

Extension of non-ambient flops to toric case is straightforward

## Application 4: Non-ambient flops for toric CIs, gCICYs

---

Extension of non-ambient flops to toric case is straightforward

Above were special cases: one side consistent with  $\mathbb{P}^n \times \dots$

## Application 4: Non-ambient flops for toric CIs, gCICYs

Extension of non-ambient flops to toric case is straightforward

Above were special cases: one side consistent with  $\mathbb{P}^n \times \dots$

$$\text{e.g. } \left\{ \begin{array}{l} F_1^{(2)} x_0 + F_2^{(3)} x_1 = 0 \\ G_1^{(4)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} F_1^{(2)} x_0 + G_1^{(4)} x_1 = 0 \\ F_2^{(3)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\}$$

## Application 4: Non-ambient flops for toric CIs, gCICYs

Extension of non-ambient flops to toric case is straightforward

Above were special cases: one side consistent with  $\mathbb{P}^n \times \dots$

$$\text{e.g. } \left\{ \begin{array}{l} F_1^{(2)} x_0 + F_2^{(3)} x_1 = 0 \\ G_1^{(4)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} F_1^{(2)} x_0 + G_1^{(4)} x_1 = 0 \\ F_2^{(3)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\}$$

gCICYs: allow negative entries [\[1507.03235\]](#)

## Application 4: Non-ambient flops for toric CIs, gCICYs

Extension of non-ambient flops to toric case is straightforward

Above were special cases: one side consistent with  $\mathbb{P}^n \times \dots$

$$\text{e.g. } \left\{ \begin{array}{l} F_1^{(2)} x_0 + F_2^{(3)} x_1 = 0 \\ G_1^{(4)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} F_1^{(2)} x_0 + G_1^{(4)} x_1 = 0 \\ F_2^{(3)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\}$$

gCICYs: allow negative entries [\[1507.03235\]](#)

For example:

gCICY on the right recently used  
for model-building [\[2010.09763\]](#)

$$\left[ \begin{array}{c|cc} \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 0 & 2 \\ \mathbb{P}^1 & 3 & -1 \end{array} \right]$$

## Application 4: Non-ambient flops for toric CIs, gCICYs

Extension of non-ambient flops to toric case is straightforward

Above were special cases: one side consistent with  $\mathbb{P}^n \times \dots$

$$\text{e.g. } \left\{ \begin{array}{l} F_1^{(2)} x_0 + F_2^{(3)} x_1 = 0 \\ G_1^{(4)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} F_1^{(2)} x_0 + G_1^{(4)} x_1 = 0 \\ F_2^{(3)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\}$$

gCICYs: allow negative entries [\[1507.03235\]](#)

For example:

gCICY on the right recently used  
for model-building [\[2010.09763\]](#)

$$\left[ \begin{array}{c|cc} \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 0 & 2 \\ \mathbb{P}^1 & 3 & -1 \end{array} \right]$$

Flops developed above can be performed here too, in rows 1-4

## Application 4: Non-ambient flops for toric CIs, gCICYs

Extension of non-ambient flops to toric case is straightforward

Above were special cases: one side consistent with  $\mathbb{P}^n \times \dots$

$$\text{e.g. } \left\{ \begin{array}{l} F_1^{(2)} x_0 + F_2^{(3)} x_1 = 0 \\ G_1^{(4)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\} \leftrightarrow \left\{ \begin{array}{l} F_1^{(2)} x_0 + G_1^{(4)} x_1 = 0 \\ F_2^{(3)} x_0 + G_2^{(5)} x_1 = 0 \end{array} \right\}$$

gCICYs: allow negative entries [\[1507.03235\]](#)

For example:

gCICY on the right recently used  
for model-building [\[2010.09763\]](#)

$$\left[ \begin{array}{c|cc} \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 1 & 1 \\ \mathbb{P}^1 & 0 & 2 \\ \mathbb{P}^1 & 3 & -1 \end{array} \right]$$

Flops developed above can be performed here too, in rows 1-4

Result is generalised toric complete intersections

# Extensions

---



# Extensions

---

- Classify non-ambient flops for all CICYs, toric CIs, gCICYs, to better our understanding of web of CY3s

# Extensions

---

- Classify non-ambient flops for all CICYs, toric CIs, gCICYs, to better our understanding of web of CY3s
- Characterise possible structures for extended Kähler moduli space (finite chamber cases, fractal cases, etc)

# Extensions

---

- Classify non-ambient flops for all CICYs, toric CIs, gCICYs, to better our understanding of web of CY3s
- Characterise possible structures for extended Kähler moduli space (finite chamber cases, fractal cases, etc)
- Connection to GLSMs (infinitely many phases, etc)

# Extensions

---

- Classify non-ambient flops for all CICYs, toric CIs, gCICYs, to better our understanding of web of CY3s
- Characterise possible structures for extended Kähler moduli space (finite chamber cases, fractal cases, etc)
- Connection to GLSMs (infinitely many phases, etc)
- Cohomology: relation of flops to higher cohomologies - read them off too?

Thanks for listening

Questions?