

# Vanishing Yukawa couplings in heterotic compactifications

Matthew Magill

Based on work with L. Anderson, J. Gray, M. Larfors and R. Schneider  
2103.10454

Uppsala University

27th April, 2021

Seminar Series on String Phenomenology



# Setup

- Compactify  $E_8 \times E_8$  heterotic string on Calabi-Yau three-fold,  $X$ , with a bundle,  $V \rightarrow X$ ;
  - Structure group of  $V$  embedded in  $E_8$  and its commutant is gauge group of effective theory;
  - Will assume  $V$  an  $SU(3)$  bundle  $\rightsquigarrow E_6$  gauge symmetry;
- 
- Effective  $N = 1$  supersymmetric gauge theory.
  - Interactions of chiral supermultiplets,  $\Phi$ , governed by **superpotential**  $\int d^2\theta W(\Phi) + h.c$ ;

# Yukawa coupling I

- Perturbative Yukawa couplings correspond to cubic term in the (perturbative) superpotential;
- **Physical** Yukawa couplings depend on correct normalisation of fields  $\implies$  explicit Calabi-Yau metric dependence;
- Whether coupling vanishes or not is normalisation independent;
- Implications for physics, e.g. top quark mass .

# Yukawa coupling II

## The holomorphic Yukawa coupling

- Matter fields correspond to bundle-valued harmonic one-forms  $\Phi^I \leftrightarrow \nu_I \in \text{Harm}^{0,1}(V)$ ;
- From 10d action, deduce cubic term  $W \supset \lambda(\nu_I, \nu_J, \nu_K) \Phi^I \Phi^J \Phi^K$ , with

$$\lambda(\nu_1, \nu_2, \nu_3) = \int_X \Omega \wedge \text{tr}(\nu_1 \wedge \nu_2 \wedge \nu_3);$$

- $\Omega \in \Omega^{3,0}$ , holomorphic three-form;
- $\text{tr} = \text{map induced from projection } V^{\otimes 3} \rightarrow \Lambda^3 V \rightarrow \mathcal{O}_X$ ;

# Yukawa coupling III

$$\lambda(\nu_1, \nu_2, \nu_3) = \int_X \Omega \wedge \text{tr}(\nu_1 \wedge \nu_2 \wedge \nu_3);$$

## Key properties

- Depends only on **cohomology** classes  $[\nu_i]$ ;
- Composition of map  $H^1(X, V)^{\otimes 3} \rightarrow H^3(X, \mathcal{O}_X)$ , and  $\int_X \Omega \wedge - : H^3(X, \mathcal{O}_X) \rightarrow \mathbb{C}$ , **isomorphism!**

## Implications

- $\lambda(\nu_1, \nu_2, \nu_3) = 0$  if and only if the **cohomology class**  $[\text{tr}(\nu_1 \wedge \nu_2 \wedge \nu_3)] = 0$ .
- Computing  $[\text{tr}(\nu_1 \wedge \nu_2 \wedge \nu_3)]$  is **very hard!**

# Outline

- 1 Background and Motivation
- 2 Sheaf Cohomology**
- 3 Vanishing theorems

# Motivation for sheaf cohomology


## Decomposing cohomology

Need a mechanism to **split up** cohomology group that is **compatible** with Yukawa computation.

# Motivation for sheaf cohomology

## Decomposing cohomology

Need a mechanism to **split up** cohomology group that is **compatible** with Yukawa computation.






# Motivation for sheaf cohomology

## Decomposing cohomology


Need a mechanism to **split up** cohomology group that is **compatible** with Yukawa computation.



# Motivation for sheaf cohomology

## Decomposing cohomology

Need a mechanism to **split up** cohomology group that is **compatible** with Yukawa computation.



We “decompose” **before** taking cohomology  $\rightsquigarrow$  gives controlled splitting.

## Examples of sheaves

- For  $X$  a complex manifold, define  $\mathcal{O}_X(U) = \{\text{holomorphic maps } U \rightarrow \mathbb{C}\}$  with the natural restriction maps: **structure sheaf**.
- For any vector bundle,  $V$ , define  $\mathcal{V}(U) = \Gamma(U, V|_U)$ : **sheaf of sections**.
- For any abelian group,  $A$ , can define  $\underline{A}(U) = \{\text{locally constant functions } U \rightarrow A\}$ : **constant sheaf**.



# Basic definitions

## Exact sequences of sheaves

A complex of sheaves

$$\cdots \mathcal{F}_2 \xrightarrow{f_2} \mathcal{F}_1 \xrightarrow{f_1} \mathcal{F}_0 \rightarrow \cdots$$

is called exact if the sequence of groups obtained by restricting to smaller and smaller sets, is exact. **Roughly:**

- $\ker(f_i(U)) = \operatorname{Im}(f_{i+1}(U))$  for all “very small”  $U$

## Example

$$0 \rightarrow \mathbb{Z} \xrightarrow{2\pi i \cdot} \mathcal{O}_{\mathbb{C}^*}(-, \mathbb{C}) \xrightarrow{\exp} \mathcal{O}_{\mathbb{C}^*}(-, \mathbb{C}^*) \rightarrow 0$$

# Global sections

## Exactness & global sections

- Exactness not preserved on **global** sections;

# Global sections

## Exactness & global sections

- Exactness not preserved on **global** sections;
- Pass to cohomology  $\rightsquigarrow$  long exact sequence of groups:

$$\cdots \rightarrow H^{i-1}(\mathcal{O}_{\mathbb{C}^*}^*) \xrightarrow{\delta} H^i(\mathbb{Z}) \rightarrow H^i(\mathcal{O}_{\mathbb{C}^*}) \rightarrow H^i(\mathcal{O}_{\mathbb{C}^*}^*) \rightarrow \cdots$$

# Global sections

## Exactness & global sections

- Exactness not preserved on **global** sections;
- Pass to cohomology  $\rightsquigarrow$  long exact sequence of groups:

$$\cdots \rightarrow H^{i-1}(\mathcal{O}_{\mathbb{C}^*}^*) \xrightarrow{\delta} H^i(\mathbb{Z}) \rightarrow H^i(\mathcal{O}_{\mathbb{C}^*}) \rightarrow H^i(\mathcal{O}_{\mathbb{C}^*}^*) \rightarrow \cdots$$

- Defined using special exact sequence of sheaves:

$$0 \rightarrow \mathcal{F} \rightarrow \mathcal{I}^0 \rightarrow \mathcal{I}^1 \rightarrow \cdots$$

$$H^i(X, \mathcal{F}) = \frac{\ker(\mathcal{I}^i(X) \rightarrow \mathcal{I}^{i+1}(X))}{\operatorname{Im}(\mathcal{I}^{i-1}(X) \rightarrow \mathcal{I}^i(X))}.$$

# Godement resolution

- We will use “Godement resolutions”
- Behaves well with tensor product (“universal exactness”):

$$\begin{array}{ccc} \mathcal{G}^\bullet(\mathcal{F}) \otimes \mathcal{G}^\bullet(\mathcal{F}') & \xrightarrow{u} & \mathcal{G}^\bullet(\mathcal{F} \otimes \mathcal{F}') \\ \uparrow & & \uparrow \\ \mathcal{F} \otimes \mathcal{F}' & \xrightarrow{\text{Id}} & \mathcal{F} \otimes \mathcal{F}' \end{array}$$

- (Also for  $\mathcal{F}, \mathcal{F}'$  complexes of sheaves)



# Cup product

## Definition

- Use Godement resolution to define cohomology  

$$H^i(X, V) = \frac{\ker(\mathcal{G}^i(V)(X) \rightarrow \mathcal{G}^{i+1}(V)(X))}{\operatorname{Im}(\mathcal{G}^{i-1}(V)(X) \rightarrow \mathcal{G}^i(V)(X))};$$
- **Cup product** is a map  $\smile: H^i(\mathcal{F}) \otimes H^j(\mathcal{F}') \rightarrow H^{i+j}(\mathcal{F} \otimes \mathcal{F}')$ ;
- Defined by  $[\nu] \otimes [\mu] \mapsto [u(\nu \otimes \mu)]$ ;
- Same operation as wedge product.

$$\begin{array}{ccc}
 \mathcal{G}^\bullet(\mathcal{F}) \otimes \mathcal{G}^\bullet(\mathcal{F}') & \xrightarrow{u} & \mathcal{G}^\bullet(\mathcal{F} \otimes \mathcal{F}') \\
 \uparrow & & \uparrow \\
 \mathcal{F} \otimes \mathcal{F}' & \xrightarrow{\operatorname{Id}} & \mathcal{F} \otimes \mathcal{F}'
 \end{array}$$

# Cup product

## Definition

- Use Godement resolution to define cohomology

$$H^i(X, V) = \frac{\ker(\mathcal{G}^i(V)(X) \rightarrow \mathcal{G}^{i+1}(V)(X))}{\operatorname{Im}(\mathcal{G}^{i-1}(V)(X) \rightarrow \mathcal{G}^i(V)(X))};$$

- **Cup product** is a map  $\smile: H^i(\mathcal{F}) \otimes H^j(\mathcal{F}') \rightarrow H^{i+j}(\mathcal{F} \otimes \mathcal{F}')$ ;

## Redefinition of Yukawa coupling

Class appearing in Yukawa coupling,

$$[\operatorname{tr}(\nu_1 \wedge \nu_2 \wedge \nu_3)]$$

is expressed as **composition** of

- Two cup products  $\smile \otimes \smile: H^1(X, V)^{\otimes 3} \rightarrow H^3(X, V^{\otimes 3})$ ;
- Map  $\operatorname{tr}: H^3(X, V^{\otimes 3}) \rightarrow H^3(X, \mathcal{O}_X)$

# Outline

- 1 Background and Motivation
- 2 Sheaf Cohomology
- 3 Vanishing theorems

# Known vanishing theorems I

## $(p, q)$ -vanishing theorem

Braun, He, Ovrut 0601204, ...

- Can be applied when  $X$  is elliptically fibred,  $E \rightarrow X \xrightarrow{p} B$ ;
- Use Leray spectral sequence to **decompose**  
 $H^1(X, V) \hookrightarrow H^0(B, R^1 p_* V) \oplus H^1(B, p_* V)$ ;
- Only allowed couplings are of form

$$H^1(B, p_* V) \otimes H^1(B, p_* V) \otimes H^0(B, R^1 p_* V)$$

- **Assuming**  $p_* V, R^1 p_* V$  are locally free.

## Stability walls

Anderson, Gray, Ovrut 1001.2317, ...

# Known vanishing theorems I

## $(p, q)$ -vanishing theorem

Braun, He, Ovrut 0601204, ...

## Stability walls

Anderson, Gray, Ovrut 1001.2317, ...

- $SU(3)$  bundle splits  $V \rightarrow K \oplus F$  at stability wall;
- **Enhanced** effective gauge group  $E_6 \rightarrow E_6 \times U(1)$ ;
- **Splits** cohomology  $H^1(V) = H^1(K) \oplus H^1(F)$ ;
- Summands have definite  $U(1)$  charge.
- Only couplings that have **zero charge** are allowed at this locus.

# Known vanishing theorems II

## Koszul complexes

Blesneag, Buchbinder, Candelas, Lukas 1512.05322, ...

- Applicable when  $X \hookrightarrow \mathcal{A} = \mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_k}$  and  $V = \mathcal{V}|_X$  for  $\mathcal{V}$  a sum of line bundles on ambient space;
- Exists **canonical resolution**, “Koszul complex”:

$$0 \rightarrow \Lambda^\bullet \mathcal{N}^* \otimes \mathcal{V} \rightarrow V \rightarrow 0$$

- Induces **decomposition**

$$H^1(X, V) \hookrightarrow \bigoplus_{\tau=1}^{\dim \mathcal{A}-2} H^\tau(\mathcal{A}, \Lambda^{\tau-1} \mathcal{N}^* \otimes \mathcal{V})$$

- Call  $\tau$  the “type” of class;
- Coupling  $\lambda(\nu_1, \nu_2, \nu_3)$  vanishes unless  $\tau_1 + \tau_2 + \tau_3 \geq \dim \mathcal{A}$ .

# Towards new vanishing theorems

## Setup

Anderson, Gray, Larfors, M.M., Schneider, 2103.10454

- Start with a resolution

$$0 \rightarrow \mathcal{F}_1 \xrightarrow{f} \mathcal{F}_0 \xrightarrow{\pi} V \rightarrow 0$$

- Use Godement resolution:

$$0 \rightarrow \mathcal{G}^\bullet(\mathcal{F}_1) \rightarrow \mathcal{G}^\bullet(\mathcal{F}_0) \rightarrow \mathcal{G}^\bullet(V) \rightarrow 0$$

# Towards new vanishing theorems

$$0 \rightarrow \mathcal{G}^\bullet(\mathcal{F}_1) \rightarrow \mathcal{G}^\bullet(\mathcal{F}_0) \rightarrow \mathcal{G}^\bullet(V) \rightarrow 0$$

## Coboundary map

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^2(V)) \longrightarrow 0 \\
 & & \uparrow & & \uparrow & & \uparrow \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^1(V)) \longrightarrow 0 \\
 & & & & & & \uparrow \\
 & & & & & & \nu
 \end{array}$$



# Towards new vanishing theorems

$$0 \rightarrow \mathcal{G}^\bullet(\mathcal{F}_1) \rightarrow \mathcal{G}^\bullet(\mathcal{F}_0) \rightarrow \mathcal{G}^\bullet(V) \rightarrow 0$$

## Coboundary map

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^2(V)) \longrightarrow 0 \\
 & & \uparrow & & \uparrow & & \uparrow \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^1(V)) \longrightarrow 0
 \end{array}$$

$$\hat{\nu} \xrightarrow{\quad\quad\quad} \nu$$

# Towards new vanishing theorems

$$0 \rightarrow \mathcal{G}^\bullet(\mathcal{F}_1) \rightarrow \mathcal{G}^\bullet(\mathcal{F}_0) \rightarrow \mathcal{G}^\bullet(V) \rightarrow 0$$

## Coboundary map

$$\begin{array}{ccccccc}
 & & \mathfrak{d}\hat{\nu} & & & & \\
 & & \downarrow & & & & \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^2(V)) \longrightarrow 0 \\
 & & \uparrow & & \uparrow & & \uparrow \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^1(V)) \longrightarrow 0 \\
 & & & & \uparrow & & \\
 & & & & \hat{\nu} & \xrightarrow{\quad \nu \quad} & \nu
 \end{array}$$

# Towards new vanishing theorems

$$0 \rightarrow \mathcal{G}^\bullet(\mathcal{F}_1) \rightarrow \mathcal{G}^\bullet(\mathcal{F}_0) \rightarrow \mathcal{G}^\bullet(V) \rightarrow 0$$

## Coboundary map

$$\begin{array}{ccccccc}
 & & \mathfrak{d}\hat{\nu} & \xrightarrow{\quad} & 0 & & \\
 & & \uparrow & & \uparrow & & \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^2(V)) \longrightarrow 0 \\
 & & \uparrow & & \uparrow & & \uparrow \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^1(V)) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & \hat{\nu} & \xrightarrow{\quad} & \nu & & 
 \end{array}$$

Curved green arrows indicate coboundary maps:  $\mathfrak{d}\hat{\nu} \rightarrow 0$ ,  $0 \rightarrow \nu$ ,  $\hat{\nu} \rightarrow \mathfrak{d}\hat{\nu}$ , and  $\nu \rightarrow \nu$ .

# Towards new vanishing theorems

$$0 \rightarrow \mathcal{G}^\bullet(\mathcal{F}_1) \rightarrow \mathcal{G}^\bullet(\mathcal{F}_0) \rightarrow \mathcal{G}^\bullet(V) \rightarrow 0$$

## Coboundary map

$$\begin{array}{ccccccc}
 \delta\nu & \xrightarrow{\quad\quad\quad} & \mathfrak{d}\hat{\nu} & & & & \\
 & & \nwarrow & & & & \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^2(V)) \longrightarrow 0 \\
 & & \uparrow & & \uparrow & & \uparrow \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^1(V)) \longrightarrow 0 \\
 & & & & \nearrow & & \\
 & & & & \hat{\nu} & \xrightarrow{\quad\quad\quad} & \nu
 \end{array}$$

# Cohomology decomposition

## Cohomological type

We will say a cohomology class,  $[\nu]$  is

- **Type 1** if  $\partial\hat{\nu} = 0$ ;
- **Type 2** else.
- Induces  $H^1(V) \hookrightarrow H^1(\mathcal{F}_0) \oplus H^2(\mathcal{F}_1)$

$$\begin{array}{ccccccc}
 \delta\nu & \xrightarrow{\quad\quad\quad} & \partial\hat{\nu} & & & & \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^2(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^2(V)) \longrightarrow 0 \\
 & & \uparrow & & \uparrow & & \uparrow \\
 0 & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_1)) & \longrightarrow & \Gamma(\mathcal{G}^1(\mathcal{F}_0)) & \longrightarrow & \Gamma(\mathcal{G}^1(V)) \longrightarrow 0 \\
 & & & & \hat{\nu} & \xrightarrow{\quad\quad\quad} & \nu
 \end{array}$$

## Yukawa contribution

- Represent classes in  $H^1(V)$  by sections

$$(\hat{\nu}, \delta\nu) \in \Gamma(\mathcal{G}^1(\mathcal{F}_0)) \oplus \Gamma(\mathcal{G}^2(\mathcal{F}_1));$$

- **Type 1** represented by  $\hat{\nu} \in \mathcal{G}^1(\mathcal{F}_0)$ ;
- (Type 1)<sup>3</sup> coupling represented by

$$[\hat{\nu}_1 \otimes \hat{\nu}_2 \otimes \hat{\nu}_3] \in \mathcal{G}^3(\Lambda^3 \mathcal{F}_0);$$

- **If**  $H^3(\Lambda^3 \mathcal{F}_0) = 0$ , then the coupling must be trivial.

## Yukawa contribution

- Represent classes in  $H^1(V)$  by sections

$$(\hat{\nu}, \delta\nu) \in \Gamma(\mathcal{G}^1(\mathcal{F}_0)) \oplus \Gamma(\mathcal{G}^2(\mathcal{F}_1));$$

- Similarly, (type 1)<sup>2</sup>(type 2) coupling is represented by

$$\begin{bmatrix} \hat{\nu}_1 \otimes \hat{\nu}_2 \otimes \hat{\nu}_3 \\ \hat{\nu}_1 \otimes \hat{\nu}_2 \otimes \delta\nu_3 \end{bmatrix} \in \Gamma(\mathcal{G}^3(\Lambda^3 \mathcal{F}_0)) \oplus \Gamma(\mathcal{G}^4(\Lambda^2 \mathcal{F}_0 \otimes \mathcal{F}_1)).$$

- If  $H^4(\Lambda^2 \mathcal{F}_0 \otimes \mathcal{F}_1) = 0$  and  $H^3(\Lambda^3 \mathcal{F}_0) = 0$ , then (type 1)<sup>2</sup>(type 2) couplings vanish.

# Vanishing theorem

## Theorem

2103.10454

- If  $H^3(\Lambda^3 \mathcal{F}_0) = 0$ , then the coupling between three type 1 fields vanishes;
- If **in addition**  $H^4(\mathcal{F}_1 \otimes \Lambda^2 \mathcal{F}_0) = 0$ , then the coupling between two type 1s and a type 2 vanishes.



# Vanishing theorem

## Theorem

2103.10454

- If  $H^3(\Lambda^3 \mathcal{F}_0) = 0$ , then the coupling between three type 1 fields vanishes;
- If **in addition**  $H^4(\mathcal{F}_1 \otimes \Lambda^2 \mathcal{F}_0) = 0$ , then the coupling between two type 1s and a type 2 vanishes.

## Higher type couplings?

- Rely on exact sequence:

$$?? \rightarrow \Lambda^2 \mathcal{F}_0 \otimes \mathcal{F}_1 \rightarrow \Lambda^3 \mathcal{F}_0 \rightarrow \Lambda^3 V \rightarrow 0;$$

- Higher type needs the unknown terms.

# Vanishing theorem

## Theorem

2103.10454

- If  $H^3(\Lambda^3 \mathcal{F}_0) = 0$ , then the coupling between three type 1 fields vanishes;
  - If **in addition**  $H^4(\mathcal{F}_1 \otimes \Lambda^2 \mathcal{F}_0) = 0$ , then the coupling between two type 1s and a type 2 vanishes.
- 
- Argument can be extended to complex of **arbitrary** length, **still** only these couplings are trustworthy;
  - Can sometimes weaken second condition.

# Summary and Outlook

## Summary & Conclusions

- New theorems constraining Yukawa couplings;
- Wide range of applicability;
- Heterotic yukawa couplings are generically textured?

# Summary and Outlook

## Summary & Conclusions

- New theorems constraining Yukawa couplings;
- Wide range of applicability;
- Heterotic yukawa couplings are generically textured?

## Outlook

- Expect these results can be strengthened;
- $SU(4)$ ,  $SU(5)$ -structure group, Wilson lines?
- Physical explanation?

# Summary and Outlook

## Summary & Conclusions

- New theorems constraining Yukawa couplings;
- Wide range of applicability;
- Heterotic yukawa couplings are generically textured?

## Outlook

- Expect these results can be strengthened;
- $SU(4)$ ,  $SU(5)$ -structure group, Wilson lines?
- Physical explanation?

Thanks for listening!