

Working with Automated Reasoning Tools – Higher-Order Resolution –

Christoph Benz Müller and Geoff Sutcliffe

Department of Computer Science, Saarland University

TPTPSYS'08

SS08, Block Course at Saarland University, Germany



Approaches to Higher-Order Resolution: Andrews' \mathcal{R}

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

We present and discuss Andrews' higher-order resolution calculus [Andrews71]; we call this calculus \mathcal{R}

λ -Conversion

- Andrews' provides two rules for α -conversion and β -reduction

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

We present and discuss Andrews' higher-order resolution calculus [Andrews71]; we call this calculus \mathcal{R}

λ -Conversion

- Andrews' provides two rules for α -conversion and β -reduction
- he does not provide a rule for η -conversion: consequently η -equality of two terms (e.g., $f_{\iota \rightarrow \iota} \doteq \lambda X_{\iota}. f X$) cannot be proven in this approach without employing the functional extensionality axiom of appropriate type

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

We present and discuss Andrews' higher-order resolution calculus [Andrews71]; we call this calculus \mathcal{R}

λ -Conversion

- Andrews' provides two rules for α -conversion and β -reduction
- he does not provide a rule for η -conversion: consequently η -equality of two terms (e.g., $f_{\iota \rightarrow \iota} \doteq \lambda X_{\iota}.f X$) cannot be proven in this approach without employing the functional extensionality axiom of appropriate type
- we omit explicit rules for α - and β -convertibility and instead treat them implicitly, i.e. we assume that the presented rules operate on input and generate output in β -normal form and we automatically identify terms which differ only with respect to the names of bound variables

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Clause Normalisation

- \mathcal{R} introduces only four rules belonging to clause normalisation: negation elimination, conjunction elimination, existential elimination, and universal elimination

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Clause Normalisation

- \mathcal{R} introduces only four rules belonging to clause normalisation: negation elimination, conjunction elimination, existential elimination, and universal elimination
- as our presentation of clauses in contrast to [Andrews71] explicitly mentions the polarities of clauses and brackets the literal atoms we need additional structural rules, e.g., the rule \vee^T

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Clause Normalisation

- \mathcal{R} introduces only four rules belonging to clause normalisation: negation elimination, conjunction elimination, existential elimination, and universal elimination
- as our presentation of clauses in contrast to [Andrews71] explicitly mentions the polarities of clauses and brackets the literal atoms we need additional structural rules, e.g., the rule \vee^T

- negation elimination:
$$\frac{C \vee [\neg A]^T}{C \vee [A]^F} \neg^T \quad \frac{C \vee [\neg A]^F}{C \vee [A]^T} \neg^F$$

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Clause Normalisation

- \mathcal{R} introduces only four rules belonging to clause normalisation: negation elimination, conjunction elimination, existential elimination, and universal elimination
- as our presentation of clauses in contrast to [Andrews71] explicitly mentions the polarities of clauses and brackets the literal atoms we need additional structural rules, e.g., the rule \vee^T

■ negation elimination:

$$\frac{C \vee [\neg A]^T}{C \vee [A]^F} \neg^T \quad \frac{C \vee [\neg A]^F}{C \vee [A]^T} \neg^F$$

- conjunction/disjunction elimination:

$$\frac{C \vee [A \vee B]^T}{C \vee [A]^T \vee [B]^T} \vee^T \quad \frac{C \vee [A \vee B]^F}{C \vee [A]^F} \vee_l^F \quad \frac{C \vee [A \vee B]^F}{C \vee [B]^F} \vee_r^F$$

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Clause Normalisation (contd.)

- existential/universal elimination:

$$\frac{C \vee [\Pi^\alpha \mathbf{A}]^T}{C \vee [\mathbf{A} \mathbf{X}_\alpha]^T} \Pi^T \qquad \frac{C \vee [\Pi^\alpha \mathbf{A}]^F}{C \vee [\mathbf{A} \mathbf{s}_\alpha]^F} \Pi^F$$

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Clause Normalisation (contd.)

- existential/universal elimination:

$$\frac{C \vee [\Pi^\alpha \mathbf{A}]^T}{C \vee [\mathbf{A} X_\alpha]^T} \Pi^T \quad \frac{C \vee [\Pi^\alpha \mathbf{A}]^F}{C \vee [\mathbf{A} s_\alpha]^F} \Pi^F$$

X_α is a new free variable and s_α is a new Skolem term

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Clause Normalisation (contd.)

- existential/universal elimination:

$$\frac{C \vee [\Pi^\alpha \mathbf{A}]^T}{C \vee [\mathbf{A} \mathbf{X}_\alpha]^T} \Pi^T \quad \frac{C \vee [\Pi^\alpha \mathbf{A}]^F}{C \vee [\mathbf{A} \mathbf{s}_\alpha]^F} \Pi^F$$

\mathbf{X}_α is a new free variable and \mathbf{s}_α is a new Skolem term

- additionally Andrews presents rules addressing commutativity and associativity of the \vee -operator connecting the clauses literals; we treat this implicit here

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Clause Normalisation (contd.)

- existential/universal elimination:

$$\frac{C \vee [\Pi^\alpha A]^\top}{C \vee [A X_\alpha]^\top} \Pi^\top \quad \frac{C \vee [\Pi^\alpha A]^F}{C \vee [A s_\alpha]^F} \Pi^F$$

X_α is a new free variable and s_α is a new Skolem term

- additionally Andrews presents rules addressing commutativity and associativity of the \vee -operator connecting the clauses literals; we treat this implicit here
- we refer with $\text{Cnf}(A)$ to the set of clauses obtained from formula A by exhaustive clause normalisation

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Resolution & Factorisation

- Instead of a resolution and a factorisation rule — which work in connection with unification — Andrews presents a simplification and a cut rule. The cut rule is only applicable to clauses with two complementary literals which have identical atoms. Similarly Sim is defined only for clauses with two identical literals. In order to generate identical literal atoms during the refutation process these two rules have to be combined with the substitution rule Sub presented below.

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Resolution & Factorisation

- Instead of a resolution and a factorisation rule — which work in connection with unification — Andrews presents a simplification and a cut rule. The cut rule is only applicable to clauses with two complementary literals which have identical atoms. Similarly Sim is defined only for clauses with two identical literals. In order to generate identical literal atoms during the refutation process these two rules have to be combined with the substitution rule Sub presented below.

- Simplification:
$$\frac{[A]^\mu \vee [A]^\mu \vee C}{[A]^\mu \vee C} \text{ Sim}$$

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Resolution & Factorisation

- Instead of a resolution and a factorisation rule — which work in connection with unification — Andrews presents a simplification and a cut rule. The cut rule is only applicable to clauses with two complementary literals which have identical atoms. Similarly Sim is defined only for clauses with two identical literals. In order to generate identical literal atoms during the refutation process these two rules have to be combined with the substitution rule Sub presented below.

- Simplification:

$$\frac{[A]^\mu \vee [A]^\mu \vee C}{[A]^\mu \vee C} \text{ Sim}$$

- Cut:

$$\frac{[A]^\mu \vee C \quad [A]^\nu \vee D}{C \vee D} \text{ Cut}$$

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Unification & Primitive Substitution

- As higher-order unification was still an open problem in 1971 calculus \mathcal{R} employs the British museum method instead, i.e. it provides a substitution rule that allows to blindly instantiate free variables by arbitrary terms. As the instantiated terms may contain logical constants, instantiation of variables in proper clauses may lead to pre-clauses, which must be normalised again with the clause normalisation rules.

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Unification & Primitive Substitution

- As higher-order unification was still an open problem in 1971 calculus \mathcal{R} employs the British museum method instead, i.e. it provides a substitution rule that allows to blindly instantiate free variables by arbitrary terms. As the instantiated terms may contain logical constants, instantiation of variables in proper clauses may lead to pre-clauses, which must be normalised again with the clause normalisation rules.

- Substitution of arbitrary terms:

$$\frac{C}{C_{[T_\alpha/X_\alpha]}} \text{ Sub}$$

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Unification & Primitive Substitution

- As higher-order unification was still an open problem in 1971 calculus \mathcal{R} employs the British museum method instead, i.e. it provides a substitution rule that allows to blindly instantiate free variables by arbitrary terms. As the instantiated terms may contain logical constants, instantiation of variables in proper clauses may lead to pre-clauses, which must be normalised again with the clause normalisation rules.

- Substitution of arbitrary terms:

$$\frac{\mathcal{C}}{\mathcal{C}_{[\mathbf{T}_\alpha/\mathbf{X}_\alpha]}} \text{ Sub}$$

\mathbf{X}_α is a free variable occurring in \mathcal{C} .

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Extensionality Treatment

- Calculus \mathcal{R} does not provide rules addressing the functional and/or Boolean extensionality principles.

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Extensionality Treatment

- Calculus \mathcal{R} does not provide rules addressing the functional and/or Boolean extensionality principles.
- Instead \mathcal{R} assumes that the following extensionality axioms are (in form of respective clauses) explicitly added to the search space. And since the functional extensionality principle is parameterised over arbitrary functional types infinitely many functional extensionality axioms are required.

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Extensionality Treatment

- Calculus \mathcal{R} does not provide rules addressing the functional and/or Boolean extensionality principles.
- Instead \mathcal{R} assumes that the following extensionality axioms are (in form of respective clauses) explicitly added to the search space. And since the functional extensionality principle is parameterised over arbitrary functional types infinitely many functional extensionality axioms are required.
- Extensionality axioms

$$\mathbf{EXT}_{\alpha \rightarrow \beta}^{\doteq}: \quad \forall F_{\alpha \rightarrow \beta}. \forall G_{\alpha \rightarrow \beta}. (\forall X_{\beta}. F X \doteq G X) \Rightarrow F \doteq G$$

$$\mathbf{EXT}_{\circ}^{\doteq}: \quad \forall A_{\circ}. \forall B_{\circ}. (A \Leftrightarrow B) \Rightarrow A \doteq^{\circ} B$$

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Extensionality Treatment (contd.)

- The extensionality clauses derived from the extensionality axioms have the following form (note the many free variables, especially at literal head position, that are introduced into the search space – they heavily increase the amount of blind search in any attempt to automate the calculus):

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Extensionality Treatment (contd.)

- The extensionality clauses derived from the extensionality axioms have the following form (note the many free variables, especially at literal head position, that are introduced into the search space – they heavily increase the amount of blind search in any attempt to automate the calculus):

$$\begin{aligned}\mathcal{E}_1^{\alpha \rightarrow \beta} &: [\text{p } (\mathbf{F} \text{ s})]^T \vee [\mathbf{Q} \mathbf{F}]^F \vee [\mathbf{Q} \mathbf{G}]^T \\ \mathcal{E}_2^{\alpha \rightarrow \beta} &: [\text{p } (\mathbf{G} \text{ s})]^F \vee [\mathbf{Q} \mathbf{F}]^F \vee [\mathbf{Q} \mathbf{G}]^T\end{aligned}$$

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Extensionality Treatment (contd.)

- The extensionality clauses derived from the extensionality axioms have the following form (note the many free variables, especially at literal head position, that are introduced into the search space – they heavily increase the amount of blind search in any attempt to automate the calculus):

$$\mathcal{E}_1^{\alpha \rightarrow \beta} : [p (\mathbf{F} s)]^T \vee [\mathbf{Q} \mathbf{F}]^F \vee [\mathbf{Q} \mathbf{G}]^T$$

$$\mathcal{E}_2^{\alpha \rightarrow \beta} : [p (\mathbf{G} s)]^F \vee [\mathbf{Q} \mathbf{F}]^F \vee [\mathbf{Q} \mathbf{G}]^T$$

$$\mathcal{E}_1^o : [\mathbf{A}]^F \vee [\mathbf{B}]^F \vee [\mathbf{P} \mathbf{A}]^F \vee [\mathbf{P} \mathbf{B}]^T$$

$$\mathcal{E}_2^o : [\mathbf{A}]^T \vee [\mathbf{B}]^T \vee [\mathbf{P} \mathbf{A}]^F \vee [\mathbf{P} \mathbf{B}]^T$$

Andrews' Higher-Order Resolution \mathcal{R} ATP in FOL and HOL

Extensionality Treatment (contd.)

- The extensionality clauses derived from the extensionality axioms have the following form (note the many free variables, especially at literal head position, that are introduced into the search space – they heavily increase the amount of blind search in any attempt to automate the calculus):

$$\mathcal{E}_1^{\alpha \rightarrow \beta} : [p (\mathbf{F} s)]^T \vee [\mathbf{Q} \mathbf{F}]^F \vee [\mathbf{Q} \mathbf{G}]^T$$

$$\mathcal{E}_2^{\alpha \rightarrow \beta} : [p (\mathbf{G} s)]^F \vee [\mathbf{Q} \mathbf{F}]^F \vee [\mathbf{Q} \mathbf{G}]^T$$

$$\mathcal{E}_1^o : [\mathbf{A}]^F \vee [\mathbf{B}]^F \vee [\mathbf{P} \mathbf{A}]^F \vee [\mathbf{P} \mathbf{B}]^T$$

$$\mathcal{E}_2^o : [\mathbf{A}]^T \vee [\mathbf{B}]^T \vee [\mathbf{P} \mathbf{A}]^F \vee [\mathbf{P} \mathbf{B}]^T$$

$p_{\beta \rightarrow o}, s_\alpha$ are Skolem terms and $A_o, B_o, P_{o \rightarrow o}, Q_{(\alpha \rightarrow \beta) \rightarrow o}$ are new free variables.

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Proof Search

- initially the proof problem is negated and normalised

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Proof Search

- initially the proof problem is negated and normalised
- proof search then starts with the normalised clauses and applies the cut and simplification rule in close connection with the substitution rule

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Proof Search

- initially the proof problem is negated and normalised
- proof search then starts with the normalised clauses and applies the cut and simplification rule in close connection with the substitution rule
- intermediate applications of the clause normalisation rules may be needed to normalise temporarily generated pre-clauses

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Proof Search


- initially the proof problem is negated and normalised
- proof search then starts with the normalised clauses and applies the cut and simplification rule in close connection with the substitution rule
- intermediate applications of the clause normalisation rules may be needed to normalise temporarily generated pre-clauses
- the extensionality treatment in \mathcal{R} simply assumes to add at the beginning of the refutation process the above clauses obtained from the extensionality axioms

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Proof Search

- initially the proof problem is negated and normalised
- proof search then starts with the normalised clauses and applies the cut and simplification rule in close connection with the substitution rule
- intermediate applications of the clause normalisation rules may be needed to normalise temporarily generated pre-clauses
- the extensionality treatment in \mathcal{R} simply assumes to add at the beginning of the refutation process the above clauses obtained from the extensionality axioms
- the proof search can be graphically illustrated as follows:



Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Completeness

- [Andrews71] gives a completeness proof for calculus \mathcal{R} with respect to the semantical notion of V-complexes (corresponds to our weakest model class $\mathfrak{M}_\beta(\Sigma)$)

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Completeness

- [Andrews71] gives a completeness proof for calculus \mathcal{R} with respect to the semantical notion of V-complexes (corresponds to our weakest model class $\mathfrak{M}_\beta(\Sigma)$)
- as the extensionality principles are not valid in this rather weak semantical structures, the extensionality axioms are not needed in this completeness proof

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Completeness

- [Andrews71] gives a completeness proof for calculus \mathcal{R} with respect to the semantical notion of V-complexes (corresponds to our weakest model class $\mathfrak{M}_\beta(\Sigma)$)
- as the extensionality principles are not valid in this rather weak semantical structures, the extensionality axioms are not needed in this completeness proof
- Theorem: (V-completeness of \mathcal{R}) The calculus \mathcal{R} is (sound and) complete with respect to the notion of V-complexes.

Proof: [Andrews71].

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Henkin Completeness

- We can also prove Henkin completeness of calculus \mathcal{R} .

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Henkin Completeness

- We can also prove Henkin completeness of calculus \mathcal{R} .
- Theorem: (Henkin completeness of \mathcal{R}) The calculus \mathcal{R} is (sound and) complete with respect to Henkin semantics provided that the infinitely many extensionality axioms are given.

Proof: exercise

Andrews' Higher-Order Resolution \mathcal{R}

ATP in FOL and HOL

Henkin Completeness

- We can also prove Henkin completeness of calculus \mathcal{R} .
- Theorem: (Henkin completeness of \mathcal{R}) The calculus \mathcal{R} is (sound and) complete with respect to Henkin semantics provided that the infinitely many extensionality axioms are given.

Proof: exercise

Exercise: How are the following theorems proved in calculus \mathcal{R} ?

- Leibniz equality and η -equality:

$$f_{\iota \rightarrow \iota} \doteq \lambda X_{\iota}. f X$$

Exercise: How are the following theorems proved in calculus \mathcal{R} ?

- Leibniz equality and η -equality:

$$f_{\iota \rightarrow \iota} \doteq \lambda X_{\iota}. f X$$

- The set of all red balls equals the set of all balls that are red:
 $\{X | \text{red } X \wedge \text{ball } X\} = \{X | \text{ball } X \wedge \text{red } X\}$. This problem can be encoded as

$$(\lambda X_{\iota}. \text{red } X \wedge \text{ball } X) = (\lambda X_{\iota}. \text{ball } X \wedge \text{red } X)$$

Exercise: How are the following theorems proved in calculus \mathcal{R} ?

- All unary logical operators $O_{o \rightarrow o}$ which map the propositions a and b to \top consequently also map $a \wedge b$ to \top :

$$\forall O_{o \rightarrow o}. (O a_o) \wedge (O b_o) \Rightarrow (O (a_o \wedge b_o))$$

Exercise: How are the following theorems proved in calculus \mathcal{R} ?

- In Henkin semantics the domain \mathcal{D}_o of all Booleans contains exactly the truth values \perp and \top . Consequently the domain of all mappings from Booleans to Booleans contains exactly contains in each Henkin model at most four elements. And because of the requirement, that the function domains in Henkin models must be rich enough such that every term has a denotation, it follows that $\mathcal{D}_{o \rightarrow o}$ contains exactly the pairwise distinct denotations of the following four terms: $\lambda X_o.X_o$, $\lambda X_o.\neg X_o$, $\lambda X_o.\perp$, and $\lambda X_o.\top$. This theorem can be formulated as follows (where $f_{o \rightarrow o}$ is a constant):

$$(f = \lambda X_o.X_o) \vee (f = \lambda X_o.\neg X_o) \vee (f = \lambda X_o.\perp) \vee (f = \lambda X_o.\top)$$



Approaches to Higher-Order
Resolution: Huet's CR

Huet's Constrained Resolution \mathcal{CR}

ATP in FOL and HOL

We transform Huet's constrained resolution approach [Huet72,Huet73]. The calculus here is the unsorted fragment of the variant of Huet's approach as presented in [Kohlhase94]. In the remainder of this paper we refer to this calculus with \mathcal{CR} .

λ -Conversion

- Calculus \mathcal{CR} assumes that terms, literals, and clauses are implicitly reduced to β -normal form.

We transform Huet's constrained resolution approach [Huet72,Huet73]. The calculus here is the unsorted fragment of the variant of Huet's approach as presented in [Kohlhase94]. In the remainder of this paper we refer to this calculus with \mathcal{CR} .

λ -Conversion

- Calculus \mathcal{CR} assumes that terms, literals, and clauses are implicitly reduced to β -normal form.
- Furthermore, we assume that α -equality is treated implicitly, i.e. we identify all terms that differ only with respect to the names of bound variables.

Clause Normalisation

- [Huet72] does not explicitly present clause normalisation rules but assumes that they are given. Here we employ the rules \neg^T , \neg^F , \vee^T , \vee_l^F , \vee_r^F , \sqcap^T , and \sqcap^F as already defined for calculus \mathcal{R} before.

Clause Normalisation

- [Huet72] does not explicitly present clause normalisation rules but assumes that they are given. Here we employ the rules \neg^T , \neg^F , \vee^T , \vee_l^F , \vee_r^F , \sqcap^T , and \sqcap^F as already defined for calculus \mathcal{R} before.

Huet's Constrained Resolution CR

ATP in FOL and HOL

Resolution & Factorisation

- As first-order unification is decidable and unitary it can be employed as a strong filter in first-order resolution [Robinson65].

Resolution & Factorisation

- As first-order unification is decidable and unitary it can be employed as a strong filter in first-order resolution [Robinson65].
- Unfortunately higher-order unification is not decidable (cf. [Lucchesi72,Huet73,Goldfarb81]) and thus it can not be applied in the sense of a terminating side computation in higher-order theorem proving.

Resolution & Factorisation

- As first-order unification is decidable and unitary it can be employed as a strong filter in first-order resolution [Robinson65].
- Unfortunately higher-order unification is not decidable (cf. [Lucchesi72,Huet73,Goldfarb81]) and thus it can not be applied in the sense of a terminating side computation in higher-order theorem proving.
- Huet therefore suggests in [Huet72,Huet73] to delay the unification process and to explicitly encode unification problems occurring during the refutation search as unification constraints.

Resolution & Factorisation (contd.)

- In his original approach Huet presented a hyper-resolution rule which simultaneously resolves on the resolution literals A^1, \dots, A^n ($1 \leq n$) and B^1, \dots, B^m ($1 \leq m$) of two given clauses and adds the unification constraint $[\neq^? (A^1, \dots, A^n, B^1, \dots, B^m)]$ to the resolvent:

$$\frac{[A^1]^\mu \vee \dots \vee [A^n]^\mu \vee C \quad [B^1]^\nu \vee \dots \vee [B^m]^\nu \vee D}{C \vee D \vee [\neq^? (A^1, \dots, A^n, B^1, \dots, B^m)]} \text{Hres}$$

(where $\mu \neq \nu$).

Resolution & Factorisation (contd.)

- In order to ease the comparison with the two other approaches discussed in this paper we instead employ a resolution rule Res and a factorisation rule Fac.

Resolution & Factorisation (contd.)

- In order to ease the comparison with the two other approaches discussed in this paper we instead employ a resolution rule Res and a factorisation rule Fac.
- Like Hres both rules encode the unification problem to be solved as a unification constraint:

Resolution & Factorisation (contd.)

- In order to ease the comparison with the two other approaches discussed in this paper we instead employ a resolution rule Res and a factorisation rule Fac.
- Like Hres both rules encode the unification problem to be solved as a unification constraint:

- Constrained resolution:

$$\frac{[A]^{\mu} \vee C \quad [B]^{\nu} \vee D}{C \vee D \vee [A \neq^? B]} \text{ Res}$$

(where $\mu \neq \nu$).

Resolution & Factorisation (contd.)

- In order to ease the comparison with the two other approaches discussed in this paper we instead employ a resolution rule Res and a factorisation rule Fac.
- Like Hres both rules encode the unification problem to be solved as a unification constraint:

- Constrained resolution:

$$\frac{[A]^{\mu} \vee C \quad [B]^{\nu} \vee D}{C \vee D \vee [A \neq^? B]} \text{ Res}$$

(where $\mu \neq \nu$).

- Constrained factorisation:

$$\frac{[A]^{\mu} \vee [B]^{\mu} \vee C}{[A]^{\mu} \vee C \vee [A \neq^? B]^F} \text{ Fac}$$

Resolution & Factorisation (contd.)

- One can easily prove by induction on $n + m$ that each proof step applying rule Hres can be replaced by a corresponding derivation employing Res and Fac.

Resolution & Factorisation (contd.)

- One can easily prove by induction on $n + m$ that each proof step applying rule Hres can be replaced by a corresponding derivation employing Res and Fac.
- For a formal proof note that the unification constraint $[\neq^? (\mathbf{A}^1, \dots \mathbf{A}^n, \mathbf{B}^1, \dots \mathbf{B}^m)]$ is equivalent to $[\mathbf{A}^1 \neq^? \mathbf{A}^2] \vee [\mathbf{A}^2 \neq^? \mathbf{A}^3] \vee \dots \vee [\mathbf{A}^{n-1} \neq^? \mathbf{A}^n] \vee [\mathbf{A}^n \neq^? \mathbf{B}^1] \vee [\mathbf{B}^1 \neq^? \mathbf{B}^2] \vee [\mathbf{B}^2 \neq^? \mathbf{B}^3] \vee \dots \vee [\mathbf{B}^{n-1} \neq^? \mathbf{B}^n]$.

Unification & Splitting

- [Huet75] introduces higher-order unification and higher-order pre-unification and shows that higher-order pre-unification is sufficient to verify the soundness of a refutation in which the occurring unification problems have been delayed until the end.

Unification & Splitting

- [Huet75] introduces higher-order unification and higher-order pre-unification and shows that higher-order pre-unification is sufficient to verify the soundness of a refutation in which the occurring unification problems have been delayed until the end.
- The higher-order pre-unification rules presented here are discussed in detail in [Benzmüller-PhD-99]. They furthermore closely reflect the rules as presented in [SnyderGallier89].

Unification & Splitting

- [Huet75] introduces higher-order unification and higher-order pre-unification and shows that higher-order pre-unification is sufficient to verify the soundness of a refutation in which the occurring unification problems have been delayed until the end.
- The higher-order pre-unification rules presented here are discussed in detail in [Benzmüller-PhD-99]. They furthermore closely reflect the rules as presented in [SnyderGallier89].
- Elimination of trivial pairs:

$$\frac{C \vee [A \neq^? A]}{C} \text{ Triv}$$

Unification & Splitting

- [Huet75] introduces higher-order unification and higher-order pre-unification and shows that higher-order pre-unification is sufficient to verify the soundness of a refutation in which the occurring unification problems have been delayed until the end.
- The higher-order pre-unification rules presented here are discussed in detail in [Benzmüller-PhD-99]. They furthermore closely reflect the rules as presented in [SnyderGallier89].

- Elimination of trivial pairs:

$$\frac{C \vee [A \neq^? A]}{C} \text{ Triv}$$
- Decomposition

$$\frac{C \vee [h\overline{U}^n \neq^? h\overline{V}^n]}{C \vee [U^1 \neq^? V^1] \vee \dots \vee [U^n \neq^? V^n]} \text{ Dec}$$

Unification & Splitting (contd.)

Elimination of λ -binders:

- (weak functional extensionality)

$$\frac{C \vee [M_{\alpha \rightarrow \beta} \neq^? N_{\alpha \rightarrow \beta}]}{C \vee [M s_{\alpha} \neq^? N s_{\alpha}]} \text{Func}$$

Unification & Splitting (contd.)

Elimination of λ -binders:

- (weak functional extensionality)

$$\frac{C \vee [M_{\alpha \rightarrow \beta} \neq^? N_{\alpha \rightarrow \beta}]}{C \vee [M s_{\alpha} \neq^? N s_{\alpha}]} \text{Func}$$

s_{α} is a new Skolem term.

Unification & Splitting (contd.)

Elimination of λ -binders:

- (weak functional extensionality)

$$\frac{C \vee [M_{\alpha \rightarrow \beta} \neq^? N_{\alpha \rightarrow \beta}]}{C \vee [M s_{\alpha} \neq^? N s_{\alpha}]} \text{ Func}$$

s_{α} is a new Skolem term.

- Imitation of rigid heads:

$$\frac{C \vee [F_{\gamma} \overline{U}^n \neq^? h \overline{V}^m] \quad G \in \mathcal{AB}_{\gamma}^h}{C \vee [F \neq^? G] \vee [F \overline{U}^n \neq^? h \overline{V}^m]} \text{ FlexRigid}$$

Unification & Splitting (contd.)

Elimination of λ -binders:

- (weak functional extensionality)

$$\frac{C \vee [M_{\alpha \rightarrow \beta} \neq^? N_{\alpha \rightarrow \beta}]}{C \vee [M s_{\alpha} \neq^? N s_{\alpha}]} \text{ Func}$$

s_{α} is a new Skolem term.

- Imitation of rigid heads:
$$\frac{C \vee [F_{\gamma} \overline{U}^n \neq^? h \overline{V}^m] \quad G \in \mathcal{AB}_{\gamma}^h}{C \vee [F \neq^? G] \vee [F \overline{U}^n \neq^? h \overline{V}^m]} \text{ FlexRigid}$$

\mathcal{AB}_{γ}^h is the set of general bindings of type γ for head h .

Unification & Splitting (contd.)

- Huet points to the usefulness of eager unification to filter out clauses with non-unifiable unification constraints or to back-propagate the solutions of easily solvable constraints (e.g., in case of first-order unification problems occurring during the proof search): many of the higher-order unification problems occurring in practice are decidable and have only finitely many solutions.

Unification & Splitting (contd.)

- Huet points to the usefulness of eager unification to filter out clauses with non-unifiable unification constraints or to back-propagate the solutions of easily solvable constraints (e.g., in case of first-order unification problems occurring during the proof search): many of the higher-order unification problems occurring in practice are decidable and have only finitely many solutions.
- Hence, even though higher-order unification is generally not decidable it is sensible in practice to apply the unification algorithm with a particular resource, such that only those unification problems which may have further solutions beyond this bound need to be delayed.

Unification & Splitting (contd.)

- In our presentation of calculus \mathcal{CR} we explicitly address the aspect of eager unification and substitution by rule Subst. This rule back-propagates eagerly computed unifiers to the literal part of a clause.

Unification & Splitting (contd.)

- In our presentation of calculus \mathcal{CR} we explicitly address the aspect of eager unification and substitution by rule Subst. This rule back-propagates eagerly computed unifiers to the literal part of a clause.
- Eager unification & substitution:

$$\frac{C \vee [X \neq^? A] \quad X \notin \text{free}(A)}{C_{[A/X]}} \text{Subst}$$

Unification & Splitting (contd.)

- The literal heads of our clauses may consist of set variables and it may be necessary to instantiate them with terms introducing new logical constant at head position in order to find a refutation.

Unification & Splitting (contd.)

- The literal heads of our clauses may consist of set variables and it may be necessary to instantiate them with terms introducing new logical constant at head position in order to find a refutation.
- Unfortunately not all appropriate instantiations can be computed with the calculus rules presented so far.

Unification & Splitting (contd.)

- The literal heads of our clauses may consist of set variables and it may be necessary to instantiate them with terms introducing new logical constant at head position in order to find a refutation.
- Unfortunately not all appropriate instantiations can be computed with the calculus rules presented so far.
- To address this problem Huet's approach provides the following splitting rules:

Huet's Constrained Resolution CR

ATP in FOL and HOL

Unification & Splitting (contd.)

- Instantiate
set variables:

Huet's Constrained Resolution \mathcal{CR}

ATP in FOL and HOL

Unification & Splitting (contd.)

- Instantiate set variables:

$$\frac{[P \ A]^T \vee C}{[Q]^T \vee [R]^T \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^T$$

Huet's Constrained Resolution \mathcal{CR}

ATP in FOL and HOL

Unification & Splitting (contd.)

- Instantiate set variables:

$$\frac{[P \ A]^{\mu} \vee C}{[Q]^{\nu} \vee C \vee [P \ A] \neq^? \neg Q_o} S_{\neg}^{TF} \quad (\text{where } \mu \neq \nu)$$

$$\frac{[P \ A]^T \vee C}{[Q]^T \vee [R]^T \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^T$$

$$\frac{[P \ A]^F \vee C}{[Q]^F \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^F$$

$$[R]^F \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)$$

Huet's Constrained Resolution \mathcal{CR}

ATP in FOL and HOL

Unification & Splitting (contd.)

- Instantiate set variables:

$$\frac{[P \ A]^{\mu} \vee C}{[Q]^{\nu} \vee C \vee [P \ A] \neq^? \neg Q_o} S_{\neg}^{TF} \quad (\text{where } \mu \neq \nu)$$

$$\frac{[P \ A]^T \vee C}{[Q]^T \vee [R]^T \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^T$$

$$\frac{[P \ A]^F \vee C}{[Q]^F \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^F$$

$$[R]^F \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)$$

$$\frac{[P \ A_{\alpha \rightarrow o}]^T \vee C}{[M_{\alpha \rightarrow o} \ Z]^T \vee C \vee [P \ A] \neq^? \Pi^{\alpha} M} S_{\Pi}^T$$

Huet's Constrained Resolution \mathcal{CR}

ATP in FOL and HOL

Unification & Splitting (contd.)

- Instantiate set variables:

$$\frac{[P \ A]^{\mu} \vee C}{[Q]^{\nu} \vee C \vee [P \ A] \neq^? \neg Q_o} S_{\neg}^{TF} \quad (\text{where } \mu \neq \nu)$$

$$\frac{[P \ A]^T \vee C}{[Q]^T \vee [R]^T \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^T$$

$$\frac{[P \ A]^F \vee C}{[Q]^F \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^F$$

$$[R]^F \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)$$

$$\frac{[P \ A_{\alpha \rightarrow o}]^T \vee C}{[M_{\alpha \rightarrow o} \ Z]^T \vee C \vee [P \ A] \neq^? \Pi^{\alpha} M} S_{\Pi}^T$$

$$\frac{[P \ A_{\alpha \rightarrow o}]^F \vee C}{[M_{\alpha \rightarrow o} \ s]^F \vee C \vee [P \ A] \neq^? \Pi^{\alpha} M} S_{\Pi}^F$$

Unification & Splitting (contd.)

- Instantiate set variables:

$$\frac{[P \ A]^T \vee C}{[Q]^T \vee [R]^T \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^T$$

$$\frac{[P \ A]^\mu \vee C}{[Q]^\nu \vee C \vee [P \ A] \neq^? \neg Q_o} S_{\neg}^{TF} \quad (\text{where } \mu \neq \nu)$$

$$\frac{[P \ A]^F \vee C}{[Q]^F \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^F$$

$$\frac{[P \ A]^F \vee C}{[R]^F \vee C \vee [P \ A] \neq^? (Q_o \vee R_o)} S_V^F$$

$$\frac{[P \ A_{\alpha \rightarrow o}]^T \vee C}{[M_{\alpha \rightarrow o} \ Z]^T \vee C \vee [P \ A] \neq^? \Pi^\alpha M} S_{\Pi}^T$$

$$\frac{[P \ A_{\alpha \rightarrow o}]^F \vee C}{[M_{\alpha \rightarrow o} \ s]^F \vee C \vee [P \ A] \neq^? \Pi^\alpha M} S_{\Pi}^F$$

- S_{Π}^T and S_{Π}^F are infinitely branching as they are parameterised over type α . $Q_o, R_o, M_{\alpha \rightarrow o}, Z_\alpha$ are new variables and s_α is a new Skolem constant.

Huet's Constrained Resolution \mathcal{CR}

ATP in FOL and HOL

Unification & Splitting (contd.)

- A theorem which is not refutable in \mathcal{CR} if the splitting rules are not available is $\exists A_o.A$:

Unification & Splitting (contd.)

- A theorem which is not refutable in \mathcal{CR} if the splitting rules are not available is $\exists A_o.A$:
- After negation this statement normalises to clause $\mathcal{C}_1 : [A]^F$, such that none but the splitting rules are applicable. With the help of rule S_{\neg}^{TF} and eager unification, however, we can derive $\mathcal{C}_2 : [A']^T$ which is then successfully resolvable against \mathcal{C}_1 .

Extensionality Treatment

- On the one hand η -convertibility is built-in in higher-order unification, such that calculus \mathcal{CR} already supports functional extensionality reasoning to a certain extend.

Extensionality Treatment

- On the one hand η -convertibility is built-in in higher-order unification, such that calculus \mathcal{CR} already supports functional extensionality reasoning to a certain extend.
- On the other hand \mathcal{CR} nevertheless fails to address full extensionality as it does not realise the required subtle interplay between the functional and Boolean extensionality principles.

Extensionality Treatment

- On the one hand η -convertibility is built-in in higher-order unification, such that calculus CR already supports functional extensionality reasoning to a certain extend.
- On the other hand CR nevertheless fails to address full extensionality as it does not realise the required subtle interplay between the functional and Boolean extensionality principles.
- Without employing additional (Boolean and functional!) extensionality axioms CR is, e.g., not able to prove the rather simple examples presented before.

Proof Search

- Initially the proof problem is negated and normalised. The main proof search then operates on the generated clauses by applying the resolution, factorisation, and splitting rules.

Proof Search

- Initially the proof problem is negated and normalised. The main proof search then operates on the generated clauses by applying the resolution, factorisation, and splitting rules.
- Despite the possibility of eager unification CR generally foresees to delay the higher-order unification process in order to overcome the undecidability problem.

Proof Search

- Initially the proof problem is negated and normalised. The main proof search then operates on the generated clauses by applying the resolution, factorisation, and splitting rules.
- Despite the possibility of eager unification CR generally foresees to delay the higher-order unification process in order to overcome the undecidability problem.
- When deriving a potentially empty clause (no normal literals), CR then tests whether the accumulated unification constraints justifying this particular refutation are solvable.

Huet's Constrained Resolution CR


ATP in FOL and HOL

Proof Search (contd.)

- Like \mathcal{R} , the extensionality treatment of CR requires to add infinitely many extensionality axioms to the search space.

Proof Search (contd.)

- Like \mathcal{R} , the extensionality treatment of CR requires to add infinitely many extensionality axioms to the search space.
- The following figure graphically illustrates the main ideas of the proof search in CR .



Completeness Results

- [Huet72,Huet73] analyses completeness of \mathcal{CR} formally only with respect to Andrews V-complexes, i.e. Huet verifies that the set of non-refutable sentences in \mathcal{CR} is an abstract consistency class for V-complexes.

Completeness Results

- [Huet72,Huet73] analyses completeness of \mathcal{CR} formally only with respect to Andrews V-complexes, i.e. Huet verifies that the set of non-refutable sentences in \mathcal{CR} is an abstract consistency class for V-complexes.
- Theorem (V-completeness of \mathcal{CR}): The calculus \mathcal{CR} is complete with respect to the notion of V-complexes.

Proof: [Huet72,Huet73]

Completeness Results

- [Huet72,Huet73] analyses completeness of \mathcal{CR} formally only with respect to Andrews V-complexes, i.e. Huet verifies that the set of non-refutable sentences in \mathcal{CR} is an abstract consistency class for V-complexes.
- Theorem (V-completeness of \mathcal{CR}): The calculus \mathcal{CR} is complete with respect to the notion of V-complexes.

Proof: [Huet72,Huet73]

- Theorem (Henkin completeness of \mathcal{CR}): The calculus \mathcal{CR} is complete wrt. Henkin semantics provided that the infinitely many extensionality axioms are given.

Proof: exercise

Exercise: How are the following theorems proved in calculus \mathcal{CR} ?

- Leibniz equality and η -equality:

$$f_{\iota \rightarrow \iota} \doteq \lambda X_{\iota}. f X$$

Exercise: How are the following theorems proved in calculus \mathcal{CR} ?

- Leibniz equality and η -equality:

$$f_{\iota \rightarrow \iota} \doteq \lambda X_{\iota}. f X$$

- The set of all red balls equals the set of all balls that are red:
 $\{X | \text{red } X \wedge \text{ball } X\} = \{X | \text{ball } X \wedge \text{red } X\}$. This problem can be encoded as

$$(\lambda X_{\iota}. \text{red } X \wedge \text{ball } X) = (\lambda X_{\iota}. \text{ball } X \wedge \text{red } X)$$

Exercise: How are the following theorems proved in calculus \mathcal{CR} ?

- All unary logical operators $O_{o \rightarrow o}$ which map the propositions a and b to \top consequently also map $a \wedge b$ to \top :

$$\forall O_{o \rightarrow o}. (O a_o) \wedge (O b_o) \Rightarrow (O (a_o \wedge b_o))$$

Exercise: How are the following theorems proved in calculus \mathcal{CR} ?

- In Henkin semantics the domain \mathcal{D}_o of all Booleans contains exactly the truth values \perp and \top . Consequently the domain of all mappings from Booleans to Booleans contains exactly contains in each Henkin model at most four elements. And because of the requirement, that the function domains in Henkin models must be rich enough such that every term has a denotation, it follows that $\mathcal{D}_{o \rightarrow o}$ contains exactly the pairwise distinct denotations of the following four terms: $\lambda X_o.X_o$, $\lambda X_o.\neg X_o$, $\lambda X_o.\perp$, and $\lambda X_o.\top$. This theorem can be formulated as follows (where $f_{o \rightarrow o}$ is a constant):

$$(f = \lambda X_o.X_o) \vee (f = \lambda X_o.\neg X_o) \vee (f = \lambda X_o.\perp) \vee (f = \lambda X_o.\top)$$



Approaches to Higher-Order
Resolution: Benzmüller's *ER*

Clause normalization

$$\begin{array}{c}
 \frac{C \vee [A \vee B]^T}{C \vee [A]^T \vee [B]^T} \vee^T \quad \frac{C \vee [A \vee B]^F}{C \vee [A]^F} \vee_I^F \quad \frac{C \vee [A \vee B]^F}{C \vee [B]^F} \vee_r^F \\
 \\
 \frac{C \vee [\neg A]^T}{C \vee [A]^F} \neg^T \quad \frac{C \vee [\neg A]^F}{C \vee [A]^T} \neg^F \\
 \\
 \frac{C \vee [\Pi^\alpha A]^T \quad X_\alpha \text{ new variable}}{C \vee [A X]^T} \Pi^T \\
 \\
 \frac{C \vee [\Pi^\alpha A]^F \quad \text{sk}_\alpha \text{ Skolem term}}{C \vee [A \text{ sk}_\alpha]^F} \Pi^F
 \end{array}$$

This rules may be combined into a single rule Cnf .

Resolution and Factorisation

$$\frac{[N]^\alpha \vee C \quad [M]^\beta \vee D \quad \alpha \neq \beta}{C \vee D \vee [N \neq^? M]} \text{ Res}$$

$$\frac{[N]^\alpha \vee [M]^\alpha \vee C \quad \alpha \in \{T, F\}}{[N]^\alpha \vee C \vee [N \neq^? M]} \text{ Fac}$$

$$\frac{[Q_\gamma \overline{U^k}]^\alpha \vee C \quad P \in \mathcal{GB}_\gamma^{\{\neg, \vee\} \cup \{\Pi^\beta \mid \beta \in \mathcal{T}^k\}}}{[Q_\gamma \overline{U^k}]^\alpha \vee C \vee [Q \neq^? P]} \text{ Prim}^k$$

(Pre-)unification rules

$$\frac{C \vee [M_{\alpha \rightarrow \beta} \neq^? N_{\alpha \rightarrow \beta}] \quad s_\alpha \text{ Skolem-Term}}{C \vee [M s \neq^? N s]} \text{ Func}$$

$$\frac{C \vee [h\overline{U}^n \neq^? h\overline{V}^n]}{C \vee [U^1 \neq^? V^1] \vee \dots \vee [U^n \neq^? V^n]} \text{ Dec} \quad \frac{C \vee [A \neq^? A]}{C} \text{ Triv}$$

$$\frac{C \vee [F_\gamma \overline{U}^n \neq^? h\overline{V}^n] \quad G \in \mathcal{GB}_\gamma^h}{C \vee [F \neq^? G] \vee [F\overline{U}^n \neq^? h\overline{V}^n]} \text{ Flex/Rigid}$$

$$\frac{C \vee E \quad E \text{ solved for } C}{\text{Cnf}(\text{subst}_E(C))} \text{ Subst}$$

Extensionality rules

$$\frac{C \vee [M_o \neq^? N_o]}{\text{Cnf}(C \vee [M_o \Leftrightarrow N_o]^F)} \text{Equiv}$$

$$\frac{C \vee [M_\alpha \neq^? N_\alpha] \quad \alpha \in \{o, \iota\}}{\text{Cnf}(C \vee [\forall P_{\alpha \rightarrow o}. PM \Rightarrow PN]^F)} \text{Leib}$$

Extensionality Treatment

- Instead of adding infinitely many extensionality axioms to the search space \mathcal{CR} provides two new extensionality rules which closely connect refutation search and eager unification.

Extensionality Treatment

- Instead of adding infinitely many extensionality axioms to the search space \mathcal{CR} provides two new extensionality rules which closely connect refutation search and eager unification.
- The idea is to allow for recursive calls from higher-order unification to the overall refutation process.

Extensionality Treatment


- Instead of adding infinitely many extensionality axioms to the search space \mathcal{CR} provides two new extensionality rules which closely connect refutation search and eager unification.
- The idea is to allow for recursive calls from higher-order unification to the overall refutation process.
- This turns the rather weak syntactical higher-order unification approach considered so far into a most general approach for *dynamic* higher-order theory unification.

Proof Search

- Initially the proof problem is negated and normalised. The main proof search then closely interleaves the refutation process on resolution layer and unification, i.e. the main proof search rules Res, Fac, and Prim and the unification rules are integrated at a common conceptual level. The calls from unification to the overall refutation process with rules *Leib* and *Equiv* introduce new clauses into the search space which can be resolved against already given ones.

Proof Search

- Initially the proof problem is negated and normalised. The main proof search then closely interleaves the refutation process on resolution layer and unification, i.e. the main proof search rules Res, Fac, and Prim and the unification rules are integrated at a common conceptual level. The calls from unification to the overall refutation process with rules *Leib* and *Equiv* introduce new clauses into the search space which can be resolved against already given ones.
- The following figure graphically illustrates the main ideas of the proof search in \mathcal{ER} .



Ex.: Extensional HO Resolution \mathcal{ER}

ATP in FOL and HOL

$$\forall P_{o \rightarrow o}. (P a_o) \wedge (P b_o) \Rightarrow (P (a_o \wedge b_o))$$

Negation and clause normalization

$$\mathcal{C}_1 : [p a]^T \quad \mathcal{C}_2 : [p b]^T \quad \mathcal{C}_3 : [p (a \wedge b)]^F$$

Resolution between \mathcal{C}_1 and \mathcal{C}_3 and between \mathcal{C}_2 and \mathcal{C}_3

$$\mathcal{C}_4 : [p a \neq^? p (a \wedge b)] \quad \mathcal{C}_5 : [p b \neq^? p (a \wedge b)]$$

Decomposition

$$\mathcal{C}_6 : [a \neq^? (a \wedge b)] \quad \mathcal{C}_7 : [b \neq^? (a \wedge b)]$$

Recursive call of proof process with rules Equiv and Cnf

$$\mathcal{C}_8 : [a]^F \vee [b]^F \quad \mathcal{C}_9 : [a]^T \vee [b]^T \quad \mathcal{C}_{10} : [a]^T \quad \mathcal{C}_{11} : [b]^T$$

Ex.: Extensional HO Resolution \mathcal{ER}

ATP in FOL and HOL

$$\forall B_{\alpha \rightarrow o}, C_{\alpha \rightarrow o}, D_{\alpha \rightarrow o}. B \cup (C \cap D) = (B \cup C) \cap (B \cup D)$$

Negation and definition expansion with

$$\cup = \lambda A_{\alpha \rightarrow o}, B_{\alpha \rightarrow o}, X_{\alpha}. (A X) \vee (B X) \quad \cap = \lambda A_{\alpha \rightarrow o}, B_{\alpha \rightarrow o}, X_{\alpha}. (A X) \wedge (B X)$$

leads to:

$$C_1 : [\lambda X_{\alpha}. (b X) \vee ((c X) \wedge (d X)) \neq? \lambda X_{\alpha}. ((b X) \vee (c X)) \wedge ((b X) \vee (d X))]$$

Goal directed functional and Boolean extensionality treatment:

$$C_2 : [(b x) \vee ((c x) \wedge (d x)) \Leftrightarrow ((b x) \vee (c x)) \wedge ((b x) \vee (d x))]^F$$

Clause normalization results then in a pure propositional, i.e. decidable, set of clauses. Only these clauses are still in the search space of LEO (in total there are 33 clauses generated and LEO finds the proof on a 2,5GHz PC in 820ms).

Similar proof in case of embedded propositions:

$$\forall P_{(\alpha \rightarrow o) \rightarrow o}, B_{\alpha \rightarrow o}, C_{\alpha \rightarrow o}, D_{\alpha \rightarrow o}. P(B \cup (C \cap D)) \Rightarrow P((B \cup C) \cap (B \cup D))$$

Ex.: Extensional HO Resolution \mathcal{ER}

ATP in FOL and HOL

Further small examples which test Henkin completeness:

$$\forall F_{o \rightarrow o}. (F \doteq \lambda X_o. X_o) \vee (F \doteq \lambda X_o. \neg X_o) \vee (F \doteq \lambda X_o. \perp) \vee (F \doteq \lambda X_o. \top)$$

$$\forall H_{o \rightarrow o}. H \perp \doteq H \ (H \top \doteq H \perp)$$

...

- LEO-II [BenzmuellerEtAl08] and LEO [Benzmueller99]:
extensional higher-order resolution (plus primitive equality
treatment) and cooperation with FO ATPs

- LEO-II [BenzmuellerEtAl08] and LEO [Benzmueller99]:
extensional higher-order resolution (plus primitive equality
treatment) and cooperation with FO ATPs
- TPS [AndrewsEtAl06]: mating method

- LEO-II [BenzmuellerEtAl08] and LEO [Benzmueller99]:
extensional higher-order resolution (plus primitive equality treatment) and cooperation with FO ATPs
- TPS [AndrewsEtAl06]: mating method
- OTTER- λ [Beeson06]: not operating simple type theory but λ -logic, which is a novel combination of λ -calculus and first-order logic; employs first-order resolution