

# The Ratio of Area to Circumference is Half the Radius

(December 26, 2025)

Pretending that we know some integral calculus but not how to measure circles, we know that the area of the first quadrant of a circle with radius  $r$  centered at the origin is

$$A = \int_0^r \sqrt{r^2 - x^2} dx$$

and its length is

$$L = \int_0^r \sqrt{1 + \left( \frac{d}{dx} (\sqrt{r^2 - x^2}) \right)^2} dx = \int_0^r \frac{r}{\sqrt{r^2 - x^2}} dx$$

This theorem connects these two integrals:

**Theorem 1.** For  $n = 2, 3, \dots$ , and  $r > 0$ ,

$$\int_0^r \frac{r^n}{\sqrt{r^n - x^n}} dx = \left(1 + \frac{2}{n}\right) \int_0^r \sqrt{r^n - x^n} dx \quad (1)$$

*Proof.* Observing that

$$\begin{aligned} \int_0^r \sqrt{r^n - x^n} dx &= \int_0^r \frac{r^n - x^n}{\sqrt{r^n - x^n}} dx \\ &= \int_0^r \frac{r^n}{\sqrt{r^n - x^n}} dx - \int_0^r \frac{x^n}{\sqrt{r^n - x^n}} dx \\ &= \int_0^r \frac{r^n}{\sqrt{1 - x^n}} dx + \int_0^r \frac{2x}{n} \cdot \frac{d}{dx} \sqrt{r^n - x^n} dx \end{aligned}$$

Applying integration by parts to the second integral on the right with  $u = 2x/n$  and  $dv = \frac{d}{dx} \sqrt{r^n - x^n} dx$  gives

$$\begin{aligned} \int_0^r \frac{2x}{n} \cdot \frac{d}{dx} \sqrt{r^n - x^n} dx &= \left[ \frac{2x}{n} \sqrt{r^n - x^n} \right]_0^r - \frac{2}{n} \int_0^r \sqrt{r^n - x^n} dx \\ &= -\frac{2}{n} \int_0^r \sqrt{r^n - x^n} dx \end{aligned}$$

and (1) follows. □

Taking  $n = 2$  gives

$$rL = \left(1 + \frac{2}{2}\right) A \text{ or } \frac{A}{L} = \frac{r}{2}$$