

# ENRICHED LAWVERE THEORIES FOR OPERATIONAL SEMANTICS

*John C. Baez*

Department of Mathematics  
University of California  
Riverside CA, USA 92521  
and  
Centre for Quantum Technologies  
National University of Singapore  
Singapore 117543

*Christian Williams*

Department of Mathematics  
University of California  
Riverside CA, USA 92521

email: baez@math.ucr.edu, cwill041@math.ucr.edu

August 13, 2019

ABSTRACT. Enriched Lawvere theories are a generalization of Lawvere theories that allow us to describe the operational semantics of formal systems. For example, a graph-enriched Lawvere theory describes structures that have a *graph* of operations of each arity, where the vertices are operations and the edges are *rewrites* between operations. Enriched theories can be used to equip systems with operational semantics, and maps between enriching categories can serve to translate between different forms of operational and denotational semantics. The Grothendieck construction lets us study all models of all enriched theories in all contexts in a single category. We illustrate these ideas with the *SKI*-combinator calculus, a variable-free version of the lambda calculus, and with Milner’s calculus of communicating processes.

## 1. INTRODUCTION

Formal systems are not always explicitly connected to how they operate in practice. Lawvere theories [16] are an excellent formalism for describing algebraic structures obeying equational laws, but they do not specify how to compute in such a structure, for example taking a complex expression and simplifying it using rewrite rules. Recall that a Lawvere theory is a category with finite products  $\mathbf{T}$  generated by a single object  $t$ , for “type”, and morphisms  $t^n \rightarrow t$  representing  $n$ -ary operations, with commutative diagrams specifying equations. There is a theory for groups, a theory for rings, and so on. We can specify algebraic structures of a given kind in some category  $\mathbf{C}$  with finite products by a product-preserving functor  $\mu: \mathbf{T} \rightarrow \mathbf{C}$ . This is a simple and elegant form of *denotational* semantics. However, Lawvere theories know nothing of *operational* semantics. Our goal here is to address this using “enriched” Lawvere theories.

In a Lawvere theory, the objects are types and the morphisms are terms; however, there are no relations between terms, only equations. The process of computing one term into another should be given by hom-objects with more structure. In operational semantics, program behavior is often specified by labelled transition systems, or labelled directed (multi)graphs [24]. The edges of such a graph represent rewrites:

$$(\lambda x.x + x \ 2) \xrightarrow{\beta} 2 + 2 \xrightarrow{+} 4$$

We can use an enhanced Lawvere theory in which, rather than merely *sets* of morphisms, there are *graphs* or perhaps *categories*. Enriched Lawvere theories are exactly for this purpose.

In a theory  $T$  enriched in a category  $V$  of some kind of “directed object”, including graphs, categories, and posets, the theory has the following interpretation:

types:	objects of $T$
terms:	morphisms of $T$
equations between terms:	commuting diagrams
rewrites between terms:	“edges” in $\text{hom}$ in $V$

To be clear, this is not a new idea. Using enriched Lawvere theories for operational semantics has been explored in the past. For example, category-enriched theories have been studied by Seely [28] for the  $\lambda$ -calculus, and poset-enriched ones by Ghani and L  th [19] for understanding “modularity” in term rewriting systems. They have been utilized extensively by Power, enriching in  $\omega$ -complete partial orders to study recursion [25] – in fact, there the simplified “natural number” enriched theories which we explore were implicitly considered.

The goal of this paper is to give a simple unified explanation of enriched Lawvere theories and some of their applications to operational semantics. We aim our explanations at readers familiar with category theory but not yet enriched categories. To reduce the technical overhead we only consider enrichment over cartesian closed categories.

Because operational semantics uses (labelled) directed graphs to represent terms and rewrites, this should be our main example of enriching category; i.e. the “thing” of  $n$ -ary operations, or  $n$ -variable terms in a theory, should be a directed graph whose edges are rewrites. However, for reasons to be explained, we need to use a slightly different concept called a *simplicial set*. For all intents and purposes, it can be understood as an ordinary directed graph; although, it does provide a more general and powerful framework for higher-dimensional rewriting, which should prove to be quite useful.


We denote by  $\mathbf{sSet} := [\Delta^{\text{op}}, \mathbf{Set}]$  the category of simplicial sets, contravariant functors from the category of linear orders and monotone maps to the category of sets and functions, and natural transformations between them. They can be visualized as spaces built from “simplices”, which are triangles of any dimension. For an introduction to simplicial sets, see [32]. We will only need those simplicial sets which have “nondegenerate”, or nontrivial, elements in dimension 1. These we give the simpler name **unital graphs**, and we denote the full subcategory of unital graphs by  $\mathbf{uGph}$ . Hereafter, by “graph” we mean unital graph, though it will usually be explicit.

In general, for a cartesian closed category  $V$ , we take a “ $V$ -theory” to be a  $V$ -enriched Lawvere theory with natural number arities. We consider  $V$  as a choice of “method of computation” for algebraic theories. There are functors between enriching categories that allow the translation between different kinds of operational and denotational semantics. There is a spectrum of choices for  $V$ , which allow us to examine the computation of term calculi at various levels of detail:

- **Unital Graphs:**  $\mathbf{uGph}$ -theories represent “small-step” operational semantics:
  - a hom-edge is a *single* term rewrite.
- **Categories:**  $\mathbf{Cat}$ -theories represent “big-step” operational semantics.
  - (Often this means a rewrite to a *normal form*. We use the term more generally.)
  - a composite of morphisms is a *big-step* rewrite.
- **Posets:**  $\mathbf{Pos}$ -theories represent “full-step” operational semantics:
  - a hom-poset boolean is the *existence* of a big-step rewrite.
- **Sets:**  $\mathbf{Set}$ -theories represent denotational semantics:
  - a hom-set element is an *equivalence class* of the symmetric closure of the big-step relation.

In Section 2 we review Lawvere theories as a more explicit, but equivalent, presentation of finitary monads. In Section 3, we recall the basics of enrichment over cartesian closed categories. In Section 4 we give the central definition of  $\mathbf{V}$ -theory, adapted from the work of Lucyshyn-Wright [18], which allows us to apply his theorem relating enriched Lawvere theories and monads.

In Section 6 we discuss how suitable functors between enriching categories induce *change-of-computation*: they transform theories, and their models, from one method of rewriting to another. Our main examples arise from this chain of adjunctions:



The right adjoints here automatically preserve finite products, and the left adjoints do as well:

- $R: \mathbf{uGph} \rightarrow \mathbf{Cat}$ , called *realization*, maps a unital graph  $G$  to the category whose morphisms are sequences of edges in  $G$ , and composition is concatenation, modulo equations for associativity and unitality. (This is where the distinction between graphs and unital graphs matters. For an explanation of this functor, see appendix.) Change of base along  $R$  maps small-step operational semantics to big-step operational semantics.
- $FP: \mathbf{Cat} \rightarrow \mathbf{Pos}$  maps a category  $C$  to the poset whose elements are objects of  $C$ , with  $c \leq c'$  iff  $C$  has a morphism from  $c$  to  $c'$ . Change of base along  $FP$  maps big-step operational semantics to full-step operational semantics.
- $FS: \mathbf{Pos} \rightarrow \mathbf{Set}$  maps a poset  $P$  to the set of “components” of  $P$ , where  $p, p' \in P$  are in the same component if  $p \leq p'$ . Change of base along  $FS$  maps full-step operational semantics to denotational semantics.

In Section 7 we show that models of all  $\mathbf{V}$ -theories for all enriching  $\mathbf{V}$  can be assimilated into one category using the Grothendieck construction. In Section 8.1 we bring all the strands together and demonstrate these concepts with the *SKI*-combinator calculus. In Section 8.4 we show how theories enriched over the category of labelled graphs can be used to study bisimulation.

**Acknowledgements.** This paper builds upon the ideas of Mike Stay and Greg Meredith presented in “Representing operational semantics with enriched Lawvere theories” [30]. We appreciate their offer to let us develop this work further for use in the innovative distributed computing system RChain, and gratefully acknowledge the support of Pyroflex Corporation. We also thank Richard Garner, Todd Trimble and others at the *n*-Category Café for classifying cartesian closed categories where every object is a finite coproduct of copies of the terminal object [2].

## 2. LAWVERE THEORIES

Algebraic structures are traditionally treated as sets equipped with operations obeying equations, but we can generalize such structures to live in any category with finite products. For example, given any category  $\mathbf{C}$  with finite products, we can define a monoid internal to  $\mathbf{C}$  to consist of:

an object	$M$
an identity element	$e: 1 \rightarrow M$
and multiplication	$m: M^2 \rightarrow M$
obeying the associative law	$m \circ (m \times M) = m \circ (M \times m)$
and the right and left unit laws	$m \circ (e \times \text{id}_M) = \text{id}_M = m \circ (\text{id}_M \times e).$

Lawvere theories formalize this idea. For example, there is a Lawvere theory  $\text{Th}(\text{Mon})$ , the category with finite products freely generated by an object  $t$  equipped with an identity element  $e: 1 \rightarrow t$  and multiplication  $m: t^2 \rightarrow t$  obeying the associative law and unit laws listed above. This captures the “Platonic idea” of a monoid internal to a category with finite products. A monoid internal to  $\mathbf{C}$  then corresponds to a functor  $\mu: \mathbf{T} \rightarrow \mathbf{C}$  that preserves finite products.

In more detail, let  $\mathbf{N}$  be any skeleton of the category of finite sets  $\mathbf{FinSet}$ . Because  $\mathbf{N}$  is the free category with finite coproducts on 1,  $\mathbf{N}^{\text{op}}$  is the free category with finite products on 1. A **Lawvere theory** is a category with finite products  $\mathbf{T}$  equipped with a functor  $\tau: \mathbf{N}^{\text{op}} \rightarrow \mathbf{T}$  that is bijective on objects and preserves finite products. Thus, a Lawvere theory is essentially a category generated by one object  $\tau(1) = t$  and  $n$ -ary operations  $t^n \rightarrow t$ , as well as the projection and diagonal morphisms of finite products.

For efficiency let us call a functor that preserves finite products **cartesian**. Lawvere theories are the objects of a category  $\mathbf{Law}$  whose morphisms are cartesian functors  $f: \mathbf{T} \rightarrow \mathbf{T}'$  that obey  $f\tau = \tau'$ . More generally, for any category with finite products  $\mathbf{C}$ , a **model** of the Lawvere theory  $\mathbf{T}$  in  $\mathbf{C}$  is a cartesian functor  $\mu: \mathbf{T} \rightarrow \mathbf{C}$ . The models of  $\mathbf{T}$  in  $\mathbf{C}$  are the objects of a category  $\text{Mod}(\mathbf{T}, \mathbf{C})$ , in which the morphisms are natural transformations.

A theory can thus have models in many different contexts. For example, there is a Lawvere theory  $\text{Th}(\text{Mon})$ , the theory of monoids, described as above. Ordinary monoids are models of this theory in  $\mathbf{Set}$ , while topological monoids are models of this theory in  $\mathbf{Top}$ .

For completeness, it is worthwhile to mention the *presentation* of a Lawvere theory: after all, we are arguing their utility in everyday programming. How exactly does the above “sketch” of  $\text{Th}(\text{Mon})$  produce a category with finite products? It is precisely analogous to the presentation of an algebra by generators and relations: we form the free category with finite products on the data given, and impose the required equations. The result is a category whose objects are powers of  $M$ , and whose morphisms are composites of products of the morphisms in  $\text{Th}(\text{Mon})$ , projections, deletions, symmetries and diagonals. A detailed account was given by Barr and Wells [5, Chap. 4]; for a more computer-science-oriented approach see Crole [9, Chap. 3].

Currently, *monads* are more widely used in computer science than Lawvere theories. However, Hyland and Power have suggested that Lawvere theories could do much of the work that monads do today [12]. In 1965, Linton [17] proved that Lawvere theories correspond to “finitary monads” on the category of sets. For every Lawvere theory  $\mathbf{T}$ , there is an adjunction:

$$\begin{array}{ccc}
 & F & \\
 \text{Set} & \xrightarrow{\quad} & \text{Mod}(\mathbf{T}, \text{Set}). \\
 & U & \\
 & \perp &
 \end{array}$$

The functor

$$U: \text{Mod}(\mathbf{T}, \mathbf{Set}) \rightarrow \mathbf{Set}$$

sends each model  $\mu$  to its underlying set,  $X = \mu(\tau(1))$ . Its left adjoint, the free model functor

$$F: \mathbf{Set} \rightarrow \text{Mod}(\mathbf{T}, \mathbf{Set}),$$

sends each finite set  $n \in \mathbf{N}$  to the representable functor  $\mathbf{T}(\tau(n), -): \mathbf{T} \rightarrow \mathbf{Set}$ , and in general any set  $X$  to the colimit of all such representables as  $n$  ranges over the poset of finite subsets of  $X$ . In rough terms,  $F(X)$  is the model of all  $n$ -ary operations from  $\mathbf{T}$  on the set  $X$ .

If we momentarily abbreviate  $\text{Mod}(\mathbf{T}, \mathbf{Set})$  as  $\text{Mod}$ , we obtain an adjunction

$$\text{Mod}(F(n), \mu) = \text{Mod}(\mathbf{T}(\tau(n), -), \mu) \cong \mu(\tau(n)) \cong \mu(\tau(1))^n = \mathbf{Set}(n, U(\mu))$$

where the left isomorphism arises from the Yoneda lemma, and the right isomorphism from the product preservation of  $\mu$ .

This adjunction induces a monad  $T$  on  $\mathbf{Set}$ :

$$(1) \quad T(X) = \int^{n \in \mathbf{N}} X^n \times \mathbf{T}(n, 1).$$

The integral here is a coend, essentially a coproduct quotiented by the equations of the theory and the equations induced by the cartesian structure of the category. This forms the set of all terms that can be constructed from applying the operations to the elements, subject to the equations of the theory. The monad constructed this way is always **finitary**: that is, it preserves filtered colimits [1], or its action on sets is determined by its action on finite sets.

Conversely, for a monad  $T$  on  $\mathbf{Set}$ , its Kleisli category  $\text{Kl}(T)$  is the category of all free algebras of the monad, which has all coproducts. There is a functor  $k: \mathbf{Set} \rightarrow \text{Kl}(T)$  that is the identity on objects and preserves coproducts. Thus,

$$k^{\text{op}}: \mathbf{Set}^{\text{op}} \rightarrow \text{Kl}(T)^{\text{op}}$$

is a cartesian functor, and restricting its domain to  $\mathbf{N}^{\text{op}}$  is a Lawvere theory  $k_T$ . To see what this is doing, note that:

$$\text{Kl}(T)^{\text{op}}(n, m) = \text{Kl}(T)(m, n) = \mathbf{Set}(m, T(n))$$

where the latter is considered as  $m$   $n$ -ary operations in the Lawvere theory  $k_T$ . When  $T$  is finitary, the monad arising from this Lawvere theory is naturally isomorphic to  $T$  itself.

This correspondence sets up an equivalence between the category **Law** of Lawvere theories and the category of finitary monads on  $\mathbf{Set}$ . There is also an equivalence between the category  $\text{Mod}(\mathbf{T}, \mathbf{Set})$  of models of a Lawvere theory  $\mathbf{T}$  and the category of algebras of the corresponding finitary monad  $T$ . Furthermore, all this generalizes with  $\mathbf{Set}$  replaced by any “locally finitely presentable” category [1]. For more details see [5, 16, 20].

### 3. ENRICHMENT

To allow more general semantics, we now turn to Lawvere theories that have hom-*objects* rather than mere hom-*sets*. To do this we use enriched category theory [14] and replace sets with objects of a cartesian closed category  $\mathbf{V}$ , called the “enriching” category or “base”. A  $\mathbf{V}$ -enriched category

or **V-category**  $\mathbf{C}$  is:

$$\begin{array}{ll} \text{a collection of objects} & \text{Ob}(\mathbf{C}) \\ \text{a hom-object function} & \mathbf{C}(-, -): \text{Ob}(\mathbf{C}) \times \text{Ob}(\mathbf{C}) \rightarrow \text{Ob}(\mathbf{V}) \\ \text{composition morphisms} & \circ_{a,b,c}: \mathbf{C}(b, c) \times \mathbf{C}(a, b) \rightarrow \mathbf{C}(a, c) \quad \forall a, b, c \in \text{Ob}(\mathbf{C}) \\ \text{identity-assigning morphisms} & i_a: 1_{\mathbf{V}} \rightarrow \mathbf{C}(a, a) \quad \forall a \in \text{Ob}(\mathbf{C}) \end{array}$$

such that composition is associative and unital. A **V-functor**  $F: \mathbf{C} \rightarrow \mathbf{D}$  is:

$$\begin{array}{ll} \text{a function} & F: \text{Ob}(\mathbf{C}) \rightarrow \text{Ob}(\mathbf{D}) \\ \text{a collection of morphisms} & F_{ab}: \mathbf{C}(a, b) \rightarrow \mathbf{D}(F(a), F(b)) \quad \forall a, b \in \mathbf{C} \end{array}$$

such that  $F$  preserves composition and identity. A **V-natural transformation**  $\alpha: F \Rightarrow G$  is:

$$\text{a family } \alpha_a: 1_{\mathbf{V}} \rightarrow \mathbf{D}(F(a), G(a)) \quad \forall a \in \text{Ob}(\mathbf{C})$$

such that  $\alpha$  is “natural” in  $a$ : an evident square commutes. There is a 2-category  $\mathbf{VCat}$  of  $\mathbf{V}$ -categories,  $\mathbf{V}$ -functors, and  $\mathbf{V}$ -natural transformations.

We can construct new  $\mathbf{V}$ -categories from old by taking products and opposites in an obvious way. There is also a  $\mathbf{V}$ -category denoted  $\underline{\mathbf{V}}$  with the same objects as  $\mathbf{V}$  and with hom-objects given by the internal hom:

$$\underline{\mathbf{V}}(v, w) = w^v \quad \forall v, w \in \mathbf{V}.$$

The concepts of adjunction and monad generalize straightforwardly to  $\mathbf{V}$ -categories, and when we speak of an adjunction or monad in the enriched context this generalization is what we mean [14]. For example, there is an adjunction

$$\underline{\mathbf{V}}(u \times v, w) \cong \underline{\mathbf{V}}(u, w^v)$$

called “currying”.

We can generalize products and coproducts to the enriched context. Given a  $\mathbf{V}$ -category  $\mathbf{C}$ , the **V-coproduct** of an  $n$ -tuple of objects  $b_1, \dots, b_n \in \text{Ob}(\mathbf{C})$  is an object  $b$  equipped with a  $\mathbf{V}$ -natural isomorphism

$$\mathbf{C}(b, -) \cong \prod_{i=1}^n \mathbf{C}(b_i, -).$$

If such an object exists, we denote it by  $\sum_{i=1}^n b_i$ . This makes sense even when  $n = 0$ : a 0-ary  $\mathbf{V}$ -coproduct in  $\mathbf{C}$  is called a **V-initial object** and denoted as  $0_{\mathbf{C}}$ . When  $\mathbf{V}$  is cartesian closed, any finite coproduct that exists in  $\mathbf{V}$  is also a  $\mathbf{V}$ -coproduct in  $\underline{\mathbf{V}}$ . In particular,

$$u^{v+w} \cong u^v \times u^w \quad \text{and} \quad w^0 \cong 1_{\mathbf{V}}$$

whenever  $0$  is an initial object of  $\mathbf{V}$ . Conversely, any finite  $\mathbf{V}$ -coproduct that exists in  $\mathbf{V}$  is also a coproduct in the usual sense.

Similarly, a **V-product** of objects  $b_1, \dots, b_n \in \text{Ob}(\mathbf{C})$  is an object  $b$  equipped with a  $\mathbf{V}$ -natural isomorphism

$$(2) \quad \mathbf{C}(-, b) \cong \prod_{i=1}^n \mathbf{C}(-, b_i).$$

If such an object  $b$  exists, we denote it by  $\prod_{i=1}^n b_i$ . A 0-ary product in  $\mathbf{C}$  is called a **V-terminal object** and denoted as  $1_{\mathbf{C}}$ . Whenever  $\mathbf{V}$  is cartesian closed, the finite products in  $\mathbf{V}$  are also  $\mathbf{V}$ -products in  $\underline{\mathbf{V}}$ . In particular,

$$(u \times v)^w \cong u^w \times v^w \quad \text{and} \quad 1_{\mathbf{V}}^w \cong 1_{\mathbf{V}}$$

where our chosen terminal object  $1_V$  is also  $V$ -terminal. Conversely, any finite  $V$ -product in  $V$  is also a product in the usual sense.

A general  $V$ -category  $C$  does not exactly have projections from a  $V$ -product to its factors, since given two objects  $c, c' \in \text{Ob}(C)$  there is not, fundamentally, a *set* of morphisms from  $c$  to  $c'$ . Instead there is the hom-object  $C(c, c')$ , which is an object of  $V$ . However, any object  $v$  of  $V$  has a set of **elements**, namely morphisms  $f: 1_V \rightarrow v$ . Elements of  $C(c, c')$  act like morphisms from  $c$  to  $c'$ .

In particular, any  $V$ -product  $b = \prod_{i=1}^n b_i$  gives rise to elements

$$p_i: 1_V \rightarrow C(b, b_i)$$

which serve as substitutes for the projections in a usual product. These elements are defined as composites

$$1_V \xrightarrow{i_b} C(b, b) \xrightarrow{\sim} \prod_{i=1}^n C(b, b_i) \rightarrow C(b, b_i)$$

where the isomorphism comes from Eq. (2) and the last arrow is a projection in  $V$ .

Even better, we can bundle up all these elements  $p_i$  into a single element

$$p: 1_V \rightarrow \prod_{i=1}^n C(b, b_i)$$

which serves as a substitute for the universal cone in a usual product. Starting from  $p$  we can recover the  $V$ -natural isomorphism in Eq. (2) as follows:

$$(3) \quad C(-, b) \xrightarrow{\sim} 1_V \times C(-, b) \xrightarrow{p \times 1} \prod_{i=1}^n C(b, b_i) \times C(-, b) \longrightarrow \prod_{i=1}^n C(-, b_i)$$

where the last arrow is given by composition. Thus, we say a **universal cone** exhibiting  $b$  as the  $V$ -product of objects  $b_1, \dots, b_n$  is an element  $p: 1_V \rightarrow \prod_{i=1}^n C(b, b_i)$  such that the  $V$ -natural transformation  $C(-, b) \rightarrow \prod_{i=1}^n C(-, b_i)$  given by Eq. (3) is an isomorphism.

The advantage of this reformulation is that we can say a  $V$ -functor  $F: C \rightarrow D$  **preserves finite  $V$ -products** if for every universal cone  $p: 1_V \rightarrow \prod_{i=1}^n C(b, b_i)$  exhibiting  $b$  as the  $V$ -product of the objects  $b_i$ , the composite

$$1_V \xrightarrow{p} \prod_{i=1}^n C(b, b_i) \xrightarrow{\prod_i F} D(F(b), F(b_i))$$

is universal cone exhibiting  $F(b)$  as the  $V$ -product of the objects  $F(b_i)$ .

A bit more subtly, generalizing the exponentials in  $V$ , a  $V$ -category  $C$  can have “powers”. Given  $v \in \text{Ob}(V)$ , we say an object  $c^v \in \text{Ob}(C)$  is a  **$v$ -power** of  $c \in \text{Ob}(C)$  if it is equipped with a  $V$ -natural isomorphism

$$(4) \quad C(-, c^v) \cong C(-, c)^v.$$

In the special case  $V = \mathbf{Set}$  this forces  $c^v$  to be the  $v$ -fold product of copies of  $c$ . As with  $V$ -products, it is useful to repackage the isomorphism of Eq. (4) so we can say what it means for a  $V$ -functor to preserve  $v$ -powers. First, note that this isomorphism gives rise to an element

$$q: 1_V \rightarrow C(c^v, c)^v,$$

namely the composite

$$1_V \xrightarrow{i_{c^v}} C(c^v, c^v) \xrightarrow{\sim} C(c^v, c)^v.$$

Conversely, any element  $q: 1_V \rightarrow C(c^v, c)^v$  determines a  $V$ -natural transformation  $e: C(-, c^v) \rightarrow C(-, c)^v$ , and we say  $e$  is a **universal cone** if this  $V$ -natural transformation is an isomorphism. Next, suppose  $C$  and  $D$  are  $V$ -categories with  $v$ -powers. We say a  $V$ -functor  $F: C \rightarrow D$  **preserves  $v$ -powers** if it maps universal cones to universal cones.

There are just a few more technicalities. A category is **locally finitely presentable** if it is the category of models for a finite limits theory, and an object is **finite** if its representable functor is **finitary**: that is, it preserves filtered colimits [1]. A  $V$ -category  $C$  is **locally finitely presentable** if its underlying category  $C_0$  is locally finitely presentable,  $C$  has finite powers, and  $(-)^x: C_0 \rightarrow C_0$  is finitary for all finitely presentable  $x$ . The details are not crucial here: all categories to be considered are locally finitely presentable. We will use  $V_f$  to denote the full subcategory of  $V$  of finite objects: in  $\mathbf{uGph}$ , these are simply unital graphs with finitely many vertices and edges.

#### 4. ENRICHED LAWVERE THEORIES

Power introduced the notion of enriched Lawvere theory about twenty years ago, “in seeking a general account of what have been called notions of computation” [26]. The original definition is as follows: for a symmetric monoidal closed category  $(V, \otimes, 1)$ , a “ $V$ -enriched Lawvere theory” is a  $V$ -category  $T$  that has powers by objects in  $V_f$ , equipped with an identity-on-objects  $V$ -functor

$$\tau: \underline{V}_f^{\text{op}} \rightarrow T$$

that preserves these powers. A “model” of a  $V$ -theory is a  $V$ -functor  $\mu: T \rightarrow V$  that preserves powers by finite objects of  $V$ . There is a category  $\mathbf{Mod}(T, V)$  whose objects are models and whose morphisms are  $V$ -natural transformations. The monadic adjunction and equivalence of Section 2 generalize to the enriched setting.

However, this sort of  $V$ -enriched Lawvere theory has arities for every finite object of  $V$ . These *generalized arities* may be very powerful—rather than only inputting  $n$ -tuples of terms, we can input any finite object of terms. Despite its potential, this generalization remains largely unexploited in computer science. Power [15] introduced “enriched sketches” as a way of presenting enriched Lawvere theories, but to the authors’ knowledge these are not yet widely understood or used. What does it mean for an operation to take in a finite (unital) graph of terms? How can we learn to use this generality? One clue that we note in Example 7 is that limits and colimits are operations whose arity is a “diagram shape” rather than a natural number. We hope that this idea is explored more widely, so that we can use more general arities in both mathematics and computer programming.

In this paper, however, we only consider *natural number* arities, while still retaining enrichment. To do this we use the work of Lucyshyn-Wright [18], who along with Power [23] has generalized Power’s original ideas to allow a more flexible choice of arities. We also limit ourselves to the case where the tensor product of  $V$  is cartesian. This has a significant simplifying effect, yet it suffices for many cases of interest in computer science.

Thus, in all that follows, we let  $(V, \times, 1_V)$  be a cartesian closed category equipped with chosen finite coproducts of the terminal object  $1_V$ , say

$$n_V = \sum_{i \in n} 1_V.$$

Define  $N_V$  to be the full subcategory of  $V$  containing just these objects  $n_V$ . There is also a  $V$ -category  $\underline{N}_V$  whose objects are those of  $N_V$  and whose hom-objects are given as in  $V$ . We define the  $V$ -category of **arities** for  $V$  to be

$$A_V := \underline{N}_V^{\text{op}}.$$



We shall soon see that  $A_V$  has finite  $V$ -products.

**Definition 1.** We define a  **$V$ -theory**  $(T, \tau)$  to be a  $V$ -category  $T$  equipped with a  $V$ -functor

$$\tau: A_V \rightarrow T$$

that is bijective on objects and preserves finite  $V$ -products.

**Definition 2.** A **model** of  $T$  in a  $V$ -category  $C$  is a  $V$ -functor

$$\mu: T \rightarrow C$$

that preserves finite  $V$ -products.

Just as all the objects of a Lawvere theory are finite products of a single object, we shall see that all the objects of  $T$  are finite  $V$ -products of the object

$$t = \tau(1_V).$$

**Definition 3.** We define  $V\text{Law}$ , the **category of  $V$ -theories**, to be the category for which an object is a  $V$ -theory and a morphism from  $(T, \tau)$  to  $(T', \tau')$  is a  $V$ -functor  $f: T \rightarrow T'$  that preserves finite  $V$ -products and has  $f\tau = \tau'$ .

**Definition 4.** For every  $V$ -theory  $(T, \tau)$  and every  $V$ -category  $C$  with finite  $V$ -products, we define  $\text{Mod}(T, C)$ , the **category of models** of  $(T, \tau)$  in  $C$ , to be the category for which an object is a  $V$ -functor  $\mu: T \rightarrow C$  that preserves finite  $V$ -products and a morphism is a  $V$ -natural transformation.

The basic monadicity results for Lawvere theories generalize to  $V$ -theories when  $V$  is complete and cocomplete, as in the main examples we consider:  $V = \mathbf{Gph}, \mathbf{Cat}, \mathbf{Pos}$ , and  $\mathbf{Set}$ . Under this extra assumption  $V\text{Law}$  and  $\text{Mod}(T, C)$  can be promoted to  $V$ -categories, which we call  $\underline{V\text{Law}}$  and  $\underline{\text{Mod}}(T, C)$ . Furthermore, there is a  $V$ -functor

$$U: \underline{\text{Mod}}(T, V) \rightarrow \underline{V}$$

sending any model  $\mu: T \rightarrow V$  to its underlying object  $\mu(t) \in V$ . Recall that monads and adjunctions make sense in  $V\text{Cat}$ , just as they do in  $\mathbf{Cat}$ . The  $V$ -functor  $U$  has a left adjoint

$$F: \underline{V} \rightarrow \underline{\text{Mod}}(T, V),$$

and  $\underline{\text{Mod}}(T, V)$  is equivalent to the  $V$ -category of algebras of the resulting monad  $T = UF$ . More precisely:

**Theorem 5.** Suppose  $V$  is cartesian closed, complete and cocomplete, and has chosen finite co-products of the terminal object. Let  $(T, \tau)$  be a  $V$ -theory. Then there is a monadic adjunction

$$\begin{array}{ccc} & F & \\ \underline{V} & \xrightarrow{\quad} & \underline{\text{Mod}}(T, V) \\ & U & \end{array} \quad \perp$$

*Proof.* This follows from Lucyshyn-Wright's general theory [18], so our task is simply to explain how. He allows  $V$  to be a symmetric monoidal category, and uses a more general concept of algebraic theory with a system of arities given by any fully faithful symmetric monoidal  $V$ -functor  $j: J \rightarrow V$ . For us  $J = \underline{N}_V$  and  $j: \underline{N}_V \rightarrow V$  is the obvious inclusion; this is his Example 3.7.

Lucyshyn-Wright defines a  **$J$ -theory** to be a  $V$ -functor  $\tau: J^{\text{op}} \rightarrow T$  that is the identity on objects and preserves powers by objects in  $J$  (or more precisely, their images under  $j$ ). For us  $J^{\text{op}} = A_V$ .

We are only demanding that  $\tau: \mathbf{A}_V \rightarrow \mathbf{T}$  be bijective on objects, but we can make it the identity on objects simply by renaming the objects of  $\mathbf{T}$ . So, to apply his theory, we need to show that a  $V$ -functor  $\tau: \mathbf{A}_V \rightarrow \mathbf{T}$  preserves powers by objects in  $\mathbf{N}_V$  if and only if it preserves finite  $V$ -products. This is Lemma 12 below.

He defines a model (or “algebra”) of a  $J$ -theory to be a  $V$ -functor  $\tau: \mathbf{T} \rightarrow \underline{V}$  that preserves powers by objects in  $J$ . He defines a morphism of models to be a  $V$ -natural transformation between such  $V$ -functors. So, to apply his theory, we also need to show that when  $J = \mathbf{N}_V$ , a  $V$ -functor  $\mu: \mathbf{T} \rightarrow \underline{V}$  preserves powers by objects of  $J$  if and only if it preserves finite  $V$ -products. This is Lemma 13 below.

A technical concept fundamental to Lucyshyn-Wright’s theory is that of an **eleutheric** system of arities  $j: J \rightarrow \underline{V}$ . This is one where the left Kan extension of any  $V$ -functor  $f: J \rightarrow \underline{V}$  along  $j$  exists and is preserved by each  $V$ -functor  $\underline{V}(x, -): \underline{V} \rightarrow \underline{V}$ . In Example 7.5.5 he shows that  $j: \mathbf{N}_V \rightarrow \underline{V}$  is eleutheric when  $V$  is countably cocomplete. In Thm. 8.9 shows that when  $j: J \rightarrow \underline{V}$  is eleutheric, and has equalizers, we may form the  $V$ -category  $\mathbf{Mod}(\mathbf{T}, V)$ , and that the forgetful  $V$ -functor

$$U: \mathbf{Mod}(\mathbf{T}, V) \rightarrow \underline{V}$$

is monadic. This is the result we need. So, our theorem actually holds whenever  $V$  is cartesian closed, with equalizers and countable colimits, and has chosen finite coproducts of the initial object.  $\square$

Before turning to examples, a word about Lucyshyn-Wright’s construction of the left adjoint  $F$  and the monad  $T$  is in order. These rely on the “free model” on an object  $n_V \in V$ . This is the enriched generalization of the free model described in Section 2: it is the composite of  $\tau^{\text{op}}: \mathbf{A}_V^{\text{op}} \rightarrow \mathbf{T}^{\text{op}}$  with the enriched Yoneda embedding  $y: \mathbf{T}^{\text{op}} \rightarrow [\mathbf{T}, V]$ :

$$\begin{aligned} \mathbf{A}_V^{\text{op}} &\xrightarrow{\tau^{\text{op}}} \mathbf{T}^{\text{op}} \xrightarrow{y} [\mathbf{T}, V] \\ n_V &\mapsto t^{n_V} \mapsto \mathbf{T}(t^{n_V}, -) \end{aligned}$$

Since an object of  $V$  does not necessarily have a “poset of finite subobjects” over which to take a filtered colimit (as in  $\mathbf{Set}$ ), the extension of this “free model” functor  $y\tau^{\text{op}}$  to all of  $V$  is specified by a somewhat higher-powered generalization: it is the left Kan extension of  $y\tau^{\text{op}}$  along  $j$ .

$$\begin{array}{ccc} \mathbf{N}_V & \xrightarrow{y\tau^{\text{op}}} & [\mathbf{T}, V] \\ & \searrow j & \downarrow \eta \\ & & V \end{array} \quad \begin{array}{c} \nearrow \\ F := \text{Lan}_j y\tau^{\text{op}} \end{array}$$

This is the universal “best solution” to the problem of making the triangle commute up to a  $V$ -natural transformation. That is, for any functor  $G: V \rightarrow [\mathbf{T}, V]$  and  $V$ -natural transformation  $\theta: y\tau^{\text{op}} \Rightarrow Gj$ , the latter factors uniquely through  $\eta$ . From the adjunction between  $V$  and the category of models  $\mathbf{Mod}(\mathbf{T}, V)$  we obtain a  $V$ -enriched monad


$$T = UF: V \rightarrow V,$$

and this has a more concrete formula as an enriched coend:

$$T(V) = \int^{n_V \in \mathbf{N}_V} V^{n_V} \times \mathbf{T}(t^{n_V}, t).$$

We next give two examples of a rather abstract nature, where we show how  $\mathbf{Cat}$ -enriched Lawvere theories can describe categories with extra structure. In Section 8 we study examples more directly connected to operational semantics.

**Example 6.** When  $\mathbf{V} = \mathbf{Cat}$ , a  $\mathbf{V}$ -category is a 2-category, so a  $\mathbf{V}$ -theory deserves to be called a **2-theory**. For example, let  $\mathbf{T} = \mathbf{Th}(\mathbf{PsMon})$  be the 2-theory of pseudomonoids [10]. A pseudomonoid is a weakened version of a monoid: rather than associativity and unitality *equations*, it has 2-isomorphisms called the associator and unitors, which we can treat as *rewrite rules*. To equate various possible rewrite sequences, these 2-isomorphisms must obey equations called “coherence laws”. Here is a presentation of the 2-theory for pseudomonoids:



We write the equations as commutative diagrams merely for convenience; they could also be written as equations in a more traditional style. The top diagram expresses the pentagon identity for the associator, while the bottom one expresses the usual coherence law involving the left and right unitors.

Models of  $\mathbf{T} = \mathbf{Th}(\mathbf{PsMon})$  in  $\mathbf{Cat}$  are monoidal categories: let us explore this example in more detail. A model of  $\mathbf{T}$  is a finite-product-preserving 2-functor  $\mu: \mathbf{T} \rightarrow \mathbf{Cat}$ , which sends

$$\begin{aligned}
 t &\mapsto \mathbf{C} \\
 m &\mapsto \otimes: \mathbf{C}^2 \rightarrow \mathbf{C} \\
 e &\mapsto I: 1 \rightarrow \mathbf{C} \\
 \alpha &\mapsto a: \otimes \circ (\otimes \times 1_{\mathbf{C}}) \Rightarrow \otimes \circ (1_{\mathbf{C}} \times \otimes) \\
 \lambda &\mapsto \ell: I \circ 1_{\mathbf{C}} \Rightarrow 1_{\mathbf{C}} \\
 \rho &\mapsto r: 1_{\mathbf{C}} \circ I \Rightarrow 1_{\mathbf{C}}
 \end{aligned}$$

such that the coherence laws of the rewrites are preserved. Thus, a model is a category equipped with a tensor product  $\otimes$  and unit object  $I$  such that these operations are associative and unital up to natural isomorphism; so these models are precisely monoidal categories.

Given two models  $\mu, \nu: \mathbf{T} \rightarrow \mathbf{Cat}$ , a morphism of models is a 2-natural transformation  $\varphi: \mu \Rightarrow \nu$ ; this amounts to a strict monoidal functor  $\varphi: (\mathbf{C}, \otimes_{\mathbf{C}}, I_{\mathbf{C}}) \rightarrow (\mathbf{D}, \otimes_{\mathbf{D}}, I_{\mathbf{D}})$ . The strictness arises because morphisms between models are 2-natural transformations rather than pseudonatural transformations. There is a substantial amount of theory on pseudomonads and pseudoalgebras [7, 11], but to the authors' knowledge the theory-monad correspondence has not yet been extended to weak enrichment.

Finally, because  $\mathbf{Cat}$  is complete and cocomplete, the category of models  $\mathbf{Mod}(\mathbf{T}, \mathbf{Cat})$  can be promoted to a 2-category  $\mathbf{Mod}(\mathbf{T}, \mathbf{Cat})$ . This is the 2-category of monoidal categories, strict monoidal functors, and monoidal natural transformations.

We can accomplish the same thing on the monad side: a  $\mathbf{Cat}$ -enriched monad is called a **2-monad**, and  $\mathbf{T}$  gives rise to the “free monoidal category” 2-monad  $T$  on  $\mathbf{Cat}$  [7]. To apply this 2-monad to  $C \in \mathbf{Cat}$  we first form the free model on  $C$  by taking a left Kan extension as above, and then evaluate this model at the generating object. In the same way that the (underlying set of the) free monoid on a set  $X$  consists of all finite strings of elements of  $X$ ,  $T(C)$  is the monoidal category consisting of all finite tensor products of objects of  $C$  and all morphisms built from those of  $C$  by composition and tensoring together with associators and unitors obeying the necessary coherence laws. Morphisms of these algebras are strict monoidal functors, while 2-morphisms are natural transformations. We thus have a 2-equivalence between  $\mathbf{Mod}(\mathbf{T}, \mathbf{Cat})$  and the 2-category of algebras of  $T$ .

In this way, 2-theories generalize equipping *set*-like objects with operations obeying equations to equipping *category*-like objects with operations obeying equations up to transformations that obey equations of their own. In particular, this gives us a way to present graphical calculi such as string diagrams – the language of monoidal categories.

**Example 7.** Enrichment generalizes operations in more ways than by weakening equations to coherent isomorphisms. We can also use 2-theories to describe other structures that make sense inside 2-categories, such as adjunctions.

For example, we may define a cartesian category  $\mathbf{X}$  to be one equipped with right adjoints to the diagonal  $\Delta_{\mathbf{X}}: \mathbf{X} \rightarrow \mathbf{X} \times \mathbf{X}$  and the unique functor  $!_{\mathbf{X}}: \mathbf{X} \rightarrow 1_{\mathbf{Cat}}$ . These right adjoints are a functor  $m: \mathbf{X}^2 \rightarrow \mathbf{X}$  describing binary products in  $\mathbf{X}$  and a functor  $e: 1 \rightarrow \mathbf{X}$  picking out the terminal object in  $\mathbf{X}$ . We can capture the fact that they are right adjoints by providing them with units and counits and imposing the triangle equations. There is thus a 2-theory  $\mathbf{Th}(\mathbf{Cart})$  whose models in  $\mathbf{Cat}$  are categories with chosen finite products. More generally a model of this 2-theory in any 2-category  $\mathbf{C}$  with finite products is called a **cartesian object** in  $\mathbf{C}$ .

$\mathbf{Th}(\mathbf{Cart})$		
<b>type</b>	$X$	cartesian object
<b>operations</b>	$m: X^2 \rightarrow X$	product
	$e: 1 \rightarrow X$	terminal element
<b>rewrites</b>	$\Delta: \mathrm{id}_X \Rightarrow m \circ \Delta_X$	unit of adjunction between $m$ and $\Delta_X$
	$\pi: \Delta_X \circ m \Rightarrow \mathrm{id}_{X^2}$	counit of adjunction between $m$ and $\Delta_X$
	$\top: \mathrm{id}_X \Rightarrow e \circ !_X$	unit of adjunction between $e$ and $!_X$
	$\epsilon: !_X \circ e \Rightarrow \mathrm{id}_1$	counit of adjunction between $e$ and $!_X$
<b>equations</b>		

$$\begin{array}{ccc}
\begin{array}{c} \Delta_X \\ \Delta_X \circ \Delta \Downarrow \\ \Delta_X \circ m \circ \Delta_X \xrightarrow{\pi \circ \Delta_X} \Delta_X \end{array} & \begin{array}{c} \searrow 1 \\ \xrightarrow{\quad} \end{array} & \Delta_X \\
\begin{array}{c} !_X \\ !_X \circ \top \Downarrow \\ !_X \circ e \circ !_X \xrightarrow{e \circ !_X} !_X \end{array} & \begin{array}{c} \searrow 1 \\ \xrightarrow{\quad} \end{array} & !_X
\end{array}
\qquad
\begin{array}{ccc}
\begin{array}{c} m \\ \Delta \circ m \Downarrow \\ m \circ \Delta_X \circ m \xrightarrow{m \circ \pi} m \end{array} & \begin{array}{c} \searrow 1 \\ \xrightarrow{\quad} \end{array} & m \\
\begin{array}{c} e \\ \top \circ e \Downarrow \\ e \circ !_X \circ e \xrightarrow{e \circ \epsilon} e \end{array} & \begin{array}{c} \searrow 1 \\ \xrightarrow{\quad} \end{array} & e
\end{array}$$

Again we write the equations as commutative diagrams, but this time commutative triangles of 2-morphisms in  $\mathbf{Th}(\mathbf{Cart})$ . These are the triangle equations that force  $m$  to be the right adjoint of  $\Delta_X$  and  $e$  to be the right adjoint of  $!_X$ . A model of  $\mathbf{Th}(\mathbf{Cart})$  is a category with chosen binary products and a chosen terminal object; morphisms in  $\mathbf{Mod}(\mathbf{Th}(\mathbf{Cart}), \mathbf{Cat})$  are functors that strictly preserve this extra structure.

The subtle interplay between the cartesian structure of  $\mathbf{Th}(\mathbf{Cart})$  and the cartesian structure of the object  $X \in \mathbf{Th}(\mathbf{Cart})$  is an example of the “microcosm principle”: objects with a given structure are most generally defined in a context that has the same sort of structure. As seen in the previous example, we can also define pseudomonoids in any 2-category with finite products, but this is excess to requirements: one can in fact define them more generally in any monoidal 2-category [10].

In fact, if we let arities be finite categories, we would have  $\mathbf{Cat}$ -theories of categories with finite limits and colimits. However, for the purposes of this paper we are using only natural number arities. This suffices for constructing  $\mathbf{Th}(\mathbf{Cart})$  and also  $\mathbf{Th}(\mathbf{CoCart})$ , the theory of categories with chosen binary coproducts and a chosen initial object. Various other kinds of categories—distributive categories, rig categories, etc.—can also be expressed using  $\mathbf{Cat}$ -theories with natural number arities. This gives a systematic formalization of these categories, internalizes them to new contexts, and allows for the generation of 2-monads that describe them.

## 5. NATURAL NUMBER ARITIES

In this section we prove the lemmas required for Theorem 5 and our study of base change in Section 6. Throughout this section  $\mathbf{V}$  is cartesian closed with chosen  $n$ -fold coproducts  $n_V$  of its terminal object.

We begin with a study of  $\mathbf{N}_V$ , the full subcategory of  $V$  on the objects  $n_V$ . First we must resolve a potential ambiguity. On the one hand, for any object  $b$  of  $V$  we can form the exponential  $b^{n_V}$ . On the other hand, we can take the product of  $n$  copies of  $b$ , which we call  $b^n$ . Luckily these are the same, or at least naturally isomorphic:

**Lemma 8.** The functors  $(-)^{n_V} : V \rightarrow V$  and  $(-)^n : V \rightarrow V$  are naturally isomorphic.

*Proof.* If  $a, b \in V$ , then

$$\begin{aligned}
 V(a, b^{n_V}) &\cong V(a \times n_V, b) && \text{hom-tensor adjunction} \\
 &= V(a \times (n \cdot 1_V), b) && \text{definition of } n_V \\
 &\cong V(n \cdot (a \times 1_V), b) && \text{products distribute over coproducts} \\
 &\cong V(n \cdot a, b) && \text{unitality} \\
 &\cong V(a, b)^n && \text{definition of coproduct} \\
 &\cong V(a, b^n) && \text{definition of product.}
 \end{aligned}$$

Each of these isomorphisms is natural in  $a$  and  $b$ , so by the Yoneda lemma  $(-)^{n_V} \cong (-)^n$ .  $\square$

We can now understand coproducts, products and exponentials in  $\mathbf{N}_V$ :

**Lemma 9.** If  $V$  is any cartesian closed category with chosen coproducts of the initial object then  $\mathbf{N}_V$  is cartesian closed, with finite coproducts. The unique initial object of  $\mathbf{N}_V$  is  $0_V$ . The binary coproducts in  $\mathbf{N}_V$  are unique, given by

$$m_V + n_V = (m + n)_V.$$

The unique terminal object of  $\mathbf{N}_V$  is  $1_V$ , and the binary products are unique, given by

$$m_V \times n_V = (mn)_V.$$

Exponentials in  $\mathbf{N}_V$  are also unique, given by

$$m_V^{n_V} = (m^n)_V.$$

*Proof.* In  $V$  we know that  $0_V$  is an initial object and  $1_V$  is a terminal object, by definition. Since the subcategory  $\mathbf{N}_V$  is skeletal  $0_V$  is the unique initial object and  $1_V$  is the unique terminal object in  $\mathbf{N}_V$ . Similarly, in  $V$  we have defined  $(m + n)_V$  to be a coproduct of  $m_V$  and  $n_V$ , so in  $\mathbf{N}_V$  it is the unique such, and we can unambiguously write

$$m_V + n_V = (m + n)_V.$$

Products distribute over coproducts in any cartesian closed category, so in  $V$  we have

$$m_V \times n_V \cong (1_V + \cdots + 1_V) \times (1_V + \cdots + 1_V) \cong (mn)_V$$

where in the second step we use the distributive law twice. It follows that  $\mathbf{N}_V$  has finite products, and since this subcategory is skeletal they are unique, given by

$$m_V \times n_V = (mn)_V.$$

Finally, by Lemma 8 we have

$$m_V^{n_V} \cong m_V^n \cong \prod_{i=1}^n m_V \cong (m^n)_V.$$

It follows that  $\mathbf{N}_V$  has exponentials, and since this subcategory is skeletal they are unique, given by

$$m_V^{n_V} = (m^n)_V. \quad \square$$

We warn the reader that  $\text{hom}(m_V, n_V)$  may not have  $n^m$  elements. It does in **Gph**, **Cat**, **Pos** and of course **Set**, but not in  $V = \text{Set}^k$ , where  $|\text{hom}(m_V, n_V)| = n^{km}$ . In fact, whenever  $N_V$  has finite hom-sets it is equivalent to  $\text{FinSet}^k$  for some  $k$ . The reason is that  $2_V$  is an internal Boolean algebra in  $V$ , so its set of elements  $\text{hom}(1_V, 2_V)$  must be some Boolean algebra  $B$  in **Set**. A further argument due to Garner and Trimble shows that  $N_V$  is completely characterized, up to equivalence, by this Boolean algebra, and any Boolean algebra can occur [2]. If this Boolean algebra is finite it must be isomorphic to  $\{0, 1\}^k$  for some  $k \geq 0$ . In this case,  $N_V$  is equivalent to  $\text{FinSet}^k$ .

Now suppose  $C$  is a  $V$ -category. The question arises whether the power of an object  $c \in C$  by  $n_V$  must also be the  $V$ -product of  $n$  copies of  $c$ . The answer is yes:

**Lemma 10.** Let  $C$  be a  $V$ -category and  $c \in \text{Ob}(C)$ . Then the power  $c^{n_V}$  exists if and only if the  $n$ -fold  $V$ -product  $c^n$  exists, in which case they are isomorphic.

*Proof.* In Section 3 we saw that an object  $b \in \text{Ob}(C)$  is an  $n$ -fold  $V$ -product of copies of  $c$  precisely when it is equipped with a universal cone

$$p: 1_V \rightarrow C(b, c)^n.$$

Similarly,  $b$  is an  $n_V$ -power of  $c$  when it is equipped with a universal cone

$$q: 1_V \rightarrow C(b, c)^{n_V}.$$

The universality properties have the same form, and by Lemma 8 the functors  $(-)^n: V \rightarrow V$  and  $(-)^{n_V}: V \rightarrow V$  are naturally isomorphic. Thus, given either sort of universal cone we get the other, so an object is an  $n$ -fold product of copies of  $c$  if and only if it is the  $n_V$ -power of  $c$ .  $\square$

**Lemma 11.** Suppose  $C$  is a  $V$ -category such that every object is the  $n$ -fold  $V$ -product  $c^n$  of some object  $c$ . Then a  $V$ -functor  $F: C \rightarrow D$  preserves finite  $V$ -products if and only if it preserves powers by all objects of  $N_V$ .

*Proof.* Define a “finite  $V$ -power” to be a finite  $V$ -product of  $n$  copies of the same object. The  $V$ -functor  $F$  preserves finite  $V$ -powers if and only if it maps any universal cone

$$p: 1_V \rightarrow C(b, c)^n$$

in  $C$  to a universal cone in  $D$ . Similarly,  $F$  preserves powers by all objects of  $N_V$  if and only if it maps any universal cone

$$q: 1_V \rightarrow C(b, c)^{n_V}$$

in  $C$  to a universal cone in  $D$ . Two kinds of universality are involved here, but since they have the same form, and since Lemma 8 says the functors  $(-)^n: V \rightarrow V$  and  $(-)^{n_V}: V \rightarrow V$  are naturally isomorphic, it follows that  $F$  preserves finite  $V$ -powers if and only if it preserves powers by all objects of  $N_V$ .

It thus suffices to show that  $F$  preserves finite  $V$ -products if and only if it preserves finite  $V$ -powers. This follows from the assumption that every object is the  $n$ -fold  $V$ -product  $c^n$  of some object  $c$ .  $\square$

**Lemma 12.** Let  $V$  be cartesian closed with chosen finite coproducts of the terminal object and let  $T$  be a  $V$ -category. These conditions for a  $V$ -functor  $\tau: A_V \rightarrow T$  are equivalent:

- (1)  $(T, \tau)$  is a  $V$ -theory,
- (2)  $\tau$  preserves finite  $V$ -products,
- (3)  $\tau$  preserves powers by objects of  $N_V$ .

*Proof.* Conditions 1 and 2 are equivalent by definition. Since  $A_V = \underline{N}_V^{\text{op}}$ , finite  $V$ -products in  $A_V$  are the same as finite  $V$ -coproducts in  $\underline{N}_V$ , which are the same as finite coproducts in  $N_V$ . Since every object in  $\underline{N}_V$  is a finite coproduct of copies of  $1_V$ , Lemma 11 implies that conditions 2 and 3 are equivalent.  $\square$

**Lemma 13.** Given a  $V$ -theory  $(T, \tau)$  and a  $V$ -functor  $\mu: T \rightarrow C$ , the following conditions are equivalent:

- $\mu$  is a model of  $(T, \tau)$ ,
- $\mu$  preserves finite  $V$ -products,
- $\mu$  preserves powers by objects of  $N_V$ .

*Proof.* Conditions 1 and 2 are equivalent by definition. Since  $\tau$  is bijective on objects and preserves  $V$ -products each object of  $T$  is of the form  $t^n$  where  $t = \tau(1_V)$ . Thus, Lemma 11 implies that conditions 2 and 3 are equivalent.  $\square$

## 6. CHANGE OF BASE

We now have the tools to formulate the main idea: a choice of enrichment for Lawvere theories corresponds to a choice of *computation*, and changing enrichments corresponds to a *change of computation*. We propose a general framework in which one can translate between different forms of computation: small-step, big-step, full-step operational semantics, and denotational semantics.

Suppose that  $V$  and  $W$  are enriching categories of the sort we are considering: cartesian closed categories equipped with chosen finite coproducts of the terminal object. Suppose  $F: V \rightarrow W$  preserves finite products. This induces a **change of base** functor  $F_*: \mathbf{VCat} \rightarrow \mathbf{WCat}$  [8] which takes any  $V$ -category  $C$  and produces a  $W$ -category  $F_*(C)$  with the same objects but with

$$F_*(C)(a, b) := F(C(a, b))$$

for all objects  $a, b$ . Composition in  $F_*(C)$  is defined by

$$F(C(b, c)) \times F(C(a, b)) \xrightarrow{\sim} F(C(b, c) \times C(a, b)) \xrightarrow{F(\circ_{a,b,c})} F(C(a, b)).$$

The identity-assigning morphisms are given by

$$1_W \xrightarrow{\sim} F(1_V) \xrightarrow{F(i_a)} F(C(a, a)).$$

Moreover, if  $f: C \rightarrow D \in \mathbf{VCat}$  is a  $V$ -functor, there is a  $W$ -functor  $F_*(f): F_*(C) \rightarrow F_*(D)$  that on objects equals  $f$  and on hom-objects equals  $F(f)$ . If  $\alpha: f \Rightarrow g$  is a  $V$ -natural transformation and  $c \in \text{Ob}(C)$ , then we define

$$F_*(\alpha)_c: 1_W \xrightarrow{\sim} F(1_V) \xrightarrow{F(\alpha_c)} F(D(f(c), g(c))).$$

Thus, change of base actually gives a 2-functor from the 2-category of  $V$ -categories,  $V$ -functors and  $V$ -natural transformations to the corresponding 2-category for  $W$ .

In fact, the change of base operation gives a 2-functor

$$\begin{array}{ccc} \mathbf{MonCat} & \xrightarrow{(-)_*} & \mathbf{2Cat} \\ (F: V \rightarrow W) & \mapsto & (F_*: \mathbf{VCat} \rightarrow \mathbf{WCat}) \end{array}$$



In particular, if  $\mathbf{V}$  has not just finite coproducts of the terminal object, but all coproducts of this object, there is a map of adjunctions

$$\begin{array}{ccccc} \mathbf{Set} & \begin{array}{c} \xrightarrow{-\cdot 1} \\ \perp \\ \xleftarrow{\mathbf{V}(1, -)} \end{array} & \mathbf{V} & \xrightarrow{\quad} & \mathbf{Cat} & \begin{array}{c} \xrightarrow{(-\cdot 1)_*} \\ \perp \\ \xleftarrow{(\mathbf{V}(1, -))_*} \end{array} & \mathbf{VCat}. \end{array}$$

Each set  $X$  is mapped to the  $X$ -indexed coproduct of the terminal object in  $\mathbf{V}$  and conversely each object  $v$  of  $\mathbf{V}$  is represented in  $\mathbf{Set}$  by the hom-set from the unit to  $v$ . The latter induces the “underlying category” change of base, which forgets the enrichment. The former induces the “free  $\mathbf{V}$ -enrichment” change of base, whereby ordinary  $\mathbf{Set}$ -categories are converted to  $\mathbf{V}$ -categories, denoted  $\mathbf{C} \mapsto \underline{\mathbf{C}}$ . These form an adjunction, because 2-functors preserve adjunctions.

We now study how change of base affects theories and their models. We start by asking when a functor  $F: \mathbf{V} \rightarrow \mathbf{W}$  induces a change of base  $F_*: \mathbf{VCat} \rightarrow \mathbf{WCat}$  that “preserves enriched theories”. That is, given a  $\mathbf{V}$ -theory

$$\tau: \mathbf{A}_{\mathbf{V}} \rightarrow \mathbf{T}$$

we want to determine conditions for the base-changed functor

$$F_*(\tau): F_*(\mathbf{A}_{\mathbf{V}}) \rightarrow F_*(\mathbf{T})$$

to induce a  $\mathbf{W}$ -theory in a canonical way. Recall that we require  $\mathbf{V}$  and  $\mathbf{W}$  to be cartesian closed, equipped with chosen finite coproducts of their terminal objects. We thus expect the following conditions to be sufficient:  $F$  should be cartesian, and it should preserve the chosen finite coproducts of the terminal object:

$$F(n_{\mathbf{V}}) = n_{\mathbf{W}}$$

for all  $n$ .

Given these conditions there is a  $\mathbf{W}$ -functor, in fact an isomorphism

$$\tilde{F}: \mathbf{A}_{\mathbf{W}} \rightarrow F_*(\mathbf{A}_{\mathbf{V}}).$$

On objects this maps  $n_{\mathbf{W}}$  to  $n_{\mathbf{V}}$ , and on hom-objects it is simply the identity from

$$\mathbf{A}_{\mathbf{W}}(m_{\mathbf{W}}, n_{\mathbf{W}}) = n_{\mathbf{W}}^{m_{\mathbf{W}}} = (n^m)_{\mathbf{W}}$$

to

$$F(\mathbf{A}_{\mathbf{V}}(m_{\mathbf{V}}, n_{\mathbf{V}})) = F(n_{\mathbf{V}}^{m_{\mathbf{V}}}) = F((n^m)_{\mathbf{V}}) = (n^m)_{\mathbf{W}}$$

where we use Lemma 9 in these computations.

Using this we obtain a composite  $\mathbf{W}$ -functor

$$\mathbf{A}_{\mathbf{W}} \xrightarrow{\tilde{F}} F_*(\mathbf{A}_{\mathbf{V}}) \xrightarrow{F_*(\tau_{\mathbf{V}})} F_*(\mathbf{T}).$$

This is a bijection on objects and preserves finite  $\mathbf{V}$ -products because each of the factors has these properties. It is thus a  $\mathbf{W}$ -theory.

**Theorem 14.** Let  $\mathbf{V}, \mathbf{W}$  be cartesian closed categories with chosen finite coproducts of their terminal objects, and let  $F: \mathbf{V} \rightarrow \mathbf{W}$  be a cartesian functor that preserves these chosen coproducts. Then  $F_*$  **preserves enriched theories**: that is, for every  $\mathbf{V}$ -theory  $\tau_{\mathbf{V}}: \mathbf{A}_{\mathbf{V}} \rightarrow \mathbf{T}$ , the  $\mathbf{W}$ -functor

$$\tau_{\mathbf{W}} := F_*(\tau_{\mathbf{V}}) \circ \tilde{F}: \mathbf{A}_{\mathbf{W}} \rightarrow F_*(\mathbf{T})$$

is a  $\mathbf{W}$ -theory. Moreover,  $F_*$  **preserves models**: for every model  $\mu: \mathbf{T} \rightarrow \mathbf{C}$  of  $(\mathbf{T}, \tau_{\mathbf{V}})$ , the  $\mathbf{W}$ -functor  $F_*(\mu): F_*(\mathbf{T}) \rightarrow F_*(\mathbf{C})$  is a model of  $(F_*(\mathbf{T}), \tau_{\mathbf{W}})$ .

*Proof.* We have shown the first part. For the second, by Lemma 13 it suffices to assume that  $\mu$  preserves finite  $\mathbf{N}_V$ -powers and check that  $F_*(\mu)$  preserves  $\mathbf{N}_W$ -powers. We leave this as an exercise to the reader.  $\square$

Hence, any cartesian functor that preserves chosen finite coproducts of the terminal object gives a “change of computation”. That is, it provides for a method of translating formal languages between various “modes of operation”. Moreover, this reasoning generalizes to **multisorted**  $V$ -theories, enriched theories which have multiple sorts: given any  $n \in \mathbb{N}$ , the monoidal subcategory  $(\mathbf{N}_V)^n$  is also an eleutheric system of arities, so Lucyshyn-Wright’s monadicity theorem still applies. In Section 8.4 we show how this is useful in the study of bisimulation.

Next we introduce two more useful kinds of translations, and demonstrate how all of these can be encapsulated in a single category. The following section can be skipped by readers eager to see applications.

## 7. THE CATEGORY OF ALL MODELS

In addition to change of base, there are two other natural and useful ways to go between models of enriched theories. Suppose  $V$  is any cartesian closed category with chosen finite coproducts of the terminal object. Let  $V\text{Mod}(\mathbf{T}, \mathbf{C})$  be the category of models of a  $V$ -theory  $\mathbf{T}$  in a  $V$ -category  $\mathbf{C}$  with finite  $V$ -products, as in Defn. 4. A morphism of  $V$ -theories  $f: \mathbf{T} \rightarrow \mathbf{T}'$  induces a **change of theory** functor between the respective categories of models

$$f^*: V\text{Mod}(\mathbf{T}', \mathbf{C}) \rightarrow V\text{Mod}(\mathbf{T}, \mathbf{C})$$

defined as pre-composition with  $f$ . Similarly, a  $V$ -product-preserving  $V$ -functor  $g: \mathbf{C} \rightarrow \mathbf{C}'$  induces a **change of context** functor

$$g_*: V\text{Mod}(\mathbf{T}, \mathbf{C}) \rightarrow V\text{Mod}(\mathbf{T}, \mathbf{C}')$$

defined as post-composition with  $g$ .

These translations, as well as change of base, can all be packed up nicely using the **Grothendieck construction**: given any functor  $F: \mathbf{D} \rightarrow \text{Cat}$ , there is a category  $\int F$  that encapsulates all of the categories in the image of  $F$ , defined as follows:

$$\begin{array}{ll} \text{objects} & (d, x): d \in \mathbf{D}, x \in F(d) \\ \text{morphisms} & (f: d \rightarrow d', a: F(f)(x) \rightarrow x') \\ \text{composition} & (f, a) \circ (f', a') = (f \circ f', a \circ F(f)(a')). \end{array}$$

Moreover there is a functor  $p_F: \int F \rightarrow \mathbf{D}$  given as follows:

$$\begin{array}{ll} \text{on objects} & p_F: (d, x) \mapsto d \\ \text{on morphisms} & p_F: (f, a) \mapsto f. \end{array}$$

For more details see [8, 13]. We noted in Section 4 that  $V\text{Law}$  and  $\text{Mod}(\mathbf{T}, \mathbf{C})$  can be promoted to  $V$ -categories when  $V$  is complete and cocomplete: this and further conditions imply that we can use the enriched Grothendieck construction [6], but we focus on the ordinary Grothendieck construction for simplicity.

First, this construction lets us bring together all models of all different  $V$ -theories in all different contexts into one category. All the  $V$ -theories are objects of  $V\text{Law}$ , as in Defn. 3. We can also create a category of all “ $V$ -contexts”.

**Definition 15.** Let  $V\text{Con}$ , the **category of  $V$ -contexts** be the category for which an object is a  $V$ -category with finite  $V$ -products and a morphism is a functor that preserves finite  $V$ -products.

There is a functor

$$\mathbf{VMod}: \mathbf{VLaw}^{\text{op}} \times \mathbf{VCon} \rightarrow \mathbf{Cat}$$

that sends any object  $(T, C)$  to  $\mathbf{VMod}(T, C)$  and any morphism  $(f, g)$  to  $f^*g_* = g_*f^*$ . The functoriality of  $\mathbf{VMod}$  summarizes the contravariant change-of-theory and the covariant change-of-context above. Applying the Grothendieck construction we obtain a category  $\int \mathbf{VMod}$ . Technically an object of  $\int \mathbf{VMod}$  is a triple  $(T, C, \mu)$ , but more intuitively it is a model  $\mu: T \rightarrow C$  of any  $V$ -theory  $T$  in any  $V$ -context  $C$ . Similarly, a morphism

$$(f, g, \alpha): (T, C, \mu) \rightarrow (T', C', \mu')$$

in  $\mathbf{VMod}$  consists of:

- a morphism of  $V$ -theories  $f: T' \rightarrow T$ ,
- a  $V$ -functor  $g: C \rightarrow C'$  that preserves finite  $V$ -products, and
- a  $V$ -natural transformation  $\alpha: g \circ \mu \circ f \Rightarrow \mu'$ .

This is a natural way to map between different models of different theories in different contexts.

We can go further by creating a category that even contains all choices of enriching categories  $V$ :

**Definition 16.** Let  $\mathbf{Enr}$  be the category for which an object is a cartesian closed category  $V$  with chosen finite coproducts of the terminal object, and a morphism is a cartesian functor  $F: V \rightarrow W$  preserving the chosen finite coproducts of the initial object.

There is a functor

$$\mathbf{Mod}: \mathbf{Enr} \rightarrow \mathbf{Cat}$$

that maps any object  $V$  to  $\int \mathbf{VMod}$  and any morphism  $F: V \rightarrow W$  to a functor

$$\mathbf{Mod}(F): \int \mathbf{VMod} \rightarrow \int \mathbf{WMod}$$

that has the following effect:

- $\mathbf{Mod}(F)$  maps any object  $(T, C, \mu)$  to the object  $(F_*(T), F_*(C), F_*(\mu))$ .
- $\mathbf{Mod}(F)$  maps any morphism  $(f, g, \alpha)$  to the morphism  $(F_*(f), F_*(g), F_*(\alpha))$ .

Thus, we can use the Grothendieck construction once more to pack up all choices of enrichment into one big category:

**Theorem 17.** There is a category  $\int \mathbf{Mod}$  in which:

- An object is a choice of cartesian closed category  $V$  with chosen finite coproducts of the terminal object, a  $V$ -theory  $T$ , a  $V$ -category  $C$  with finite  $V$ -products, and a model  $\mu: T \rightarrow C$ .
- A morphism is a cartesian functor  $F: V \rightarrow W$  preserving the chosen finite coproducts of the terminal object and a morphism  $(f, g, \alpha): (F_*(T), F_*(C), F_*(\mu)) \rightarrow (T, C, \mu)$  in  $\mathbf{WMod}$ .

This category allows us to formally treat morphisms between objects of “different kinds”, something we often use informally, for example when speaking of a map from a set to a ring, or a group to a topological group. There are many unexplored questions about the large, heterogeneous categories which arise from the Grothendieck construction, regarding what unusual structure may be gained, such as limits and colimits with objects of different types, or identifying “processes” in which the kinds of objects change in an essential way. However, for our purposes we need only recognize that enriched Lawvere theories can be assimilated into one category, which provides a unified context for change of base, change of theory, and change of context.

## 8. APPLICATIONS

In computer science literature, enriched algebraic theories have primarily been studied in the context of “computational effects” [25]. Stay and Meredith have proposed that enriched Lawvere theories can be utilized for the design of programming languages [31]. The primary difference between programming languages and enriched theories, however, is that the latter do not have a notion of *variable binding*—there has certainly been progress on this idea [?], but it leads us beyond our present focus. There are other approaches which instead use an enriched theory as a “compiler”, which can translate a higher language with binding to one without—this idea comes the subject of combinatory logic.

**8.1. The *SKI*-combinator calculus.** The  $\lambda$ -calculus is an elegant formal language which is the foundation of functional computation, the model of intuitionistic logic, and the internal logic of cartesian closed categories: this is the Curry–Howard–Lambek correspondence [3].

Terms are constructed recursively by *variables*, *application*, and *abstraction*, and the basic rewrite is *beta reduction*, which substitutes the applied term for the bound variable:

$$M, N := x \mid (M N) \mid \lambda x.M$$

$$(\lambda x.M N) \Rightarrow M[N/x].$$

Despite the apparent simplicity, there are complications regarding substitution. Consider the term  $M = \lambda x.(\lambda y.(xy))$ : if this is applied to the variable  $y$ , then  $(M y) \Rightarrow \lambda y.(y y)$  — but this is not intended, because the  $y$  in  $M$  is just a placeholder, it is “bound” by whatever will be plugged in, while the  $y$  being substituted is “free”, meaning it can refer to some other value or function in the program. Hence whenever a free variable is to be substituted for a bound variable, we need to rename the bound variable to prevent “variable capture” (e.g.  $(My) \Rightarrow \lambda z.(y z)$ ).

This problem was noticed early in the history of mathematical foundations, even before the  $\lambda$ -calculus, and so Moses Schönfinkel invented **combinatory logic** [27], a basic form of logic without the red tape of variable binding, hence without functions in the usual sense. The *SKI*-calculus is the “variable-free” representation of the  $\lambda$ -calculus;  $\lambda$ -terms are translated via “abstraction elimination” into strings of combinators and applications. This is a technique for programming languages to minimize the subtleties of variables. A great introduction into the strange world of combinators is given by Smullyan [29].

The insight of Stay and Meredith [30] is that even though Lawvere theories have no variables, through abstraction elimination a programming language can be made into an algebraic object. When representing a computational calculus as a **Gph**-theory, the general rewrite rules are simply edges in the hom-graphs  $t^n \rightarrow t$ , with the object  $t$  serving in place of the variable. Below is the theory of the *SKI*-calculus:

	<b>Th(SKI)</b>
<b>type</b>	$t$
<b>term constructors</b>	$S: 1 \rightarrow t$ $K: 1 \rightarrow t$ $I: 1 \rightarrow t$ $(- -): t^2 \rightarrow t$
<b>structural congruence</b>	n/a
<b>rewrites</b>	$\sigma: (((S -) =) \equiv) \Rightarrow ((- \equiv) (= \equiv))$ $\kappa: ((K -) =) \Rightarrow -$ $\iota: (I -) \Rightarrow -$

These rewrites are implicitly universally quantified; i.e. they apply to arbitrary subterms  $-$ ,  $=$ ,  $\equiv$  without any variable binding involved, by using the cartesian structure of the category. (Here  $l, r$  denote the unitors and  $\sigma$  the symmetry of the product.) They are simply edges with vertices:

$$\begin{array}{ccc}
(((S -) =) \equiv): & t^3 \xrightarrow{l^{-1} \times t^3} 1 \times t^3 \xrightarrow{S \times t^3} t^4 \xrightarrow{(- -) \times t^2} t^3 \xrightarrow{(- -) \times t} t^2 \xrightarrow{(- -)} t & \\
\sigma \Downarrow & \Downarrow & \\
((- \equiv) (= \equiv)): & t^3 \xrightarrow{t^2 \times \Delta} t^4 \xrightarrow{t \times \tau \times t} t^4 \xrightarrow{(- -) \times (- -)} t^2 \xrightarrow{(- -)} t & 
\end{array}$$
  

$$\begin{array}{ccc}
((K -) =): & t^2 \xrightarrow{l^{-1} \times t^2} 1 \times t^2 \xrightarrow{K \times t^2} t^3 \xrightarrow{(- -) \times t} t^2 \xrightarrow{(- -)} t & \\
\kappa \Downarrow & \Downarrow & \\
-: & t^2 \xrightarrow{t \times !} t \times 1 \xrightarrow{r} t & 
\end{array}$$
  

$$\begin{array}{ccc}
(I -): & t \xrightarrow{l^{-1}} 1 \times t \xrightarrow{I \times t} t^2 \xrightarrow{(- -)} t & \\
\iota \Downarrow & \Downarrow & \\
-: & t \xrightarrow{t} t & 
\end{array}$$

These abstract rules are evaluated on concrete terms by “plugging in” via precomposition:

$$\begin{array}{ccc}
((KS)I): & 1 \xrightarrow{S \times I} t^2 \xrightarrow{((K -) =)} t & \\
\kappa \circ (S \times I) \Downarrow & \Downarrow & \\
S: & 1 \xrightarrow{S \times I} t^2 \xrightarrow{-} t & 
\end{array}$$

(Morphisms  $1 \rightarrow t$  are the “closed” terms, meaning they have no holes in which to substitute terms; in general morphisms  $t^n \rightarrow t$  are terms with  $n$  holes, or  $n$ -ary operations, and the same reasoning applies.)

A model of this theory is a power-preserving **uGph**-functor  $\mu: \text{Th(SKI)} \rightarrow \mathbf{uGph}$ . This gives a (unital) graph  $\mu(t)$  of all terms and rewrites in the *SKI*-calculus as follows:

$$1 \cong \mu(1) \xrightarrow{\mu(S)} \mu(t) \xrightarrow{\mu((- -)}} \mu(t^2) \cong \mu(t)^2$$

The images of the nullary operations  $S, K, I$  are distinguished vertices of the graph  $\mu(t)$ , because  $\mu$  preserves the terminal object which “points out” vertices. The image of the binary operation  $(- -)$  gives for every pair of vertices  $(u, v) \in \mu(t)^2$ , through the isomorphism  $\mu(t)^2 \cong \mu(t^2)$ , a vertex  $(u \ v)$  in  $\mu(t)$  which is their application. In this way we get all possible terms (writing  $\mu(S), \mu(K), \mu(I)$  as  $S, K, I$  for simplicity):

$$(((S (K (I I))) S) \dots).$$

The rewrites are transferred by the enrichment of the functor: rather than functions between hom-sets, the morphism component of  $\mu$  consists of graph homomorphisms between hom-graphs. So,

$$\mu_{1,t}: \mathbf{Th}(\mathbf{SKI})(1, t) \rightarrow \mathbf{Gph}(1, \mu(t))$$

maps the “syntactic” graph of all closed terms and rewrites coherently into the “semantic” graph, meaning a rewrite in the theory  $a \Rightarrow b$  is sent to a rewrite in the model  $\mu(a) \Rightarrow \mu(b)$ .

These rewrites in the image of  $\mu$  are *graph transformations*, which are just like natural transformations of functors, without the commuting diagram: given two graph homomorphisms  $f, g: G \rightarrow H$ , a graph transformation  $\alpha: f \Rightarrow g$  is a function  $G_0 \rightarrow H_1$  which sends a vertex  $v \in G$  to an edge  $\alpha(v)$  with source  $f(v)$  and target  $g(v)$ .

This is how  $\mu$  realizes  $\mathbf{Th}(\mathbf{SKI})$  as a (unital) graph of terms and rewrites: in the same way that a natural transformation of two constant functors  $a \Rightarrow b: 1 \rightarrow \mathbf{C}$  is a morphism  $a(1) \rightarrow b(1)$  in  $\mathbf{C}$ , a rewrite of closed terms  $a \Rightarrow b: 1 \rightarrow \mu(t)$  corresponds to an edge in  $\mu(t)$ :

$$\mu((I S)) \bullet \xrightarrow{\mu(\iota)} \bullet \mu(S)$$

Finally, the fact that  $\mu((- -))$  is not just a function but a graph homomorphism means that pairs of edges (rewrites)  $(a \rightarrow b, c \rightarrow d)$  are sent to rewrites  $(a \ b) \rightarrow (c \ d)$ . This gives the full complexity of the theory: given a large term (program), there are many different ways it can be computed — and some are better than others:

$$\begin{array}{ccc}
 ((K S) (((S K) I) (I K))) & \xrightarrow{((K S) \sigma)} & ((K S) ((K (I K)) (I (I K)))) \\
 \downarrow \kappa & & \downarrow (((K S) \iota) (I (I K))) \\
 & & ((K S) ((K K) (I (I K)))) \\
 & & \downarrow ((K S) ((K K) (I \iota))) \\
 & & ((K S) ((K K) (I K))) \\
 & & \downarrow ((K S) ((K K) \iota)) \\
 & & ((K S) ((K K) K)) \\
 & & \downarrow ((K S) \kappa) \\
 S & \xleftarrow{\kappa} & ((K S) K)
 \end{array}$$

This process is intuitive, but how do we actually define the model, as a functor, to pick out a specific (unital) graph? There are many models of  $\mathbf{Th}(\mathbf{SKI})$ , but in particular we care about the canonical *free* model, which means that  $\mu(t)$  is simply the graph of all closed terms and rewrites in

the *SKI*-calculus. This utilizes the enriched adjunction of Thm. 5:

$$\begin{array}{ccc} & f_{\text{Gph}} & \\ \text{Gph} & \xrightarrow{\quad} & \text{Mod}(\text{Th}(\text{SKI}), \text{Gph}) \\ & \perp & \\ & u_{\text{Gph}} & \end{array}$$

Then the canonical model of closed terms and rewrites is simply the free model on the empty graph,  $f_{\text{Gph}}(\emptyset)$ , i.e. the  $\mathbf{V}$ -functor  $\mathbf{T}(1, -): \mathbf{T} \rightarrow \mathbf{V}$ . Hence for us, the syntax and semantics of the *SKI* combinator calculus are unified in the model

$$\mu_{\text{SKI}}^{\text{Gph}} := \text{Th}(\text{SKI})(1, -): \text{Th}(\text{SKI}) \rightarrow \text{Gph}.$$

Here we reap the benefits of the abstract construction: the graph  $\mu_{\text{SKI}}^{\text{Gph}}(t)$  is the *transition system* which represents the **small-step operational semantics** of the *SKI*-calculus:

$$(\mu(a) \rightarrow \mu(b) \in \mu_{\text{SKI}}^{\text{Gph}}(t)) \iff (a \Rightarrow b \in \text{Th}(\text{SKI})(1, t)).$$

Interestingly, in the free model on a nonempty graph, the vertices represent designated “ground variables”, and edges represent rewrites of one variable into another. This is potentially useful for “building in” a language with other basic features not intrinsic to the theory.

**8.2. Change of base.** We can now succinctly characterize the transformation from small-step to **big-step** operational semantics. Letting  $i: \Delta \rightarrow \mathbf{Cat}$  denote the inclusion of linear orders into categories, the “realization” functor  $R: \mathbf{uGph} \rightarrow \mathbf{Cat}$  is computed:


$$R(G) = \int^{n \in \Delta} G(n) \times i(n) = \coprod_{n \in \Delta} G(n) \times i(n) / \sim$$

where  $(x, j) \sim (y, k)$  iff there exists  $f: j \rightarrow k$  such that

The same reasoning applies to the “free poset” functor  $\text{FP}: \mathbf{Cat} \rightarrow \mathbf{Pos}$ ; it is a change-of-computation because the product of posets is defined in the same way. This induces the lesser-known **full-step semantics**, which collapses hom-sets to truth values, simply asserting the existence of a rewrite sequence between terms, without distinguishing between different paths. Starting from a free category, this is simply adding the property that all distinct paths between two terms are equal, while retaining transitivity.

Finally, we can pass to the purely abstract realm where all computation is already complete. We take the right adjoint  $\text{FS}: \mathbf{Pos} \rightarrow \mathbf{Set}$  to the functor  $\text{UP}: \mathbf{Set} \rightarrow \mathbf{Pos}$  sending any set to the discrete poset on that set. The functor  $\text{FS}$  collapses every connected component of the poset to a point. This extracts the denotational semantics of the language, by identifying all terms related by rewrites. If the rewrites are terminating and confluent, we can choose a representative term for each equivalence class: the unique term that admits no nontrivial rewrites.

Thus from this simple sequence of functors, we can translate between several important kinds of computation for the *SKI*-calculus. For example, we have the following computation:



The solid arrows are the one-step rewrites of the initial **Gph**-theory; applying  $\text{FC}_*$  gives the dotted composites, and  $\text{FP}_*$  asserts that all composites between any two objects are equal. Finally,  $\text{FS}_*$  collapses the whole diagram to  $S$ . This is a simple demonstration of the basic stages of computation: small-step, big-step, full-step, and denotational semantics.

**8.3. Change-of-theory.** We can equip term calculi with *reduction contexts*, which determine when rewrites are valid, thus giving the language a certain **evaluation strategy**. For example, the “weak head normal form” is given by only allowing rewrites on the left-hand side of the term.

We can do this for  $\text{Th}(\text{SKI})$  by adding a reduction context marker as a unary operation, and a structural congruence rule which pushes the marker to the left-hand side of an application; lastly we modify the rewrite rules to be valid only when the marker is present:

$\text{Th}(\text{SKI} + \text{R})$	
<b>sort</b>	$t$
<b>term constructors</b>	$S, K, I: 1 \rightarrow t$ $R: t \rightarrow t$ $(- -): t^2 \rightarrow t$
<b>structural congruence</b>	$R(- =) = (R- =)$ $RR = R$
<b>rewrites</b>	$\sigma_r: (((RS -) =) \equiv) \Rightarrow ((R- \equiv) (= \equiv))$ $\kappa_r: ((RK -) =) \Rightarrow R-$ $\iota_r: (RI -) \Rightarrow R-$ $n_r: (- R =) \Rightarrow (- =)$

The *SKI*-calculus is thereby equipped with “lazy evaluation”, an essential paradigm in modern programming. This represents a broad potential application of equipping theories with computational methods, such as evaluation strategies.

Moreover, these equipments can be added or removed as needed: using change-of-theory, we can utilize a “free reduction” **Gph**-functor  $f_R: \text{Th}(\text{SKI}) \rightarrow \text{Th}(\text{SKI} + \text{R})$ :

objects	$t^n \mapsto t^n$
hom-vertices	$S, K, I \mapsto S, K, I$ $(- -) \mapsto R(- -)$
hom-edges	$\sigma, \kappa, \iota \mapsto \sigma_r, \kappa_r, \iota_r$



This essentially interprets ordinary *SKI* as having every subterm be a reduction context. This is a **Gph**-functor because its hom component consists of graph-homomorphisms

$$f_{n,m}: \text{Th}(\text{SKI})(t^n, t^m) \rightarrow \text{Th}(\text{SKI} + R)(t^n, t^m)$$

which simply send each application to its postcomposition with  $R$ , and each rewrite to its “marked” correspondent.

So, by precomposition this induces the change of theory on categories of models:

$$f_R^*: \text{Mod}(\text{Th}(\text{SKI} + R), \mathbf{C}) \rightarrow \text{Mod}(\text{Th}(\text{SKI}), \mathbf{C})$$

for all semantic categories  $\mathbf{C}$ , which forgets the reduction contexts.

Similarly, there is a **Gph**-functor  $u_R: \text{Th}(\text{SKI} + R) \rightarrow \text{Th}(\text{SKI})$  which forgets reduction contexts, by sending  $\sigma_r, \kappa_r, \iota_r \mapsto \sigma, \kappa, \iota$  and  $R \mapsto id_t$ ; this latter is the only way that the marked reductions can be mapped coherently to the unmarked. However, this means that  $u_R^*$  does not give the desired change-of-theory of “freely adjoining contexts”, because collapsing  $R$  to the identity eliminates the significance of the marker.

This illustrates a key aspect of categorical universal algebra: because change-of-theory is given by precomposition and is thus contravariant, *properties* (equations) and *structure* (operations) can only be removed. This is a necessary limitation, at least in the present setup, but there are ways to make do. These abstract theories are not floating in isolation but are implemented in code: one can simply use a “maximal theory” with all pertinent structure, then selectively forget as needed.

**8.4. Bisimulation.** This paper uses simple functors to illustrate the basic idea of changing methods of computation. Of course, there are many interesting and useful change-of-base functors. As demonstrated, any functor  $F: \mathbf{V} \rightarrow \mathbf{W}$  which preserves finite products and finite coproducts of the terminal object can be considered as a change in computation. For example, if we enrich in labelled directed graphs, we can utilize the important concept of *bisimulation*.

A labelled transition system consists of a set  $G$ , a label alphabet  $A$ , and a rewrite relation  $\rightarrow \subset G \times A \times G$ , equivalently a graph labelled by elements of  $A$ . The elements of  $G$  represent terms or processes, and the elements of  $A$  represent rewrite rules, in order to actually keep track of which kinds of rewrites are being used in a computation. An element  $(p, a, q)$  is denoted  $p \xrightarrow{a} q$ .

In particular, labelled transition systems allow for the correct definition of process equivalence. A **bisimilarity relation**  $\equiv \subset G \times G$  consists of pairs of processes  $(p, q)$ , written  $p \equiv q$ , defined:

$$\begin{aligned} & \forall a \in A, \quad p', q' \in G \\ & (p \xrightarrow{a} p') \text{ implies } (\exists q' \in G \quad (q \xrightarrow{a} q') \wedge p' \equiv q') \\ & (q \xrightarrow{a} q') \text{ implies } (\exists p' \in G \quad (p \xrightarrow{a} p') \wedge p' \equiv q') \end{aligned}$$

Intuitively, this means that the processes  $p$  and  $q$  can always “match each other’s moves” as they evolve. Then for all intents and purposes, these processes behave the same way, and hence should be considered as operationally equivalent. The **bisimulation** on  $G$  is the largest bisimilarity relation which is also a *congruence*, meaning that processes are bisimilar iff they are so in every context, i.e. when substituted into any one-hole term.

This concept, as well as the calculus of communicating processes, were invented and demonstrated by Milner [22]. The latter can be expressed as an **LTS**-theory. The category of labelled transition systems is just like **Gph**, except of course we now keep track of labels. Morphisms in **LTS**, operations in **LTS**-theories, and **LTS**-functors all preserve labels; for example, when we compose and multiply rewrite rules, we retain this information by labelling with the actual denotation for that composite/product. Modulo these details,  $\mathbf{V} = \mathbf{LTS}$  is exactly like the cases considered above.

## Th(CCS)

<b>types</b>	$P$ $N$ $\overline{N}$	processes actions coactions
<b>operations</b>	$0: 1 \rightarrow P$ $\tau: 1 \rightarrow P$ $ : P^2 \rightarrow P$ $+: P^2 \rightarrow P$ $\cdot: N \times P \rightarrow P$ $\cdot: \overline{N} \times P \rightarrow P$	nullity internal action parallel choice input output
<b>congruence</b>	$(P,  , 0)$ $(P, +, 0)$	commutative monoid commutative monoid
<b>rewrites</b>	$\text{tau}: \cdot \circ (\tau \times P) \circ l^{-1} \Rightarrow id_P$ $\text{inter}: + \circ   \circ (\cdot \times \cdot) \Rightarrow  $	$(\tau.P \xrightarrow{\text{tau}} P)$ $(P' + a.P \overline{a}.Q \xrightarrow{\text{react}} P Q)$

The theory is summarized in the two rewrite rules: *tau* is an “unobservable” action, a process evolving in a way that is private to the ambient context; *inter* is interaction or communication - the action  $a$  is being triggered by the coaction  $\overline{a}$ , they are used up and the sequential processes continue in parallel. This calculus is the precursor to the  $\pi$  calculus [21], and is a very simple and general framework for understanding systems of interacting automata.

There is an endofunctor  $B: \text{LTS} \rightarrow \text{LTS}$  which quotients by the bisimulation relation. It preserves products,  $B(G \times H) \cong B(G) \times B(H)$ , because  $(p_1, p_2) \equiv (q_1, q_2)$  iff  $(p_1 \equiv q_1 \text{ and } p_2 \equiv q_2)$ . Thus we can utilize base change to perform a very useful transformation on our computation: from  $\text{Th}(\text{CCS})$ , we get a new theory  $B_*(\text{Th}(\text{CCS}))$ , the hom-LTS’s of which consist of bisimulation equivalence classes of terms and rewrites in the calculus of communicating systems.

## 9. CONCLUSION

We have established the basics of how enriched Lawvere theories provide a framework for unifying the structure and behavior of formal languages. Enriching theories in category-like structures reifies operational semantics by incorporating rewrites between terms, and cartesian functors between enriching categories induce change-of-semantics functors between categories of models—this simplified condition is obtained by using only natural number arities.

Change of base, along with change of theory and change of context, can be used to create a single category  $\text{Mod}$ , which consists of all models of all enriched Lawvere theories in all contexts. We have demonstrated these concepts with the theory of combinatory logic,  $\text{Th}(\text{SKI})$ , describing a change of computation from small-step to big-step to full-step to denotational semantics. Finally, we suggest that there are many interesting change-of-computation functors, by considering an endofunctor on the category of labelled transition systems, which quotients by the bisimulation relation and is indeed a change of computation.

## REFERENCES

- [1] J. Adámek and J. Rosický, *Locally Presentable and Accessible Categories*, Cambridge U. Press, Cambridge, 1994. (Referred to on page 5, 8.)
- [2] J. Baez, Can 1+1 have more than two points?, *The n-Category Café*, April 19, 2019. Available at [https://golem.ph.utexas.edu/category/2019/04/can\\_11\\_have\\_more\\_than\\_two\\_poin.html](https://golem.ph.utexas.edu/category/2019/04/can_11_have_more_than_two_poin.html). (Referred to on page 3, 16.)
- [3] J. Baez and M. Stay, Physics, topology, logic and computation: a Rosetta Stone, in *New Structures for Physics*, ed. B. Coecke, Springer, Berlin, 2011, pp. 95–172. Available as [arXiv:0903.0340](https://arxiv.org/abs/0903.0340). (Referred to on page 21.)
- [4] H.P. Barendregt, The Lambda Calculus, its syntax and semantics, in *Studies in Logic and The Foundations of Mathematics*, Elsevier, London, 1984. (Referred to on page .)
- [5] M. Barr and C. Wells, *Toposes, Triples and Theories*, reprinted in *Repr. Theory Appl. Categ.* **12** (2005). Available at <http://www.tac.mta.ca/tac/reprints/articles/12/tr12abs.html>. (Referred to on page 4, 5.)
- [6] J. Beardsley and L. Z. Wong, The enriched Grothendieck construction. Available as [arXiv:1804.03829](https://arxiv.org/abs/1804.03829). (Referred to on page 19.)
- [7] R. Blackwell, G. M. Kelly and A. J. Power, Two-dimensional monad theory, *J. Pure Appl. Algebra* **59** (1989), 1–41. (Referred to on page 13.)
- [8] F. Borceux, *Handbook of Categorical Algebra*, vol. 2, Cambridge U. Press, Cambridge, 1994. (Referred to on page 17, 19.)
- [9] R. Crole, *Categories for Types*, Cambridge U. Press, Cambridge, 1993. (Referred to on page 4.)
- [10] B. Day and R. Street, Monoidal categories and Hopf algebroids, *Adv. Math.* **127** (1997), 99–157. (Referred to on page 11, 14.)
- [11] E. J. Dubuc, *Kan Extensions in Enriched Category Theory*, Lecture Notes in Mathematics, Springer, Berlin, 1970. (Referred to on page 13.)
- [12] M. Hyland and J. Power, The category theoretic understanding of universal algebra: Lawvere theories and monads, in *Electron. Notes Theor. Comput. Sci.* **172** (2007), 437–458. (Referred to on page 4.)
- [13] B. Jacobs, *Categorical Logic and Type Theory*, Elsevier, Amsterdam, 1999. (Referred to on page 19.)
- [14] G. M. Kelly, *Basic Concepts of Enriched Category Theory*, reprinted in *Repr. Theory Appl. Categ.* **10** (2005), 1–136. Available at <http://www.tac.mta.ca/tac/reprints/articles/10/tr10abs.html>. (Referred to on page 5, 6.)
- [15] Y. Kinoshita, J. Power and M. Takeyama, Sketches, *J. Pure Appl. Algebra* **143** (1999), 275–291. (Referred to on page 8.)
- [16] F. W. Lawvere, Functorial semantics of algebraic theories, reprinted in *Repr. Theory Appl. Categ.* **5** (2004), 1–121. Available at <http://tac.mta.ca/tac/reprints/articles/5/tr5abs.html>. (Referred to on page 1, 5.)
- [17] F. E. J. Linton, Some aspects of equational theories, in *Proceedings of the Conference on Categorical Algebra*, eds. S. Eilenberg et al., Springer, Berlin, 1965. (Referred to on page 4.)
- [18] R. B. B. Lucyshyn-Wright, Enriched algebraic theories and monads for a system of arities, *Theory Appl. Categ.* **31** (2016), 101–137. Available at <http://www.tac.mta.ca/tac/volumes/31/5/31-05abs.html>. (Referred to on page 3, 8, 9.)
- [19] C. Lüth and N. Ghani, Monads and modular term rewriting, *Category Theory and Computer Science (Santa Margherita Ligure, 1997)*, Springer, Berlin, 1997, pp. 69–86. Available at <http://www.informatik.uni-bremen.de/ctl/papers/ctcs97l.pdf>. (Referred to on page 2.)
- [20] B. Milewski, *Category Theory for Programmers*, Chap. 14: Lawvere theories, 2017. Available at <https://bartoszmilewski.com/2017/08/26/lawvere-theories/>. (Referred to on page 5.)
- [21] R. Milner, The polyadic  $\pi$ -calculus: a tutorial, in *Logic and Algebra of Specification*, Springer, Berlin, 1993, 203–246. (Referred to on page 27.)
- [22] R. Milner, *Communicating and Mobile Systems: The Pi Calculus*, in *Cambridge University Press*, Cambridge, UK, 1999. (Referred to on page 26.)
- [23] K. Nishizawa and J. Power, Lawvere theories enriched over a general base, *J. Pure Appl. Algebra* **213** (2009), 377–386. (Referred to on page 8.)
- [24] G. D. Plotkin, A structural approach to operational semantics, *J. Log. Algebr. Program.* **60/61** (2004) 17–139. Available at [http://homepages.inf.ed.ac.uk/gdp/publications/sos\\_jlap.pdf](http://homepages.inf.ed.ac.uk/gdp/publications/sos_jlap.pdf). (Referred to on page 2.)
- [25] M. Hyland and J. Power, Discrete Lawvere theories and computational effects, in *Theoretical Comp. Sci.* **366** (2006), 144–162. Available at . (Referred to on page 2, 21.)
- [26] J. Power, Enriched Lawvere theories, *Theory Appl. Categ.* **6** (1999), 83–93. (Referred to on page 8.)
- [27] M. Schönfinkel, Über die Bausteine der mathematischen Logik, *Mathematische Annalen* **92** (1924), 305–316. Available at <http://www.digizeitschriften.de/dms/img/?PID=GDZPPN002270110>. (Referred to on page 21.)

- [28] R. A. G. Seely, Modelling computations: a 2-categorical framework, in *Proceedings of the Second Annual IEEE Symposium on Logic in Computer Science (LICS 1987)*, IEEE Computer Society Press, Ithaca, New York, pp. 22–25. (Referred to on page [2](#).)
- [29] R. Smullyan, *To Mock a Mockingbird: and Other Logic Puzzles Including an Amazing Adventure in Combinatory Logic*, Oxford U. Press, Oxford, 2000. (Referred to on page [21](#).)
- [30] M. Stay and L. G. Meredith, Representing operational semantics with enriched Lawvere theories. Available as [arXiv:1704.03080](#). (Referred to on page [3](#), [21](#).)
- [31] M. Stay and L. G. Meredith, Logic as a distributive law. Available as [arXiv:1610.02247](#). (Referred to on page [21](#).)
- [32] G. Friedman, An elementary introduction to simplicial sets. Available as [.](#) (Referred to on page [2](#).)