

MATH 136: Linear Algebra I

Charles Shen

Winter 2015, University of Waterloo

Quick reference sheet.

Contents

| | | |
|----------|--------------------------------------|----------|
| 1 | Matrices | 1 |
| 1.1 | Elementary Row Operations | 1 |
| 1.2 | Reduced Row Echelon Form | 1 |
| 1.3 | System Rank Theorem | 1 |
| 2 | Inverses | 2 |
| 2.1 | Inverse of a 2-by-2 Matrix | 2 |
| 2.2 | Invertible Matrix Theorem | 2 |

1 Matrices

Definition. An **augmented matrix** has the form $\begin{bmatrix} A & | & \vec{b} \end{bmatrix}$

Definition. A **homogeneous matrix** has the form $\begin{bmatrix} A & | & \vec{0} \end{bmatrix}$

A homogeneous system is always consistent, $\vec{x} = \vec{0}$ is always a solution.

1.1 Elementary Row Operations

1. Add a multiple of one row to another: $R_i + cR_j$
2. Multiple a row by a non-zero constant: cR_i
3. Exchange two rows: $R_i \leftrightarrow R_j$

1.2 Reduced Row Echelon Form

Definition. A matrix R is said to be in **Reduced Row Echelon Form** (RREF) if:

1. All rows containing a non-zero entry are above rows which only contains zeros
2. The first non-zero entry in each non-zero row is 1, called a **leading one** (or a pivot)
3. The leading one in each non-zero row is to the right of the leading one in any row above it
4. A leading one is the only non-zero entry in its column

Definition. The **rank** of a matrix A is the number of leading ones in the RREF of the matrix.

1.3 System Rank Theorem

Theorem. Let A be the coefficient matrix of a system of m linear equations in n unknowns $\begin{bmatrix} A & | & \vec{b} \end{bmatrix}$

1. If the rank of A is less than the rank of the augmented matrix $\begin{bmatrix} A & | & \vec{b} \end{bmatrix}$, then the system is inconsistent

2. If the system $\begin{bmatrix} A & | & \vec{b} \end{bmatrix}$ is consistent, then the system contains $(n - \text{rank}(A))$ free variables
3. $\text{rank}(A) = m$ iff the system $\begin{bmatrix} A & | & \vec{b} \end{bmatrix}$ is consistent for every $\vec{b} \in \mathbb{R}^m$

2 Inverses

2.1 Inverse of a 2-by-2 Matrix

$$\text{Suppose } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\text{Then } A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

2.2 Invertible Matrix Theorem

Theorem. For any $n \times n$ matrix A , the following are equivalent:

1. A is invertible
2. $\text{rank}(A) = n$
3. RREF of $A = I$
4. $A\vec{x} = \vec{b}$ is consistent for all $\vec{b} \in \mathbb{R}^n$
5. The columns of A form a linearly independent set
6. The columns of A span \mathbb{R}^n
7. The columns of A form a basis for \mathbb{R}^n
8. The rows of A form a basis for \mathbb{R}^n
9. $\text{Col}(A) = \mathbb{R}^n$
10. $\text{Null}(A) = \{\vec{0}\}$
11. A^T is invertible
12. $\text{Row}(A) = \mathbb{R}^n$
13. $\text{Null}(A^T) = \{\vec{0}\}$