

COMMONWEALTH OF AUSTRALIA

Copyright Regulations 1969

WARNING

This material has been reproduced and communicated to you by or on behalf of the University of Sydney pursuant to Part VB of the Copyright Act 1968 (**the Act**). The material in this communication may be subject to copyright under the Act. Any further copying or communication of this material by you may be the subject of copyright protection under the Act.


Do not remove this notice.

COMPx270: Randomised and
Advanced Algorithms
Lecture 12: Learning from
experts

Clément Canonne
School of Computer Science



THE UNIVERSITY OF
SYDNEY



A question

You invest in the stock market. Each morning, you have to decide whether to **sell** or **buy**.

A question

You invest in the stock market. Each morning, you have to decide whether to **sell** or **buy**. At the end of the day, you see if you made the right decision. If you did, great: that day, you made money.

A question

You invest in the stock market. Each morning, you have to decide whether to **sell** or **buy**. At the end of the day, you see if you made the right decision. If you did, great: that day, you made money.

However, you don't know anything about the stock market.

A question

You invest in the stock market. Each morning, you have to decide whether to **sell** or **buy**. At the end of the day, you see if you made the right decision. If you did, great: that day, you made money.

However, you don't know anything about the stock market.

But you have many friends who do: they're all "experts."

A question

So every morning, before you make your decision, all those friends will give you their advice.



A question

So every morning, before you make your decision, all those friends will give you their advice.



Some might collude, or be completely wrong, or even try to make you lose money. But each of them will tell you to either **sell** or **buy**.

A question

Then, based on those many pieces of advice, **you** decide.

(And you do that again, every day.)



A question

Then, based on those many pieces of advice, **you** decide.

(And you do that again, every day.)

What is a good strategy to make money?



A question


Then, based on those many pieces of advice, **you** decide.

(And you do that again, every day.)

What is a **provably** good strategy to make money?



Let's make this formal



- There are n experts.
- Each day, $t = 1, \dots, T$, each of them makes a prediction $v_{i,t} \in \{0,1\}$
- Based on those, you make your own prediction $\hat{u}_t \in \{0,1\}$
- Then the “true” value $u_t \in \{0,1\}$ is revealed
- If $\hat{u}_t \neq u_t$, this counts as a **mistake** (mistakes are bad)



Goal: minimise **total number of mistakes** $M = \sum_{t=1}^T 1_{\hat{u}_t \neq u_t}$

- There are n experts.
- Each day, $t = 1, \dots, T$, each of them makes a prediction $v_{i,t} \in \{0,1\}$
- Based on those, you make your own prediction $\hat{u}_t \in \{0,1\}$
- Then the “true” value $u_t \in \{0,1\}$ is revealed
- If $\hat{u}_t \neq u_t$, this counts as a **mistake** (mistakes are bad)



Goal: minimise **total number of mistakes** $M = \sum_{t=1}^T 1_{\hat{u}_t \neq u_t}$

But what do we mean by this? We don't assume **anything** on the experts or on the true values. **They could even all be adversarial!**

- There are n experts.
- Each day, $t = 1, \dots, T$, each of them makes a prediction $v_{i,t} \in \{0,1\}$
- Based on those, you make your own prediction $\hat{u}_t \in \{0,1\}$
- Then the “true” value $u_t \in \{0,1\}$ is revealed
- If $\hat{u}_t \neq u_t$, this counts as a **mistake** (mistakes are bad)




Goal: minimise **total number of mistakes** $M = \sum_{t=1}^T 1_{\hat{u}_t \neq u_t}$ compared to the **best expert** (whoever that is).

Not make much more mistakes than the **best advice in hindsight**.

Warmup: a Perfect Expert




- There are n experts. **Suppose one of them (unknown) is always right.**
- Each day, $t = 1, \dots, T$, each of them makes a prediction $v_{i,t} \in \{0,1\}$
- Based on those, you make your own prediction $\hat{u}_t \in \{0,1\}$
- Then the “true” value $u_t \in \{0,1\}$ is revealed
- If $\hat{u}_t \neq u_t$, this counts as a **mistake** (mistakes are bad)




Goal: minimise **total number of mistakes** $M = \sum_{t=1}^T 1_{\hat{u}_t \neq u_t}$

Theorem. There is a strategy guaranteeing $M \leq n - 1$, regardless of T (even for $T = \infty$).




```
Set  $S \leftarrow [n]$ 
for all  $1 \leq t \leq T$  do
  Receive  $v_{1,t}, \dots, v_{n,t}$ 
  if  $|S| \geq 1$  then
    Pick any  $i \in S$                                 ▷ Lexicographically, for instance
    Choose  $\hat{u}_t \leftarrow v_{i,t}$ 
  else
    Choose  $\hat{u}_t \leftarrow 0$                                 ▷ Arbitrary
  Receive  $u_t$                                 ▷ Observe the truth
   $S \leftarrow S \setminus \{i \in S : v_{i,t} \neq u_t\}$     ▷ Remove all mistaken experts
```

Theorem. There is a strategy guaranteeing $M \leq n - 1$, regardless of T (even for $T = \infty$).




Proof.

Theorem. There is a strategy guaranteeing $M \leq \log_2 n$, regardless of T (even for $T = \infty$).




Claim. There is a strategy guaranteeing $M \leq \log_2 n$, regardless of T (even for $T = \infty$).




Algorithm: Start with $S = \{1, 2, \dots, n\}$. Each day, choose \hat{u}_t to be the **majority** of advices from experts still in S . At the end of the day, remove from S all experts who predicted wrong.

Claim. There is a strategy guaranteeing $M \leq \log_2 n$, regardless of T (even for $T = \infty$).



```
Set  $S \leftarrow [n]$ 
for all  $1 \leq t \leq T$  do
    Receive  $v_{1,t}, \dots, v_{n,t}$ 
    if  $|S| \geq 1$  then
        Choose  $\hat{u}_t \leftarrow \text{maj}_{i \in S} v_{i,t}$            ▷ Take the majority advice
    else
        Choose  $\hat{u}_t \leftarrow 0$                              ▷ Arbitrary
    Receive  $u_t$                                              ▷ Observe the truth
     $S \leftarrow S \setminus \{i \in S : v_{i,t} \neq u_t\}$     ▷ Remove all mistaken experts
```


Claim. There is a strategy guaranteeing $M \leq \log_2 n$, regardless of T (even for $T = \infty$).



Algorithm: Start with $S = \{1, 2, \dots, n\}$. Each day, choose \hat{u}_t to be the **majority** of advices from experts still in S . At the end of the day, remove from S all experts who predicted wrong.

Proof of correctness. Every time we make a mistake, at least half the experts in S must have been wrong (we took the majority vote). So after each mistake the size of S is at least halved.


Claim. There is a strategy guaranteeing $M \leq \log_2 n$, regardless of T (even for $T = \infty$).



Algorithm: Start with $S = \{1, 2, \dots, n\}$. Each day, choose \hat{u}_t to be the **majority** of advices from experts still in S . At the end of the day, remove from S all experts who predicted wrong.

Proof of correctness. Every time we make a mistake, **at least half** the experts in S must have been wrong (we took the majority vote). So after each mistake the size of S is at least **halved**. But we always have $|S| \geq 1$, since (by assumption) there exists an expert who is always right (and therefore never gets removed).

Claim. There is a strategy guaranteeing $M \leq \log_2 n$, regardless of T (even for $T = \infty$).



Algorithm: Start with $S = \{1, 2, \dots, n\}$. Each day, choose \hat{u}_t to be the **majority** of advices from experts still in S . At the end of the day, remove from S all experts who predicted wrong.

Proof of correctness. Since we started with $|S| = n$, our total number M of mistakes must then satisfy

$$\frac{n}{2^M} \geq 1$$

that is, $M \leq \log_2 n$.

Nobody's Perfect



This is great! But... things completely fail if there is no “perfect expert.”

What if even the **best** expert made some mistakes? Can we make things **robust**?

This is great! But... things completely fail if there is no “perfect expert.”

What if even the **best** expert made some mistakes? Can we make things **robust**?

Let's revisit the algorithm.

We had n **weights** w_1, \dots, w_n initialised to 1.

At day t , our prediction was $\hat{u}_t \leftarrow \text{Maj}(w_1 v_{1,t} + \dots + w_n v_{n,t})$

Whenever expert i made a mistake, we set $w_i \leftarrow 0 \cdot w_i$.

This is great! But... things completely fail if there is no “perfect expert.”

What if even the **best** expert made some mistakes? Can we make things **robust**?

Let's revisit the algorithm.

We had n **weights** w_1, \dots, w_n initialised to 1.

At day t , our prediction was $\hat{u}_t \leftarrow \text{Maj}(w_1 v_{1,t} + \dots + w_n v_{n,t})$

Whenever expert i made a mistake, we set $w_i \leftarrow \mathbf{0} \cdot w_i$.

This is great! But... things completely fail if there is no “perfect expert.”

What if even the **best** expert made some mistakes? Can we make things **robust**?

Let's revisit the algorithm.

We have n **weights** w_1, \dots, w_n initialised to 1.

At day t , our prediction is $\hat{u}_t \leftarrow \text{Maj}(w_1 v_{1,t} + \dots + w_n v_{n,t})$

Whenever expert i made a mistake, we set $w_i \leftarrow \frac{1}{2} \cdot w_i$.

Algorithm (Multiplicative Weights Update).

Start with n **weights** w_1, \dots, w_n initialised to 1.

Each day, choose the **weighted majority** $\hat{u}_t \leftarrow \text{Maj}(w_1 v_{1,t} + \dots + w_n v_{n,t})$

At the end of the day, set $w_i \leftarrow \frac{1}{2} \cdot w_i$ for expert i made a mistake.

Set $w_1, \dots, w_n \leftarrow 1$

for all $1 \leq t \leq T$ **do**

Receive $v_{1,t}, \dots, v_{n,t}$

Choose $\hat{u}_t \leftarrow \text{sign}\left(\sum_{i=1}^n w_i v_{i,t} \geq \frac{1}{2} \sum_{i=1}^n w_i\right)$ \triangleright Weighted majority

Receive u_t

\triangleright Observe the truth

for all $1 \leq i \leq n$ **do**

\triangleright Penalise all mistaken experts

$$w_i \leftarrow \begin{cases} \frac{1}{2} w_i & \text{if } v_{i,t} \neq u_t \\ w_i & \text{otherwise.} \end{cases}$$

```

Set  $w_1, \dots, w_n \leftarrow 1$ 
for all  $1 \leq t \leq T$  do
    Receive  $v_{1,t}, \dots, v_{n,t}$ 
    Choose  $\hat{u}_t \leftarrow \text{sign}\left(\sum_{i=1}^n w_i v_{i,t} \geq \frac{1}{2} \sum_{i=1}^n w_i\right)$   $\triangleright$  Weighted
    majority
    Receive  $u_t$   $\triangleright$  Observe the truth
    for all  $1 \leq i \leq n$  do  $\triangleright$  Penalise all mistaken experts
         $w_i \leftarrow \begin{cases} \frac{1}{2} w_i & \text{if } v_{i,t} \neq u_t \\ w_i & \text{otherwise.} \end{cases}$ 

```

Theorem 59. *There is a (deterministic) algorithm (Algorithm 24) such that*

$$C(T) \leq \frac{C^*(T) + \log_2 n}{\log_2 \frac{4}{3}} \leq 2.41(C^*(T) + \log_2 n).$$

Moreover, this holds even when $T = \infty$.



Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Proof. Let W_t be the total weights of experts on day t . Initially, $W_0 = n$. Every time we make a mistake, this means **at least half the weight** was on experts who did a mistake (since we took the weighted majority). So if we made a mistake at day t ,

$$W_{t+1} = W_t^{\text{good}} + \frac{1}{2} W_t^{\text{bad}} \leq \frac{1}{2} W_t + \frac{1}{2} \cdot \frac{1}{2} W_t = \frac{3}{4} W_t$$

Now, look at the **best expert** (in hindsight). They made M^* mistakes, so their final weight is $(1/2)^{M^*}$.

Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Proof. Let W_t be the total weights of experts on day t . Initially, $W_0 = n$. Every time we make a mistake, this means **at least half the weight** was on experts who did a mistake (since we took the weighted majority). So if we made a mistake at day t ,

$$W_{t+1} = W_t^{\text{good}} + \frac{1}{2} W_t^{\text{bad}} \leq \frac{1}{2} W_t + \frac{1}{2} \cdot \frac{1}{2} W_t = \frac{3}{4} W_t$$

Now, look at the **best expert** (in hindsight). They made M^* mistakes, so their final weight is $(1/2)^{M^*}$. So the final **total** weight is $W_T \geq (1/2)^{M^*}$.

Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Proof. Let W_t be the total weights of experts on day t . Initially, $W_0 = n$. Every time we make a mistake, this means **at least half the weight** was on experts who did a mistake (since we took the weighted majority). So if we made a mistake at day t ,

$$W_{t+1} = W_t^{\text{good}} + \frac{1}{2} W_t^{\text{bad}} \leq \frac{1}{2} W_t + \frac{1}{2} \cdot \frac{1}{2} W_t = \frac{3}{4} W_t$$

Now, look at the **best expert** (in hindsight). They made M^* mistakes, so their final weight is $(1/2)^{M^*}$. So the final **total** weight is $W_T \geq (1/2)^{M^*}$.

Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Proof. Putting it all together:

$$\left(\frac{1}{2}\right)^{M^*} \leq W_T \leq \left(\frac{3}{4}\right)^M W_0 = \left(\frac{3}{4}\right)^M n$$

Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Proof. Putting it all together:

$$\left(\frac{1}{2}\right)^{M^*} \leq W_T \leq \left(\frac{3}{4}\right)^M W_0 = \left(\frac{3}{4}\right)^M n$$

Now, we take the logarithm:

$$-M^* \leq M \log_2 \left(\frac{3}{4}\right) + \log_2 n$$

Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Proof. Putting it all together:

$$\left(\frac{1}{2}\right)^{M^*} \leq W_T \leq \left(\frac{3}{4}\right)^M W_0 = \left(\frac{3}{4}\right)^M n$$

Now, we take the logarithm:

$$-M^* \leq M \log_2 \left(\frac{3}{4}\right) + \log_2 n$$

and get

$$M \leq \frac{M^* + \log_2 n}{-\log_2 \left(\frac{3}{4}\right)}$$

Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Proof. Putting it all together:

$$\left(\frac{1}{2}\right)^{M^*} \leq W_T \leq \left(\frac{3}{4}\right)^M W_0 = \left(\frac{3}{4}\right)^M n$$

Now, we take the logarithm:


$$-M^* \leq M \log_2 \left(\frac{3}{4}\right) + \log_2 n$$

and get

$$M \leq \frac{M^* + \log_2 n}{-\log_2 \left(\frac{3}{4}\right)} \leq 2.41(M^* + \log_2 n)$$



Let's go further!



This is what we proved:

Algorithm (Multiplicative Weights Update).

Start with n **weights** w_1, \dots, w_n initialised to 1.

Each day, choose the **weighted majority** $\hat{u}_t \leftarrow \text{Maj}(w_1 v_{1,t} + \dots + w_n v_{n,t})$

At the end of the day, set $w_i \leftarrow \frac{1}{2} \cdot w_i$ for expert i made a mistake.

Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

This is what we proved:

Algorithm (Multiplicative Weights Update).

Start with n **weights** w_1, \dots, w_n initialised to 1.

Each day, choose the **weighted majority** $\hat{u}_t \leftarrow \text{Maj}(w_1 v_{1,t} + \dots + w_n v_{n,t})$

At the end of the day, set $w_i \leftarrow \frac{1}{2} \cdot w_i$ for expert i made a mistake.

Theorem. The MWU algorithm guarantees $M \leq 2.41(M^* + \log_2 n)$, where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Using exactly the same argument (*try it!*), we get, for any $\beta \in (0,1)$:

Algorithm (Multiplicative Weights Update).

Start with n **weights** w_1, \dots, w_n initialised to 1.

Each day, choose the **weighted majority** $\hat{u}_t \leftarrow \text{Maj}(w_1 v_{1,t} + \dots + w_n v_{n,t})$

At the end of the day, set $w_i \leftarrow \beta \cdot w_i$ for expert i made a mistake.

Theorem. The MWU algorithm guarantees $M \leq \frac{M^* \log_2(1/\beta) + \log_2 n}{\log_2(\frac{2}{1+\beta})}$,

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Using exactly the same argument we get, for any $\beta \in (0,1)$:

Theorem. The MWU algorithm guarantees

$$M \leq \frac{M^* \log_2 \left(\frac{1}{\beta} \right) + \log_2 n}{\log_2 \left(\frac{2}{1 + \beta} \right)}$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).


Using exactly the same argument we get, for any $\beta = 1 - \varepsilon \in (0,1)$:

Theorem. The MWU algorithm guarantees

$$M \leq \frac{M^* \log_2 \left(\frac{1}{\beta} \right) + \log_2 n}{\log_2 \left(\frac{2}{1 + \beta} \right)} \approx 2 \left(M^* + \frac{\ln n}{\varepsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Is that tight?



Theorem. The MWU algorithm guarantees

$$M \approx 2 \left(M^* + \frac{\ln n}{\varepsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Theorem. The MWU algorithm guarantees

$$M \approx 2 \left(M^* + \frac{\ln n}{\varepsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Can we improve that factor **2**?

Theorem. The MWU algorithm guarantees

$$M \approx 2 \left(M^* + \frac{\ln n}{\varepsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Can we improve that factor **2**? **No.**

Theorem. The MWU algorithm guarantees

$$M \approx 2 \left(M^* + \frac{\ln n}{\varepsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Can we improve that factor **2**? **No**. Consider two sets of $n/2$ experts, where experts in the first set are wrong on odd-numbered days, and those in the second set are wrong on even days. That will force T mistakes (while the best experts make $T/2$).

Theorem. The MWU algorithm guarantees

$$M \approx 2 \left(M^* + \frac{\ln n}{\varepsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).


Can we improve that factor **2**? **Yes.**

Theorem. The MWU algorithm guarantees

$$M \approx 2 \left(M^* + \frac{\ln n}{\varepsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Can we improve that factor **2**? **Yes.** With randomisation! Instead of deterministically choosing the weighted majority, pick the answer **at random according to the weights.**




Theorem. The MWU algorithm guarantees

$$M \approx 2 \left(M^* + \frac{\ln n}{\epsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Can we improve that factor **2**? **Yes.** With randomisation! Instead of deterministically choosing the weighted majority, pick the answer **at random according to the weights**. Improves the constant 2 to some $c < 2$.





Theorem. The MWU algorithm guarantees

$$M \approx 2 \left(M^* + \frac{\ln n}{\epsilon} \right)$$

where M^* is the # of mistakes made by the best expert. This holds regardless of T (even for $T = \infty$).

Can we improve that factor **2**? **Yes.** With randomisation! Instead of deterministically choosing the weighted majority, pick the answer **at random according to the weights**. Improves the constant 2 to some $c < 2$. (But only guarantee on **expected** number of mistakes).





Input: Penalty parameter $\beta \in (0, 1)$

Set $w_1, \dots, w_n \leftarrow 1$

for all $1 \leq t \leq T$ **do**

Receive $v_{1,t}, \dots, v_{n,t}$

Draw $I \in [n]$ according to the weights:

$$\Pr[I = i] = \frac{w_i}{\sum_{i=1}^n w_i}, \quad i \in [n]$$

Choose $\hat{u}_t \leftarrow v_{I,t}$

▷ One expert gets the vote

Receive u_t

▷ Observe the truth

for all $1 \leq i \leq n$ **do**

▷ Penalise all mistaken experts

$$w_i \leftarrow \begin{cases} \beta w_i & \text{if } v_{i,t} \neq u_t \\ w_i & \text{otherwise.} \end{cases}$$

Input: Penalty parameter $\beta \in (0, 1)$
 Set $w_1, \dots, w_n \leftarrow 1$
for all $1 \leq t \leq T$ **do**
 Receive $v_{1,t}, \dots, v_{n,t}$
 Draw $I \in [n]$ according to the weights:

$$\Pr[I = i] = \frac{w_i}{\sum_{i=1}^n w_i}, \quad i \in [n]$$

 Choose $\hat{u}_t \leftarrow v_{I,t}$ ▷ One expert gets the vote
 Receive u_t ▷ Observe the truth
 for all $1 \leq i \leq n$ **do** ▷ Penalise all mistaken experts

$$w_i \leftarrow \begin{cases} \beta w_i & \text{if } v_{i,t} \neq u_t \\ w_i & \text{otherwise.} \end{cases}$$

Theorem 61. *There is a (randomised) algorithm (Algorithm 26) such that*

$$\mathbb{E}[\mathcal{C}(T)] \leq \frac{\mathcal{C}^*(T) \ln(1/\beta) + \ln n}{1 - \beta}.$$

Moreover, this holds even when $T = \infty$.

Theorem 61. *There is a (randomised) algorithm (Algorithm 26) such that*

$$\mathbb{E}[\mathcal{C}(T)] \leq \frac{\mathcal{C}^*(T) \ln(1/\beta) + \ln n}{1 - \beta}.$$

Moreover, this holds even when $T = \infty$.

Input: Penalty parameter $\beta \in (0, 1)$

Set $w_1, \dots, w_n \leftarrow 1$

for all $1 \leq t \leq T$ **do**

 Receive $v_{1,t}, \dots, v_{n,t}$

 Draw $I \in [n]$ according to the weights:

$$\Pr[I = i] = \frac{w_i}{\sum_{i=1}^n w_i}, \quad i \in [n]$$

 Choose $\hat{u}_t \leftarrow v_{I,t}$

 ▷ One expert gets the vote

 Receive u_t

 ▷ Observe the truth

for all $1 \leq i \leq n$ **do**

 ▷ Penalise all mistaken experts

$$w_i \leftarrow \begin{cases} \beta w_i & \text{if } v_{i,t} \neq u_t \\ w_i & \text{otherwise.} \end{cases}$$

Theorem 61. *There is a (randomised) algorithm (Algorithm 26) such that*

$$\mathbb{E}[\mathcal{C}(T)] \leq \frac{\mathcal{C}^*(T) \ln(1/\beta) + \ln n}{1 - \beta}.$$

Moreover, this holds even when $T = \infty$.

Input: Penalty parameter $\beta \in (0, 1)$

Set $w_1, \dots, w_n \leftarrow 1$

for all $1 \leq t \leq T$ **do**

Receive $v_{1,t}, \dots, v_{n,t}$

Draw $I \in [n]$ according to the weights:

$$\Pr[I = i] = \frac{w_i}{\sum_{i=1}^n w_i}, \quad i \in [n]$$

Choose $\hat{u}_t \leftarrow v_{I,t}$

▷ One expert gets the vote

Receive u_t

▷ Observe the truth

for all $1 \leq i \leq n$ **do**

▷ Penalise all mistaken experts

$$w_i \leftarrow \begin{cases} \beta w_i & \text{if } v_{i,t} \neq u_t \\ w_i & \text{otherwise.} \end{cases}$$

Concluding remarks



- This was a **short** intro to the Multiplicative Weights Update Algorithms. Much more to say!
 - Different **predictions** (not only binary)
 - Different **payoffs** (not just 0-1 loss: correct/incorrect)
 - **Randomised** version!
- Discovered/rediscovered in many areas: **learning theory**, **game theory**/economics, **computational geometry**, **convex optimisation**...
- Many (sometimes unexpected) **applications**: online learning/bandits, semidefinite programming, flow algorithms, zero-sum games, algorithmic takes on evolution (!)

Some pointers if you have questions or want to know more about any of those (or connections to some of those topics):

- *The Multiplicative Weights Update Method: a Meta-Algorithm and Applications*. Arora, Hazan, Kale (2012):
<https://theoryofcomputing.org/articles/v008a006/>
- Lecture notes by Daniel Hsu (2017), Chapter 1:
<https://www.cs.columbia.edu/~djhsu/coms6998-f17/notes.pdf>

