

Simple statistical tests

This page demonstrates how to conduct simple statistical tests using **base R**, **rstatix**, and **gtsummary**.

- T-test
- Shapiro-Wilk test
- Wilcoxon rank sum test
- Kruskal-Wallis test
- Chi-squared test
- Correlations between numeric variables

...many other tests can be performed, but we showcase just these common ones and link to further documentation.

Each of the above packages bring certain advantages and disadvantages:

- Use **base R** functions to print a statistical outputs to the R Console
- Use **rstatix** functions to return results in a data frame, or if you want tests to run by group
- Use **gtsummary** if you want to quickly print publication-ready tables

Preparation

Load packages

```
pacman::p_load(  
  rio,          # File import  
  here,         # File locator  
  skimr,        # get overview of data  
  tidyverse,    # data management + ggplot2 graphics,  
  gtsummary,    # summary statistics and tests  
  rstatix,      # statistics  
  corrr,        # correlation analysis for numeric variables  
  janitor,      # adding totals and percents to tables
```

```
flextable      # converting tables to HTML
)
```

Import data

We import the dataset of cases from a simulated Ebola epidemic. If you want to follow along, click to download the “clean” linelist (as .rds file).

```
# import the linelist
linelist <- import("linelist_cleaned.rds")
```

base R

You can use **base** R functions to conduct statistical tests. The commands are relatively simple and results will print to the R Console for simple viewing. However, the outputs are usually lists and so are harder to manipulate if you want to use the results in subsequent operations.

T-tests

A **t-test**, also called “Student’s t-Test”, is typically used to determine if there is a significant difference between the means of some numeric variable between two groups. Here we’ll show the syntax to do this test depending on whether the columns are in the same data frame.

Syntax 1: This is the syntax when your numeric and categorical columns are in the same data frame. Provide the numeric column on the left side of the equation and the categorical column on the right side. Specify the dataset to `data =`. Optionally, set `paired = TRUE`, and `conf.level = (0.95 default)`, and `alternative =` (either “two.sided”, “less”, or “greater”). Enter `?t.test` for more details.

```
## compare mean age by outcome group with a t-test
t.test(age_years ~ gender, data = linelist)
```

Welch Two Sample t-test

data: age_years by gender

t = -21.344, df = 4902.3, p-value < 2.2e-16

alternative hypothesis: true difference in means between group f and group m is not equal to
95 percent confidence interval:

```

-7.571920 -6.297975
sample estimates:
mean in group f mean in group m
      12.60207      19.53701

```

Syntax 2: You can compare two separate numeric vectors using this alternative syntax. For example, if the two columns are in different data sets.

```
t.test(df1$age_years, df2$age_years)
```

You can also use a t-test to determine whether a sample mean is significantly different from some specific value. Here we conduct a one-sample t-test with the known/hypothesized population mean as `mu` =:

```
t.test(linelist$age_years, mu = 45)
```

Shapiro-Wilk test

The [Shapiro-Wilk test](#) can be used to determine whether a sample came from a normally-distributed population (an assumption of many other tests and analysis, such as the t-test). However, this can only be used on a sample between 3 and 5000 observations. For larger samples a [quantile-quantile plot](#) may be helpful.

```
shapiro.test(linelist$age_years)
```

Wilcoxon rank sum test

The Wilcoxon rank sum test, also called the [Mann-Whitney U test](#), is often used to help determine if two numeric samples are from the same distribution when their populations are not normally distributed or have unequal variance.

```
## compare age distribution by outcome group with a wilcox test
wilcox.test(age_years ~ outcome, data = linelist)
```

Wilcoxon rank sum test with continuity correction

```

data:  age_years by outcome
W = 2501868, p-value = 0.8308
alternative hypothesis: true location shift is not equal to 0

```

Kruskal-Wallis test

The [Kruskal-Wallis test](#) is an extension of the Wilcoxon rank sum test that can be used to test for differences in the distribution of more than two samples. When only two samples are used it gives identical results to the Wilcoxon rank sum test.

```
## compare age distribution by outcome group with a kruskal-wallis test  
kruskal.test(age_years ~ outcome, linelist)
```

Kruskal-Wallis rank sum test

data: age_years by outcome

Kruskal-Wallis chi-squared = 0.045675, df = 1, p-value = 0.8308

Chi-squared test

[Pearson's Chi-squared test](#) is used in testing for significant differences between categorical groups.

```
## compare the proportions in each group with a chi-squared test  
chisq.test(linelist$gender, linelist$outcome)
```

Pearson's Chi-squared test with Yates' continuity correction

data: linelist\$gender and linelist\$outcome

X-squared = 0.0011841, df = 1, p-value = 0.9725

Correlations

Correlation between numeric variables can be investigated using the **tidyverse** **corr** package. It allows you to compute correlations using Pearson, Kendall tau or Spearman rho. The package creates a table and also has a function to automatically plot the values.

```
correlation_tab <- linelist %>%  
  select(generation, age, ct_blood, days_onset_hosp, wt_kg, ht_cm) %>% # keep numeric variables  
  correlate() # create correlation table (using default pearson)  
  
correlation_tab # print
```

```
# A tibble: 6 x 7
  term          generation      age ct_blood days_onset_hosp  wt_kg  ht_cm
  <chr>          <dbl>      <dbl>   <dbl>      <dbl>    <dbl>  <dbl>
1 generation      NA      -2.22e-2  0.179      -0.288   -0.0302 -0.00942
2 age            -0.0222    NA      0.00849   -0.000635  0.833   0.877
3 ct_blood        0.179    8.49e-3    NA      -0.600   -0.00636  0.0181
4 days_onset_hosp -0.288   -6.35e-4 -0.600      NA      0.0153 -0.00953
5 wt_kg          -0.0302    8.33e-1 -0.00636    0.0153    NA      0.884
6 ht_cm          -0.00942   8.77e-1  0.0181   -0.00953   0.884    NA
```

```
## remove duplicate entries (the table above is mirrored)
```

```
correlation_tab <- correlation_tab %>%
  shave()
```


```
## view correlation table
```

```
correlation_tab
```

```
# A tibble: 6 x 7
  term          generation      age ct_blood days_onset_hosp  wt_kg  ht_cm
  <chr>          <dbl>      <dbl>   <dbl>      <dbl>    <dbl>  <dbl>
1 generation      NA      NA      NA      NA      NA      NA
2 age            -0.0222    NA      NA      NA      NA      NA
3 ct_blood        0.179    0.00849    NA      NA      NA      NA
4 days_onset_hosp -0.288   -0.000635 -0.600      NA      NA      NA
5 wt_kg          -0.0302    0.833   -0.00636    0.0153    NA      NA
6 ht_cm          -0.00942   0.877    0.0181   -0.00953   0.884    NA
```

```
## plot correlations
```

```
rplot(correlation_tab)
```



Resources

Much of the information in this page is adapted from these resources and vignettes online:

[gtsummary](#) [dplyr](#) [corrr](#) [sthda](#) [correlation](#)