

Research and Applications

Defining dyadic cancer pain concordance using participant-initiated interactions with a remote health monitoring system

Mina Ostovari , PhD^{1,*}, Natalie Crimp , BA¹, Sarah J. Ratcliffe , PhD²,
Virginia LeBaron , PhD, APRN¹

¹School of Nursing, University of Virginia (UVA), Charlottesville, VA 22903, United States, ²School of Medicine, University of Virginia (UVA), Charlottesville, VA 22903, United States

*Corresponding author: Mina Ostovari, PhD, SSIE, 4400 Vestal Pkwy E, Binghamton, NY 13902, United States (mostovari@binghamton.edu)

Abstract

Background: Studies on symptom concordance between patients and their caregivers often use cross-sectional designs, which may fail to capture the longitudinal, dynamic symptom experience. The Behavioral and Environmental Sensing and Intervention for Cancer (BESI-C) is a remote health monitoring system that utilizes smartwatches and ecological momentary assessments (EMAs) to empower patients and caregivers to monitor and manage cancer pain at home. BESI-C collects real-time symptom data in naturalistic settings, enabling longitudinal tracking and analysis of symptom patterns over time.

Objective: To define and examine dyadic concordance using participant-initiated symptom reports collected via remote health monitoring.

Methods: Dyads of patients with advanced cancer and their family caregivers were recruited to use BESI-C for 2 weeks, reporting pain in real time through EMAs. We used Bangdiwala's B statistic to determine the concordance of patient-reported pain and caregiver-reported perceived patient pain under different contextual criteria (eg, co-location of participants; user engagement with BESI-C) that we hypothesized would impact concordance. We also explored a hypothesis that concordance would improve between study week 1 versus week 2.

Results: Data from 21 patient-caregiver dyads were used for analysis. The reporting of pain events was highly variable between patients and their caregivers. Concordance of pain reporting improved when patients and caregivers were co-located and both wearing their BESI-C smartwatches. We did not observe consistent patterns in patient-caregiver concordance between week 1 and week 2.

Conclusion: We propose an analytical approach to define and evaluate concordance between patients' and caregivers' real-time symptom reports that can be applied to dyadic, longitudinal symptom data collected using remote health monitoring. Future work should examine the relationship between patient-caregiver symptom concordance with key quality-of-life metrics and sociodemographic factors that impact participant engagement with remote health monitoring technologies.

Lay Summary

Cancer symptoms are mostly managed at home by patients and their families. Family members and patients may have different perceptions of patient symptoms, and this could impact their communication and symptom management. Previous studies about symptom management often collected data at specific points in time and might have missed how pain levels and perceptions changed day-to-day. Our study used a remote health monitoring system designed by our research team, which uses smartwatches and surveys to allow patients and their caregivers to report on cancer pain in real time, from their homes. Over a 2-week period, 21 patient-caregiver pairs used our system to track patient pain and caregiver's perception of patient pain throughout their daily lives. We found that patients and caregivers' perceptions of pain occurring were more similar when they were physically together and actively using the system. This research introduces a new way to analyze how closely patients and caregivers agree on symptoms over time using real-world data. Our approach may help improve how we support families coping with cancer at home and guide future research into how these dynamics affect quality of life.

Key words: cancer; pain; patient-caregiver dyad; palliative care; symptom management; remote health monitoring; concordance.

Background

Cancer pain is often managed at home by patients and their families.^{1,2} For patients with advanced cancer, family members play a significant role in managing pain, such as administering medications or other pain-alleviating therapies.^{3,4} Managing cancer pain can be challenging and cause emotional distress, anxiety, and depression for both patients and their family caregivers.^{5,6} Implementing a dyadic approach in

pain assessment and management—one that considers a patient and their caregiver as a unit of care and incorporates both perspectives—can encourage collaborative decision-making, improve communication,^{7,8} and optimize health.⁹

Patients and their family caregivers often have different perceptions of a patient's pain which can impact symptom management.^{10–12} Previous studies have assessed concordance between patient-reported and caregiver-reported symptoms

Received: December 27, 2024; Revised: June 18, 2025; Editorial Decision: July 14, 2025; Accepted: July 28, 2025

© The Author(s) 2025. Published by Oxford University Press on behalf of the American Medical Informatics Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

using cross-sectional approaches that may not effectively capture the dynamic symptom experience.^{13–15} To design tailored interventions for both patients and their caregivers, longitudinal studies are needed to understand the dynamic symptom experiences at the dyadic level within the context of everyday life.^{10,16,17} Remote health monitoring systems (RHMS) can enhance the dyadic pain management approach by empowering patients and their caregivers to track symptoms, such as pain, in real-time in their daily lives for a more accurate and longitudinal depiction of their experiences.¹⁸

In this study, we propose an analytical approach to determine and analyze the concordance between patient-reported pain and caregiver-reported perceived patient pain by leveraging real-time data collected using the Behavioral and Environmental Sensing and Intervention for Cancer (BESI-C). BESI-C is an innovative example of RHMS designed to improve cancer pain management in the home setting. BESI-C consists of several components, including smartwatches worn by both patients and caregivers programmed with a custom application to facilitate the reporting of pain events using on-demand ecological momentary assessments (EMAs)—repeated brief assessments in real-time¹⁹ (Figure 1).

Participants mark patient pain as it occurs using an on-demand EMA and then receive an automatically generated follow-up EMA 40 minutes later to assess any changes in the pain or any interventions utilized to reduce the pain. Participants also receive a daily, scheduled EMA that assesses other quality of life factors, such as the amount of time patients and caregivers spent together over the past 24 hours. The smartwatches also collect physiological data (eg, heart rate) and activity data (eg, accelerometer). BESI-C also includes environmental sensors that collect ambient environmental data (eg, temperature) and interact with the smartwatches to determine the location of the patient or caregiver in their home when a pain event is recorded. Details of BESI-C design, feasibility testing, and early pilot results have been previously reported.^{20–23}

This study fills a gap in the literature by examining the longitudinal, temporal concordance between patient-reported pain and caregiver-reported perceived patient pain. The proposed analytical approach can be applied to other symptom data collected using RHMS.

Methods

Study design and participants

Approval was granted by the University of Virginia Health Sciences IRB (#21017), and all participants signed informed consent before data collection. Dyads of patients with

advanced cancer (ie, locally or metastatic cancer, any type, and estimated prognosis of <12 months) and their primary family caregivers were recruited from an outpatient palliative care clinic and a home hospice program. Patients and their caregivers used BESI-C smartwatches to record and characterize cancer-related pain as it occurred over 2 weeks. Participants could use on-demand EMAs to report patient cancer-related pain on their respective BESI-C smartwatch regardless of their location (inside or outside their homes).

We posed 2 primary hypotheses for this study: (1) patient-caregiver concordance of pain reports would increase under different contextual criteria related to engagement with BESI-C and co-location of participants (see “Determining concordance and testing hypotheses” section for specific contextual criteria), and (2) patient-caregiver concordance of pain reports would increase between weeks 1 and 2 as participants became more comfortable using the BESI-C system.

Study variables

We made a series of decisions regarding the operationalization of key variables, such as defining pain events, clustering and overlapping rules, and the concordance time window. We cannot overemphasize the importance of ensuring clarity with variable definitions when working with complex, longitudinal, dyadic RHMS symptom data. We considered patient-reported pain events (ie, on-demand EMAs submitted via the smartwatch application) as the anchor/reference and clustered patient-reported pain events occurred within a 40-minute window. Caregiver-reported perceived patient pain events (EMAs) within a 40-minute window were also clustered.

We chose a 40-minute window as 40 minutes after each initial pain event recorded by the patient/caregiver, the BESI-C application sends a follow-up EMA to re-assess the pain severity and any pharmacological or non-pharmacological actions taken to mitigate the pain. During this 40-minute window, participants can record additional pain events, for example, due to escalating pain. If a pain event was recorded within a 40-minute window of more than one cluster, it was grouped with the first/earlier cluster. We defined an “overlap” as when a caregiver-reported perceived patient pain event was recorded within 30 minutes before or after a patient-reported pain event or cluster. Operational definitions of pain events, clusters, and overlapping events are summarized and illustrated with diagram examples in Table 1.

Table 2 describes the different overlap statuses we considered for patient and caregiver-reported pain events. The







Figure 1. (Left) BESI-C environmental sensors. (Right) Examples of the BESI-C smartwatch EMA interface to record pain events.

Table 1. Definition of pain events, clusters, and overlapping events with diagram examples.

	Definition	Diagram representation
Event	A discrete episode of pain recorded by a patient or caregiver on their respective smartwatch using an on-demand EMA.	
Cluster	Pain events recorded by the patient or caregiver within a 40-minute window. The time window starts from the timestamp of the first pain event and ends 40 minutes later.	
Overlap	Any caregiver-recorded pain event or cluster recorded within a 30-minute window before or after a patient-recorded pain event or cluster. *Note: Patient-recorded pain events are used as the reference/anchor to determine overlap.	

CT, caregiver; EMA, ecological momentary assessment; PT, patient.

overlap status of 1, 2, and 3 were used in a contingency table to calculate the Bangdiwala's B statistic²⁴ for symptom concordance (see Determining concordance and testing hypotheses section).

Determining concordance and testing hypotheses

We determined concordance by calculating: (1) the Bangdiwala's B statistic and (2) the proportion of "pt=yes, cg=yes" (overlap status 3) divided by the total number of cases across overlap statuses 1, 2, and 3, which we termed "consensus" (see Table 2). The Bangdiwala's B statistic is a quantitative measure to assess inter-rater reliability between 2 raters (ie, patient and caregiver),²⁴ which is more suitable for data with unbalanced marginal distributions than Cohen's d.²⁵ B statistic can range from 0 to +1. A value of 0-0.05 corresponds to poor agreement; 0.05-0.15 corresponds to fair; 0.15-0.35 corresponds to moderate; 0.35-0.65 is considered substantial; values ≥ 0.65 show perfect agreement between 2 raters.²⁶

We calculated consensus and B statistic under 3 contextual criteria to investigate our hypothesis that concordance would increase given changes in patient and caregiver co-location and engagement with the system: criterion 1 (C1): without consideration of patient/caregiver locations or smartwatch use; criterion 2 (C2): patient and caregiver both at home, but without consideration of smartwatch use; and criterion 3 (C3): patient and caregiver both at home and wearing study smartwatches. To examine participant location ("at home" or "not at home"), we first checked the self-reported location on the pain EMA and if that was not available, we checked the localization data captured by the interaction between environmental sensors and the smartwatches. We utilized the presence of accelerometer and heart rate data on the smartwatches to confirm patients and caregivers were wearing their smartwatches. To test our second hypothesis, we used Wilcoxon signed-rank test²⁷ to examine if concordance for each dyad improved from study week 1 to week 2 (eg,

comparing consensus [the percentage of pt=yes, cg=yes events] in week 1 versus week 2 from the same dyad). All data analysis was done using R version 4.4.0 (2024-04-24).²⁸

Contextualizing concordance

To further contextualize our concordance results, we examined participants' responses to selected items on a post-deployment survey that inquired whether they recorded pain events independently or in consultation with their partner (note: before the deployment, participants are encouraged to enter data independently so we can best assess their individual experience, but this is not required). We also examined the patient's self-reported Eastern Cooperative Oncology Group (ECOG) Performance Status collected from the baseline pre-deployment survey to understand their physical capability for recording pain events.²⁹

Additionally, we analyzed responses to questions about participant employment status (from the baseline pre-deployment survey) and how much time patients and caregivers reported spending together (from the end-of-day scheduled EMAs) to understand how these factors related to patient and caregiver physical proximity might impact the reporting of pain events and subsequent concordance.

Results

Data from 21 patient-caregiver dyads (42 participants) were collected between February 2022 and October 2024. The average age of patients was 63 (SD= ± 13) and the average age of caregivers was 57 (SD= ± 15). The majority of patients were male (n=11) while the majority of caregivers were female (n=14). Most caregivers were patients' spouses or domestic partners (n=15). Table 3 summarizes the sociodemographic characteristics of the study participants.

Bangdiwala's B statistic between patient and caregiver-reported pain based on contextual criteria 1, 2, and 3 are presented in Table 4. For example, dyad 8, under criterion 1,

Table 2. Overlap statuses used to code pain events and clusters with diagram examples.

Overlap status	Event/Cluster code	Diagram representation
1) Patient pain event or cluster with no caregiver event or cluster 30-minutes before or after.	PT ^a – yes, CG – no (no overlap)	
2) Caregiver pain event or cluster, with no patient event or cluster 30 minutes before or after.	PT – no, CG – yes (no overlap)	
3) Caregiver pain event or cluster 30- minutes before or after a patient event or cluster.	PT – yes, CG – yes (yes overlap)	
4) No patient or caregiver pain event or cluster marked in a 30- minute window	PT – no, CG – no (no overlap)	

^a CG, caregiver; PT, patient. The dashed ovals represent the overlap status being defined.

had 31 patient-reported pain events, 38 caregiver-reported perceived patient pain events, and 9 overlapping events, resulting in 12% consensus (9/78). Under criterion 2, there were 4 patient-reported pain events, 8 caregiver-reported pain events, and 9 overlapping pain events, yielding 43% in consensus (9/21). Under criterion 3, there were 4 patient-reported events, 7 caregiver-reported events, and 9 overlapping events, leading to 45% in consensus (9/20). The consensus range for criterion 1 spans from 0% to 67%, while for criteria 2 and 3, it ranges from 0% to 92%. The B statistic and consensus under criteria 2 and 3 were equal for all dyads except dyads 2, 8, 13, 14, and 20. All dyads had better concordance (higher B statistics and consensus) under criteria 2 and 3 versus criterion 1. Only 4 dyads (dyads 3, 10, 11, and 21) had poor concordance (B statistic < 0.05) under all 3 criteria. Bolded dyad numbers represent dyads with substantial to perfect concordance across all criteria (Table 4).

Figure 2 presents the B statistics across all dyads and how they changed under the 3 criteria (see Methods, Determining concordance and testing hypotheses section). The B statistic values improved (ie, approached 1) under criterion 2 (when both patients and caregivers were wearing their smartwatches) and

further improved under criterion 3 (when patients and caregivers were both at home and wearing their smartwatches.)

The amount of time that the patient and caregiver spent together during the study period is summarized in Table S1 to further contextualize concordance between patient and caregiver symptom reporting. Patient and caregiver “time spent together” is based on their responses to a question in the scheduled end-of-day EMAs (“Time spent with the patient/caregiver?”) and their self-reported employment status. For example, Table S1 distinguishes between caregivers who are working full-time and those who are retired, potentially allowing them to spend more time with the patient. For instance, dyad 19 caregiver and patient were both retired and indicated that they spent more than 4 hours together during most of the study period. Dyad 4 patient and caregiver were both retired and stated that they did not answer the pain questions independently; however, their concordance was fair under criterion 1 and moderate under criteria 2 and 3. Interestingly, we did not observe any differences in concordance of pain reports within dyads where caregivers were employed and those with caregivers who were unemployed/retired, despite the impact we anticipated this would have on pain

Table 3. Sociodemographic characteristics of study participants, overall sample, and by patients and caregivers.

	Total, n (%) 42 (100%)	Patients, n (%) 21 (50%)	Caregivers, n (%) 21 (50%)
<i>Column percentages</i>			
Age band			
20-40	4 (10%)	1 (5%)	3 (14%)
41-60	11 (26%)	5 (24%)	6 (29%)
61-80	27 (64%)	15 (71%)	12 (57%)
Gender			
Female	24 (57%)	10 (48%)	14 (67%)
Male	18 (43%)	11 (52%)	7 (33%)
Race			
Other	1 (2%)	0 (0)	1 (5%)
Black/African American	2 (5%)	1 (5%)	1 (5%)
White	39 (93%)	20 (95%)	19 (90%)
Ethnicity			
Latino/Hispanic	1 (2%)	0 (0)	1 (5%)
Non-Latino/Hispanic	41 (98%)	21 (100%)	20 (95%)
Highest education level			
Less than high school	1 (2%)	0 (0)	1 (5%)
High school graduate	18 (43%)	11 (52%)	7 (33%)
Some college/college degree	18 (43%)	8 (38%)	10 (48%)
Professional/graduate degree	4 (10%)	1 (5%)	3 (14%)
Missing	1 (2%)	1 (5%)	
Current employment			
Employed full-time	9 (21%)	2 (10%)	7 (33%)
Employed part-time	5 (12%)	1 (5%)	4 (19%)
Retired	21 (50%)	13 (61%)	8 (38%)
Other (disabled/unemployed)	7 (17%)	5 (24%)	2 (10%)
Relationship to patient			
Spouse/partner		N/A	15 (71%)
Parent			1 (5%)
Child			3 (14%)
Friend			1 (5%)
Other family			1 (5%)
Primary cancer diagnosis			
Hematological malignancies		5 (24%)	NA
Solid tumors		16 (76%)	
Self-reported ECOG^{a,29}			
I am fully active		1 (5%)	
I can't do heavy work, but I can do some light work		9 (43%)	
I can't do any work, but I can care for myself		6 (28%)	
I need some help caring for myself, and I spend most of the day in bed or in a chair		4 (19%)	
I need much help caring for myself and I spend nearly all day in bed or in a chair.		1 (5%)	

^a Eastern Cooperative Oncology Group, commonly used scale to assess patient's functional status.

reporting (ie, employment status may change the amount of time patient and caregiver spend together in the home, thus impacting likelihood of concordant pain reporting).

Table S2 describes the percentage of overlapping patient and caregiver-reported pain events (see Table 1 for how overlapping pain events were determined) for each dyad in week 1 compared to week 2 of the study period. For example, dyad 2 had 8 overlapping pain events out of 27, 4 of which occurred in week 1 and the other 4 in week 2. The Wilcoxon signed rank test *P*-value was only significant under criterion 2 (*P*-value < .05), suggesting a statistically significant difference in the percentage of pt=yes, cg=yes events between week 1 and week 2, when both patient and caregiver were at home (C2).

Discussion


In this study, we propose an analytical approach to define and analyze the concordance of dyadic pain reporting between patients with advanced cancer and their primary

family caregivers, using real-time, longitudinal data collected through BESI-C, an innovative RHMS that facilitates cancer pain tracking and management in the home setting.²² Our study makes an important contribution to the symptom science literature by offering an approach that can be helpful to others engaged in similar research that utilizes longitudinal, dyadic symptom reporting data. Analytical techniques to analyze complex symptom data are particularly needed as the use of RHMSs become more prevalent.³⁰

Overall, our results demonstrate that the reporting of pain events can be highly variable between patients and their caregivers. As noted in previous studies, family caregivers often experience difficulties in detecting and describing patient symptoms, such as pain, due to their subjective nature.³¹ Caregivers may rely on different sources of information, including their previous knowledge about the patient's typical behavior and expressions of pain, which can lead to discrepancies between patient and caregiver perceptions of the symptom experience?³²

Table 4. B statistics, and percent consensus between patient and caregiver reported pain events, per dyad, for criterion 1, 2, and 3.

Dyad	PT—yes, CG-no (no overlap), status 1			PT—no, CG-yes (no overlap), status 2			PT—yes, CG-yes (overlapping), status 3			Consensus	B	Consensus	B	Consensus	B
	C1 ^a	C2	C3	C1	C2	C3	C1	C2	C3	C1	C1	C2	C2	C3	C3
Dyad 1	28	1	1	2	1	1	1	1	1	3% (1/31)	0.01	33% (1/3)	0.2	33% (1/3)	0.2
Dyad 2	18	7	4	1	0	0	8	5	2	30% (8/27)	0.25	42% (5/12)	0.42	33% (2/6)	0.33
Dyad 3	18	12	12	15	7	7	3	3	3	8% (3/36)	0.01	13% (3/22)	0.04	13% (3/22)	0.04
Dyad 4	70	20	20	23	9	9	26	22	22	21% (26/119)	0.11	43% (22/51)	0.33	43% (22/51)	0.33
Dyad 5	39	27	27	29	22	22	14	13	13	17% (14/82)	0.06	20% (13/62)	0.08	20% (13/62)	0.08
Dyad 6 ^b	5	0	0	3	2	2	12	12	12	60% (12/20)	0.53	86% (12/14)	0.85	86% (12/14)	0.85
Dyad 7 ^b	17	0	0	34	5	5	5	5	5	9% (5/56)	0.017	50% (5/10)	0.5	50% (5/10)	0.5
Dyad 8	31	4	4	38	8	7	9	9	9	12% (9/78)	0.03	43% (9/21)	0.32	45% (9/20)	0.34
Dyad 9	12	11	11	12	5	5	8	8	8	25% (8/32)	0.12	33% (8/24)	0.21	33% (8/24)	0.21
Dyad 10 ^a	4	3	3	21	9	9	0	0	0	0	0	0	0	0	0
Dyad 11	36	24	24	15	13	13	4	4	4	7% (4/55)	0.012	10% (4/41)	0.02	10% (4/41)	0.02
Dyad 12 ^{a,b}	5	1	1	6	0	0	7	7	7	39% (7/18)	0.26	88% (7/8)	0.87	88% (7/8)	0.87
Dyad 13	22	12	12	30	28	28	9	9	8	15% (9/61)	0.04	18% (9/49)	0.07	17% (8/48)	0.06
Dyad 14	55	17	17	6	4	4	9	9	7	13% (9/70)	0.06	30% (9/30)	0.2	25% (7/28)	0.15
Dyad 15	23	17	17	24	21	21	12	11	11	20% (12/59)	0.08	22% (11/49)	0.09	22% (11/49)	0.09
Dyad 16 ^b	12	2	2	0	0	0	24	24	22	67% (24/36)	0.67	92% (24/26)	0.92	92% (22/24)	0.92
Dyad 17 ^{a,b}	8	2	2	3	1	1	5	5	5	31% (5/16)	0.19	62% (5/8)	0.57	62% (5/8)	0.57
Dyad 18	37	32	32	6	6	6	11	11	11	20% (11/54)	0.12	22% (11/49)	0.13	22% (11/49)	0.13
Dyad 19 ^{a,b}	13	5	5	12	5	5	16	10	10	39% (16/41)	0.26	50% (10/20)	0.4	50% (10/20)	0.4
Dyad 20	30	16	16	31	21	21	14	13	12	18% (14/75)	0.06	26% (13/50)	0.13	24% (12/49)	0.11
Dyad 21	49	25	25	10	5	5	3	3	3	4% (3/62)	0.01	9% (3/33)	0.03	9% (3/33)	0.03

^a Dyads recruited from hospice.^b Dyads with substantial or perfect concordance under criteria 2 and 3; Bold dyad numbers represent dyads with substantial to perfect concordance across all criteria. Under C1, we did not consider patient or caregiver location or watch-wearing. Under C2, we only considered pain events that were reported when both patient and caregiver were at home and did not consider watch-wearing. Under C3, we considered pain events reported when both patient and caregiver were at home, and both were wearing their smartwatches.**Figure 2.** B statistic across all dyads based on the 3 contextual criteria described in the Methods section. Under criterion 1 (C1), we did not consider patient or caregiver location or watch-wearing. Under criterion 2 (C2), we only considered pain events that were reported when both patient and caregiver were at home and did not consider watch-wearing. Under criterion 3 (C3), we considered pain events reported when both patient and caregiver were at home, and both were wearing their smartwatches.

We examined concordance for pain events using the Bangdiwala's B²⁴ under 3 criteria to further contextualize the pain experience. Not surprisingly, when we refined our criteria to include only pain events where both the patient and caregiver were at home and wearing their smartwatches (criterion 3), the B statistic values improved (approached one). This suggests that the physical proximity/co-location of both parties and using the monitoring devices might reduce discrepancies (eg, improve concordance) in pain reporting. In other words, being in the same location and consistent engagement with RHMSs (eg, wearing the BESI-C smartwatches) helped ensure both parties were actively engaged in symptom monitoring.

Across all dyads, a third of our sample (10/21; almost 50%) had moderate to perfect concordance under criteria 2 (both at home) and 3 (both at home and wearing smartwatch), and the rest had poor to moderate concordance under the same criteria. We also qualitatively examined the sociodemographic characteristics of dyads (eg, age, gender) with substantial to perfect concordance to identify any specific patterns or traits that set them apart from others. However, we did not find any notable differences. The various symptom experiences across dyads underscore the complexity of pain perception and highlight the need for tailored interventions that can bridge the perceptual gap between patients and caregivers.^{33,34}

We hypothesized that concordance would improve from week 1 to week 2 as participants gained more familiarity with the BESI-C system. We found significant variability in the patient-caregiver consensus (the percentage of pt=yes, cg=yes events) by week; some dyads showed higher consensus in week 1 and others in week 2. Additionally, when we analyzed concordance among dyads with varying amounts of time spent together (as detailed in Table S1), we observed variability in concordance levels. This variability may be attributed to different levels of engagement with the RHMS among participants which may be influenced by sociodemographic variables, an important factor we plan to investigate further in future research.

Limitations

The results presented in this study are based on data collected from patients with advanced cancer and their family caregivers. While this work focuses on a specific disease and symptom (cancer pain), the overall analytic approach is applicable to understanding the concordance of symptom reports from other patient populations and diseases. Researchers using different RHMS or data collection procedures may need to modify the approach to suit their specific contexts. Study data were collected from healthcare institutions in Central Virginia with predominantly White, non-Hispanic populations which limited racial and ethnic diversity of the study sample. The lack of diversity may affect the generalizability of findings with respect to factors such as racial and ethnic differences.³⁵ Future studies should aim to include a more diverse sample to ensure the broader applicability of the findings. To define concordance, we used patient-reported pain events as the anchor/reference. However, if we had considered caregiver-reported perceived patient pain as the reference, the results might have differed. These results reflect an interim analysis of prospectively collected data; results may change once the entire final data set is available and analyzed.

Future directions

Future work should investigate what classifies as “high” or “low” symptom concordance, and how it correlates with patient and caregiver quality-of-life metrics. For example, does being “in sync” with your partner regarding symptom reporting have a positive impact on metrics, such as overall distress levels or mood? In this foundational work using longitudinal, dyadic symptom report data, we sought to answer the question: “what is concordance and how should we quantify it?” Future work will extend this analysis by exploring “how much does concordance matter?” Additionally, we plan to further investigate how patients and caregivers characterized pain events/clusters by exploring differences in reported pain severity and distress levels.

Conclusion

Our study addresses a critical gap in symptom science by proposing an analytical approach to assess concordance between real-time patient and caregiver symptom reports using data from RHMS. Developing standardized methods that are appropriate for complex, longitudinal symptom data collected in real-time is essential for optimizing pain management strategies. Future research should build on this framework to explore additional symptom domains and the broader impact of concordance on the quality of life for both patients and their caregivers.

Acknowledgments

The authors would like to thank Amber Steen, Project Manager, and Penny Amos, Clinical Research Coordinator, for their contributions to this work.

Author contributions

Mina Ostovari (Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing—original draft), Natalie Crimp (Conceptualization, Data curation, Methodology, Visualization, Writing—review & editing), Sarah J. Ratcliffe (Conceptualization, Funding acquisition, Methodology, Supervision, Validation, Writing—review & editing), and Virginia LeBaron (Conceptualization, Funding acquisition, Methodology, Project administration, Supervision, Writing—review & editing)

Supplementary material

[Supplementary material](#) is available at *JAMIA Open* online.

Funding

This work was supported by the National Institute of Nursing Research of the National Institutes of Health under award number R01NR019639. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflicts of interest

The authors have no conflicts of interest to declare.

Data availability

The study data cannot be shared publicly due to privacy and ethical restrictions, as they include sensitive personal health information. Data may be shared upon reasonable requests to the corresponding author, subject to institutional review board approval and a data use agreement.

Sanction

- Are you or any of your co-authors named on the following list relating to trade sanctions: the OFSI Consolidated List, the US Specially Designated Nationals or Blocked Persons list, and/or the EU Consolidated List? No
- Are you or any of your co-authors employed by an institution or entity which is named on the following lists relating to trade sanctions: the OFSI Consolidated List, the US Specially Designated Nationals or Blocked Persons list, and/or the EU Consolidated List? No
- Does your article contain military or defense-related or other controlled technical information which is restricted from publication by applicable export control laws? No

References

1. Chi N-C, Nakad L, Han S, et al. Family caregivers' challenges in cancer pain management for patients receiving palliative care. *Am J Hosp Palliat Care*. 2023;40:43-51.
2. Ullgren H, Tsitsi T, Papastavrou E, Charalambous A. How family caregivers of cancer patients manage symptoms at home: a systematic review. *Int J Nurs Stud*. 2018;85:68-79.
3. Chi N-C, Demiris G. Family caregivers' pain management in end-of-life care: a systematic review. *Am J Hosp Palliat Care*. 2017;34:470-485.
4. Mather H, Kleijwegt H, Bollens-Lund E, Kelley AS, Ornstein KA. Symptom management experience of end-of-life family caregivers: a population-based study. *J Pain Symptom Manage*. 2022;64:513-520.
5. Strang P. Cancer pain—a provoker of emotional, social and existential distress. *Acta Oncol*. 1998;37:641-644.
6. Zaza C, Baine N. Cancer pain and psychosocial factors: a critical review of the literature. *J Pain Symptom Manage*. 2002;24:526-542.
7. McPherson CJ, Hadjistavropoulos T, Lobchuk MM, Kilgour KN. Cancer-related pain in older adults receiving palliative care: patient and family caregiver perspectives on the experience of pain. *Pain Res Manag*. 2013;18:293-300.
8. Hackett J, Godfrey M, Bennett MI. Patient and caregiver perspectives on managing pain in advanced cancer: a qualitative longitudinal study. *Palliat Med*. 2016;30:711-719.
9. Lyons KS, Lee CS. The theory of dyadic illness management. *J Fam Nurs*. 2018;24:8-28.
10. Chen Y, Zhong J, Hong Y, Yuan J, Zhang J. Patient-family caregiver concordance of symptom assessment for esophageal cancer patients undergoing esophagectomy. *Cancer Nurs*. 2024;47:141-150.
11. Horio F, Ikeda T, Arake Y, et al. Consistency between patients and families in recognizing cancer chemotherapy side effects: a questionnaire survey. *Cancer Rep (Hoboken)*. 2022;5:e1451.
12. Yeşilbalkan ÖU, Okgün A. Patients' self reports and caregivers' perception of symptoms in Turkish cancer patients. *Eur J Oncol Nurs*. 2010;14:119-124.
13. Michael N, Gorelik A, Georgousopoulou E, et al. Patient-caregiver communication concordance in cancer—refinement of the cancer communication assessment tool in an Australian sample. *Support Care Cancer*. 2022;30:7387-7396.
14. Raffi F, Taleghani F, Khatooni M. Barriers to effective cancer pain management in home setting: a qualitative study. *Pain Manag Nurs*. 2021;22:531-538.
15. Ma X, Yu W, Lu Y, et al. Congruence of cancer pain experience between patients and family caregivers and associated factors: a multicenter cross-sectional study in China. *Support Care Cancer*. 2021;29:5983-5990.
16. Cui P, Shi J, Li S, et al. Family resilience and its influencing factors among advanced cancer patients and their family caregivers: a multilevel modeling analysis. *BMC Cancer*. 2023;23:623.
17. Lyons KS, Lee CS. The association of dyadic symptom appraisal with physical and mental health over time in care dyads living with lung cancer. *J Fam Nurs*. 2020;26:15-25.
18. McGregor BA, Vidal GA, Shah SA, Mitchell JD, Hendifar AE. Remote oncology care: review of current technology and future directions. *Cureus*. 2020;12:e10156.
19. Shiffman S, Stone AA, Hufford MR. Ecological momentary assessment. *Annu Rev Clin Psychol*. 2008;4:1-32.
20. LeBaron V, Hayes J, Gordon K, et al. Leveraging smart health technology to empower patients and family caregivers in managing cancer pain: protocol for a feasibility study. *JMIR Res Protoc*. 2019;8:e16178.
21. LeBaron V, Bennett R, Alam R, et al. Understanding the experience of cancer pain from the perspective of patients and family caregivers to inform design of an in-home smart health system: multimethod approach. *JMIR Form Res*. 2020;4:e20836.
22. LeBaron V, Alam R, Bennett R, et al. Deploying the behavioral and environmental sensing and intervention for cancer smart health system to support patients and family caregivers in managing pain: feasibility and acceptability study. *JMIR Cancer*. 2022;8:e36879.
23. LeBaron V, Homdee N, Ogunjirin E, et al. Describing and visualizing the patient and caregiver experience of cancer pain in the home context using ecological momentary assessments. *Digit Health*. 2023;9:20552076231194936-20552076231194936.
24. Bangdiwala SI. A graphical test for observer agreement. In: *Proceedings of the 45th International Statistical Institute Meeting*. Amsterdam: International Statistical Institute. Vol. 1. 1985:307-308.
25. Shankar V, Bangdiwala SI. Behavior of agreement measures in the presence of zero cells and biased marginal distributions. *J Appl Stat*. 2008;35:445-464.
26. Munoz SR, Bangdiwala SI. Interpretation of kappa and B statistics measures of agreement. *J Appl Stat*. 1997;24:105-112.
27. Holmes DT. Chapter 2—Statistical methods in laboratory medicine. In: Clarke W, Marzinke MA, eds. *Contemporary Practice in Clinical Chemistry*. 4th ed. Academic Press; 2020:15-35. <https://doi.org/10.1016/B978-0-12-815499-1.00002-8>
28. R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing; 2013.
29. Young J, Badgery-Parker T, Dobbins T, et al. Comparison of ECOG/WHO performance status and ASA score as a measure of functional status. *J Pain Symptom Manage*. 2015;49:258-264.
30. Vitabile S, et al. Medical data processing and analysis for remote health and activities monitoring. In: Kołodziej J, González-Vélez H, eds. *High-Performance Modelling and Simulation for Big Data Applications: Selected Results of the COST Action IC1406 cHiP-Set*. Cham: Springer International Publishing, 2019:186-220. https://doi.org/10.1007/978-3-030-16272-6_7
31. McPherson CJ, Addington-Hall JM. Evaluating palliative care: bereaved family members' evaluations of patients' pain, anxiety and depression. *J Pain Symptom Manage*. 2004;28:104-114.
32. McPherson CJ, Wilson KG, Lobchuk MM, Brajtman S. Family caregivers' assessment of symptoms in patients with advanced cancer: concordance with patients and factors affecting accuracy. *J Pain Symptom Manage*. 2008;35:70-82.

33. Chi N-C, Fu Y-K, Nakad L, et al. Family caregiver challenges in pain management for patients with advanced illnesses: a systematic review. *J Palliat Med*. 2022;25:1865-1876.
34. Simblett S, Greer B, Matcham F, et al. Barriers to and facilitators of engagement with remote measurement technology for managing health: systematic review and content analysis of findings. *J Med Internet Res*. 2018;20:e10480.
35. Krukowski RA, Ross KM, Western MJ, et al. Digital health interventions for all? Examining inclusivity across all stages of the digital health intervention research process. *Trials*. 2024;25:98.