

**Question 1:****a)**

Product rule, for

$$f(x) = g(x)h(x)$$

$$f'(x) = g(x)h'(x) + h(x)g'(x)$$

$$f(x) = (x^5 + 3x^3 + 2x + 1)e^x$$

Using

$$g = x^5 + 3x^3 + 2x + 1$$

and

$$h = e^x$$

Differentiating

$$g' = 5x^4 + 9x^2 + 2$$

and

$$h' = e^x$$



3/3

Using the product rule

$$f'(x) = (5x^4 + 9x^2 + 2)e^x + e^x(x^5 + 3x^3 + 2x + 1)$$



Simplifying

$$= (x^5 + 5x^4 + 3x^3 + 9x^2 + 2x + 3)e^x$$

**b)**

Chain rule, for

$$g(y) = i(h(y))$$

$$g'(y) = i'(h(y))h'(y)$$

$$g(y) = (\ln(y) + \sin(y))^6$$

Using

$$h(y) = \ln(y) + \sin(y)$$

and

$$i(h) = h^6$$

Differentiating

$$h'(y) = \left(\frac{1}{y} + \cos y\right)$$

and

$$i'(h) = 6h^5$$

3/3

Using the chain rule

$$g'(y) = 6 [\ln(y) + \sin(y)]^5 \left( \frac{1}{y} + \cos(y) \right)$$

c)

Quotient rule, for

$$h(z) = \frac{e^{5z}}{(2 + \cos(10z))}$$

$$h(z) = \frac{i(z)}{j(z)}$$

$$h'(z) = \frac{j(z)i'(z) - i(z)j'(z)}{(j(z))^2}$$

Using

$$i(z) = e^{5z}$$

and

$$j(z) = (2 + \cos(10z))$$

3/3

Differentiating

$$i'(z) = 5e^{5z}$$

and

$$j'(z) = -10 \sin(10z)$$

Using the quotient rule

$$h'(z) = \frac{(2 + \cos(10z))5e^{5z} - e^{5z}(-10 \sin(10z))}{(2 + \cos(10z))^2}$$

Simplifying

$$= \frac{5e^{5z}((2 + \cos(10z)) + (2 \sin(10z)))}{(2 + \cos(10z))^2}$$

d)

$$k(x) = x^2 \sin(\cos x)$$

Using

$$l(x) = x^2 \quad \text{and} \quad m(x) = \sin(\cos x)$$

Differentiating using the product rule

$$l'(x) = 2x \quad \checkmark$$

Now using the chain rule to find  $m'(x)$ , using  $u = \sin(x)$  and  $v = \cos(x)$

$$u' = \cos(x)$$

and

$$v' = -\sin(x) \quad \checkmark$$

Thus

$$m' = \cos(\cos x) (-\sin(x)) \quad \checkmark \quad \checkmark$$

Applying the product rule

$$k'(x) = x^2 (\cos(\cos x) (-\sin(x))) + 2x \sin(\cos x) \quad \checkmark$$

Simplifying

$$= 2x \sin(\cos x) - x^2 \sin(x) \cos(\cos x) \quad \checkmark$$

Product rule:

$$k(x) = l(x)m(x)$$

$$k'(x) = l(x)m'(x) + l'(x)m(x)$$

Chain rule


$$m'(x) = u'(v(x))v'(x)$$

6/6

Q1 15/15

**Question 2:**

Given the L-shaped enclosure



a)

2/2

Using the assumption that Steven uses all the fencing he has exactly the perimeter is 74 m

$$\text{perimeter} = (x + 5) + y + 5 + y + x + 2y$$

$$74 = 4y + 2x + 10$$

$$64 = 4y + 2x$$



$$4y = 64 - 2x$$


$$y = 16 - \frac{x}{2}$$



$$= \frac{1}{2} (32 - x)$$

as required

b)



The area of the L-shape is given by the total of the two shapes shown above.

$$\begin{aligned}
 A &= x(2y) + 5(y) \quad \checkmark \\
 &= x \left( 2 \left( \frac{1}{2} (32 - x) \right) \right) + 5 \left( \frac{1}{2} (32 - x) \right) \\
 &= x(32 - x) + \left( 80 - \frac{5x}{2} \right) \\
 &= 32x - x^2 + 80 - \frac{5x}{2} \quad \checkmark
 \end{aligned}$$

Multiply by 2

$$2A = 160 + 64x - 5x - 2x^2 \quad 2/2$$

Collect like terms


$$= 160 + 59x - 2x^2$$

simplify

$$A = \frac{1}{2} (160 + 59x - 2x^2)$$

*as required*

c)



Based on the shape of the curve for this graph we need only consider the stationary point at which  $dA/dx = 0$  to find the maximum area.

Given

$$A = \frac{1}{2}(160 + 59x - 2x^2)$$

Differentiating

$$A' = -2x + \frac{59}{2} \quad \checkmark$$

Setting  $A' = 0$  to find the stationary point

$$0 = -2x + \frac{59}{2} \quad \checkmark$$

$$2x = \frac{59}{2}$$

$$x = \frac{59}{4} \quad \checkmark$$

Substituting this into the original equation

$$\begin{aligned}
 A &= \frac{1}{2} (160 + 59x - 2x^2) \\
 &= \frac{1}{2} \left( 160 + 59 \left( \frac{59}{4} \right) - 2 \left( \frac{59}{4} \right)^2 \right) \\
 &= 80 + \frac{3481}{8} - \left( \frac{59}{4} \right)^2 \\
 &= 80 + \frac{3481}{8} - \left( \frac{3481}{16} \right) \\
 &= \frac{1280}{16} + \frac{6962}{16} - \left( \frac{3481}{16} \right) \\
 &= \frac{1280}{16} + \frac{3481}{16} \\
 &= \frac{4761}{16} \quad \checkmark
 \end{aligned}$$

6/6

Applying the second derivative test

$$\begin{aligned}
 A' &= -2x + \frac{59}{2} \\
 A'' &= -2 \quad \checkmark
 \end{aligned}$$

Showing that this is a maximum stationary point



Hence the maximum area of the L-shape is

$$A = \frac{4761}{16} \text{m}^2$$

Q2 10/10

**Question 3:****a)**

$$f(x) = x^2 + 2x + 5$$

Then the indefinite integral is

2/2

$$F(x) = \frac{x^3}{3} + x^2 + 5x + c \quad \checkmark \quad \checkmark$$

**b)**

$$g(\theta) = 5e^\theta + \frac{1}{5\theta} \quad \checkmark$$

Then the indefinite integral is

1.5/2 Careful here as ln theta should be ln |theta|.

$$G(\theta) = 5e^\theta + \frac{\ln \theta}{5} + c \quad \checkmark$$

**c)**

$$h(t) = 2 \sin(t) + \frac{1}{3 + 3t^2} + 3$$

3/3

$$= 2 \left( \int \sin(t) \right) dt + \frac{1}{3} \left( \int \frac{1}{1 + t^2} \right) dt + 3 \quad \checkmark$$

Then the indefinite integral is

$$H(t) = -2 \cos(t) + \frac{1}{3} \tan^{-1}(t) + 3t + c \quad \checkmark$$

$$= \frac{1}{3} (\tan^{-1}(t) - 6 \cos(t) + 9t + c) \quad \checkmark$$



d)

$$j(y) = (y - 2) \left( y^{\frac{-1}{2}} \right) + 3$$

Expand the brackets

$$= y^{\frac{1}{2}} + 3y - 2y^{\frac{-1}{2}} - 6 \quad \checkmark$$

$$= y^{\frac{1}{2}} + \left( 3 \int y \right) dy - \left( 2 \int y^{\frac{-1}{2}} \right) dy - 6$$

3/3 I have condoned the use of the letter 't' here but do be careful with writing your variables

Then the indefinite integral is

$$J(y) = \frac{1}{3} y^{\frac{3}{2}} + 3 \left( \frac{1}{2} y^2 \right) - 2 \left( \frac{1}{\frac{1}{2}} y^{\frac{1}{2}} \right) - 6y + c$$

$$= \frac{2t^{\frac{3}{2}}}{3} + \frac{3y^2}{2} - 4\sqrt{y} - 6y + c \quad \checkmark \quad \checkmark$$

Q3 10/10

**Question 4:**

$$f(x) = -x^2 + 4x + 12$$

**a)**

As the function is an inverted U parabola the x-intersection points will show where the curve crosses to below the x-axis.

$$f(x) = -x^2 + 4x + 12$$

Substituting both  $-2$  and  $6$  into the equation

$$f(-2) = -(-2)^2 + 4(-2) + 12$$

$$= -4 - 8 + 12$$

$$= 0$$

and

$$f(6) = -(6)^2 + 4(6) + 12$$

$$= -36 + 24 + 12$$



1/1

$$= 0$$

Hence the graph between and not including these points are above the x-axis.

**b)**

$$f(x) = -x^2 + 4x + 12$$

$$= \int_1^3 (-x^2 + 4x + 12) dx$$



1/1 Just the definite integral was required here

$$= \left( -\int x^2 + 4 \int x + 12 \int 1 \right) dx$$

$$= \left( \frac{-1}{3} x^3 + 4 \left( \frac{1}{2} x^2 \right) + 12(x) \right)$$

$$= \left[ \frac{-1}{3} x^3 + 8x^2 + 12x \right]_1^3$$

c)

Using this to find the area under the curve between  $-2 < x < 6$ 

$$f(x) = -x^2 + 4x + 12$$

$$= \int_{-2}^6 (-x^2 + 4x + 12) dx \quad \checkmark$$

1/3 The area to the part b) was required, ie between when x = 3 and 1

$$= \left[ \frac{-1}{3}x^3 + 4\left(\frac{1}{2}\right)x^2 + 12x \right]_{-2}^6 \quad \times$$

$$= \left( \frac{-1}{3}6^3 + 2(6)^2 + 12(6) \right) - \left( \frac{-1}{3}(-2)^3 + 2(-2)^2 + 12(-2) \right)$$

$$= \left( \frac{-1}{3}(216) + (2)36 + 72 \right) - \left( \frac{-1}{3}(-8) + (2)4 - 24 \right)$$

$$= (-72 + 72 + 72) - \left( \frac{8}{3} + 8 - 24 \right)$$

$$= 72 - \left( \frac{-40}{3} \right)$$

Hence the area under the curve between  $x = -2$  and  $x = 6$  is

Q4 3/5

$$= \frac{256}{3} \quad \times$$

The required integral is

$$\begin{aligned} \int_1^3 (-x^2 + 4x + 12) dx &= \left[ -\frac{1}{3}x^3 + 2x^2 + 12x \right]_1^3 \\ &= (-9 + 18 + 36) - \left( -\frac{1}{3} + 2 + 12 \right) \\ &= 45 - \frac{41}{3} \\ &= \frac{94}{3}. \end{aligned}$$

The area is therefore  $\frac{94}{3}$ .*References*

Activity 11 on page 140 of Book C (Unit 8).

Standard indefinite integrals, *Handbook* page 7.

**Question 5:**

a)

$$\int \frac{\cos(3x) - \sin(3x)}{(\sin(3x) + \cos(3x))^2}$$

substitute  $u = 3x$  and  $du = 3 dx$ 

$$= \frac{1}{3} \int \frac{\cos(u) - \sin(u)}{(\sin(u) + \cos(u))^2} du$$

4/4

Substitute  $v = \sin(u) + \cos(u)$ ,  $dv = \cos(u) - \sin(u) du$ 

$$= \frac{1}{3} \int \frac{1}{v^2} dv$$

$$= \frac{1}{3} \left( -\frac{1}{v} \right) + C$$

Substituting  $v$  back in

$$= \frac{-1}{3(\sin u + \cos u)} + C$$

Substituting  $u$  back in

$$= \frac{-1}{3(\sin 3x + \cos 3x)} + C$$



b)

$$\int_0^{\frac{1}{3} \ln 5} e^{3x} \sqrt{e^{3x} + 2} \, dx$$

Substitute  $u = 3x$ ,  $du = 3 \, dx$ 

$$= \frac{1}{3} \int_0^{\ln 5} e^u \sqrt{e^u + 2} \, du$$

Substitute  $v = e^u + 2$ ,  $dv = e^u \, du$ 

$$= \frac{1}{3} \int_0^{\ln 5} \sqrt{v} \, dv$$

The integrand of  $v$  is  $\frac{2}{3} v^{\frac{3}{2}}$ 

$$\begin{aligned} &= \frac{1}{3} \left( \frac{2}{3} v^{\frac{3}{2}} \right) \\ &= \frac{2}{9} v^{\frac{3}{2}} \end{aligned}$$

Substitute  $v$  back in

$$= \frac{2}{9} (e^u + 2)^{\frac{3}{2}}$$

Substitute  $u$  back in

$$= \frac{2}{9} (e^{3x} + 2)^{\frac{3}{2}}$$

It follows that

$$\begin{aligned}\int_0^{\frac{1}{3} \ln 5} e^{3x} \sqrt{e^{3x} + 2} \, dx &= \left[ \frac{2}{9} (e^{3x} + 2)^{\frac{3}{2}} \right]_0^{\frac{1}{3} \ln 5} \quad \checkmark \\&= \left[ \frac{2}{9} \left( e^{3(\frac{1}{3} \ln 5)} + 2 \right)^{\frac{3}{2}} \right] - \left[ \frac{2}{9} \left( e^{3(0)} + 2 \right)^{\frac{3}{2}} \right] \\&= \left[ \frac{2}{9} (5 + 2)^{\frac{3}{2}} \right] - \left[ \frac{2}{9} (1 + 2)^{\frac{3}{2}} \right] \quad \checkmark \\&= 4.115 \dots - 1.154 \dots \\&= 2.96 \quad \checkmark\end{aligned}$$

to 2 d.p

6/6 GMC alert!  
Remember to give full answers  
before rounding, ie  
2.9609.....  
2.96 (2dp)

Q5 10/10

**Question 6:**

Integration py parts

**a)**

$$\int f(x)g(x) dx = f(x)G(x) - \int f'G(x) dx$$

$$\int 81x^8 \ln(x) dx$$

Let,  $f(x) = \ln(x)$  and  $g(x) = x^8$ Then,  $f'(x) = \frac{1}{x}$  and  $G(x) = \frac{x^9}{9}$ 

✓

$$= 81 \int x^8 \ln(x) dx$$

$$= 81 \left[ \ln(x) \frac{x^9}{9} - \int \left( \frac{x^9}{9x} \right) dx \right]$$

$$= 9 \left( \ln(x) x^9 - \int x^8 dx \right)$$

$$= 9 \left( \ln(x) x^9 - \frac{x^9}{9} \right)$$

$$= 9 \ln(x) x^9 - x^9$$

$$= 9x^9 (\ln(x) - 1)$$

✓

4/4

✓

✓

**b)**

$$\int e^{3y} \sin(2y) dy =$$

Let,  $f(y) = \sin(2y)$  and  $g(y) = e^{3y}$ 

✓

Then,  $f'(y) = 2 \cos(2y)$  and  $G(y) = \frac{e^{3y}}{3}$ 

$$= \frac{1}{3} e^{3y} \sin(2y) - \frac{2}{3} \int e^{3y} \cos(2y) dy$$

✓

Let,  $h(y) = \cos(2y)$  and  $i(y) = e^{3y}$

Then,  $h'(y) = -2 \sin(2y)$  and  $I(y) = \frac{e^{3y}}{3}$  ✓

$$= \frac{1}{3} e^{3y} \sin 2y - \frac{2}{3} \left[ \frac{e^{3y} \cos(2y)}{3} - \frac{2}{3} \int e^{3y} \sin(2y) dy \right] \quad \checkmark$$

$$= \frac{1}{3} e^{3y} \sin(2y) - \frac{2}{9} e^{3y} \cos(2y) - \frac{4}{9} \int e^{3y} \sin(2y) dy \quad \checkmark$$

Add  $\frac{4}{9} \int e^{3y} \sin(2y) dy$  to both sides

$$\frac{13}{9} \int e^{3y} \sin(2y) dy = \frac{1}{3} e^{3y} \sin 2y - \frac{2}{9} e^{3y} \cos(2y)$$

6/6

Multiply both sides by  $\frac{9}{13}$

$$\int e^{3y} \sin(2y) dy = \frac{9}{13} \left[ \frac{1}{3} e^{3y} \sin(2y) - \frac{2}{9} e^{3y} \cos(2y) \right]$$

$$= \frac{3}{13} e^{3y} \sin(2y) - \frac{2}{13} e^{3y} \cos(2y)$$

$$= \frac{e^{3y}}{13} (3 \sin(2y) - 2 \cos(2y)) \quad \checkmark$$

Q6 10/10

Type text here



**Question 7:**

Given the function

(% i1)  $f(x) := (3x + 15x^2 - x^4) / (9x^2 + 1);$

(% o1)  $f(x) := \frac{3x + 15x^2 - x^4}{9x^2 + 1}$


(a) The graph of  $f(x)$  is

(% i2) `wxplot2d(f(x), [x, -5, 5], [y, -1, 2]);`

*plot2d : some values will be clipped.*

3/3

(% t2)



(% o2)

(b) the derivative of  $f(x)$  is

(% i3) `df(x) := "(diff(f(x), x));"`



(% o3)  $df(x) := \frac{-(4x^3) + 30x + 3}{9x^2 + 1} - \frac{18x(-x^4 + 15x^2 + 3x)}{(9x^2 + 1)^2}$

1/1

(c) The positive local maximum of  $f(x)$  is

(% i4) pos\_root:find\_root(df(x), x, 0, 5);

(pos\_root) 0.8288158368533624



Substituting this back into the  $f(x)$

(% i5) f(pos\_root);

(% o5) 1.71510439942526



Finding the second derivative of  $f(x)$

(% i6) ddf(x):="(diff(df(x), x));

(% o6)

$$ddf(x) := \frac{30 - 12x^2}{9x^2 + 1} - \frac{18(-x^4 + 15x^2 + 3x)}{(9x^2 + 1)^2} + \frac{648x^2(-x^4 + 15x^2 + 3x)}{(9x^2 + 1)^3} - \frac{36x(-4x^3 + 30x + 3)}{(9x^2 + 1)^2}$$

And substituting our x variable

(% i7) ddf(0.829);

(% o7) -1.2681397331452517



As this is  $<0$  the point is confirmed to be a local maximum of  $f(x)$  Hence the local maximum is at (0.828, 1.715), to 3 d.p. (d) To find the root to the right of  $x=0$

2.5/3 To 3dp the x coordinate is 0.829

(% i8) float(realroots(f(x)));

(% o8)

$[x = -3.7688187062740326, x = -0.20053765177726746, x = 3.9693563878536224, x = 0.0]$

Therefore the graph crosses the x-axis at 3.969, to 3 d.p. The area under the graph between  $x=0$  and this point is

3/3

(% i10) quad\_qags(f(x), x, 0, 3.969);



(% o10)  $[4.342959640603124, 2.562177785637564210^{-9}, 105, 0]$

Hence the area enclosed by the graph of  $f(x)$  between  $0 \leq x \leq 3.969$  is 4.343, to 3 d.p.



**Question 8:****a)****i.**

	Not at all confident	Slightly confident	Somewhat confident	Fairly confident	Very confident
Unit 1					✓
Unit 2					✓
Unit 3			✓		
Unit 4				✓	
Unit 5					✓
Unit 6				✓	
Unit 7			✓		
Unit 8				✓	

1/1

✓

**ii.**

I have a different room as my study, so I am separated from the rest of the house and all the distractions that comes with it. I like to set out short 30 minute time slots with a 15 minute break over the course of a few hours.

I will need to work on the different methods of integration and Taylor polynomials.

✓

✓

2/2 Do also consider attending tutorials and practicing using past papers

**b)****i.**

Section 1  $2\% \times 25 = 0.5 \times 180 = 90$

Section 2  $3\% \times 10 = 0.3 \times 180 = 54$

Section 3  $4\% \times 5 = 0.2 \times 180 = 36$

For section A, I should be averaging about 3.6 minutes per question,

For section B, I should be averaging about 5.4 minutes per question, For section C, I should be averaging about 7.2 minutes per question.

✓

1/1

**ii.**

- Review the material for the sections I am least confident in.
- Some questions might take longer than others, so I should not spend too long on any one question.
- If I am struggling with a question, I should move on and come back to it later.
- Keep track of the questions I do quickly, so I know how much I can spend on harder ones


✓

1/1


Q8 5/5


**Question 9:***Section A*

**Question 1:** A  0/2 correct option is B


**Question 2:** B  2/2

**Question 3:** C  2/2

**Question 4:** D  2/2


**Question 5:** E  2/2

*Section B*

**Question 6:** F  3/3

**Question 7:**

$$f'(x) = 9x^2 - 4$$

The x-coordinates of one stationary point is at  $x = \frac{2}{3}$ . It is a **local minimum**. The x-coordinates of the other stationary point is at  $x = -\frac{2}{3}$ . It is a **local maximum**.  3/3

*Section C***Question 8:**

- A.  $\frac{1}{8}$
- B.  $\frac{1}{2}$   4/4
- C.  $\frac{3}{8}$

Q9 18/20

Q10 5/5 Very good use of GMC throughout, just be careful with accuracy (see Q5)