

1300 Centre Street Boston, MA, 20131

Dear Dr. Gibson:

Please consider our manuscript, 'False spring damage to temperate tree saplings is amplified with winter warming', as a Research Article for the *Journal of Ecology*.

Biological spring is advancing with climate change but late spring freeze dates are not predicted to advance at the same rate, leading to renewed interest in late spring freeze events—commonly called 'false springs'—which shape the life history of many temperate and boreal plant species. Additionally, over-winter chilling is predicted to decrease with warming winters, which could impact plant phenology and, ultimately, growth.

Here, we present an experiment on the interplay of false springs and warmer winters (generally expected to reduce chilling) across eight temperate deciduous tree species examining a suite of phenological, growth and leaf tissue traits. We found that false springs slowed budburst to leafout timing—extending the period of maximum freezing risk, increased damage to the shoot apical meristem, and decreased leaf toughness, leaf thickness and chlorophyll content. Chilling accelerated budburst and leafout, even under false spring conditions, thus chilling compensated for the adverse phenological effects of false springs. Despite major shifts in phenology from false springs and chilling we did not find evidence of phenological reordering within a community. Our results instead suggest climate change will reshape forest communities through impacts on growth and leaf traits from the coupled effects of false springs and decreased over-winter chilling under future climate change.

We believe our research is important and timely because our findings contribute to fundamental knowledge on how chilling and freeze events shape tree growth across species, and suggest climate change could diminish or reverse the positive effects of carbon storage in temperate forests. Damage from false spring events in conjunction with decreases in overwinter chilling may have cascading effects to pollinators, nutrient cycling and carbon uptake as well as forest recruitment.

This manuscript is not under consideration elsewhere. Both authors approved of this version for submission. We hope that you will find it suitable for publication in the *Journal of Ecology*. Thank you for your consideration.

Sincerely,

Cot Cah

Catherine Chamberlain (on behalf of my co-authors)

Authors:

C. J. Chamberlain 1,2 & E. M. Wolkovich 1,2,3

Author affiliations:

¹Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA; ²Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, USA;

³Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4

*Corresponding author: 248.953.0189; cchamberlain@g.harvard.edu

References

- [1] Polgar, C., Gallinat, A., and Primack, R. B. Drivers of leaf-out phenology and their implications for species invasions: Insights from Thoreau's Concord. *New Phytologist* **202**(1), 106–115 (2014).
- [2] Fu, Y. H., Zhao, H., Piao, S., Peaucelle, M., Peng, S., Zhou, G., Ciais, P., Huang, M., Menzel, A., Peñuelas, J., and et al. Declining global warming effects on the phenology of spring leaf unfolding. *Nature* 526(7571), 104–107, Sep (2015).
- [3] Labe, Z., Ault, T., and Zurita-Milla, R. Identifying anomalously early spring onsets in the CESM large ensemble project. *Climate Dynamics* **48**(11-12), 3949–3966, Aug (2016).
- [4] Sgubin, G., Swingedouw, D., Dayon, G., de Cortázar-Atauri, I. G., Ollat, N., Pagé, C., and van Leeuwen, C. The risk of tardive frost damage in french vineyards in a changing climate. *Agricultural and Forest Meteorology* **250-251**, 226 242 (2018).
- [5] Boggs, C. L. and Inouye, D. W. A single climate driver has direct and indirect effects on insect population dynamics. *Ecology Letters* 15(5), 502–508, Mar (2012).
- [6] Pardee, G. L., Inouye, D. W., and Irwin, R. E. Direct and indirect effects of episodic frost on plant growth and reproduction in subalpine wildflowers. *Global Change Biology* 24(2), 848–857, Sep (2017).
- [7] Hufkens, K., Friedl, M. A., Keenan, T. F., Sonnentag, O., Bailey, A., O'Keefe, J., and Richardson, A. D. Ecological impacts of a widespread frost event following early spring leaf-out. *Global Change Biology* **18**(7), 2365–2377 (2012).
- [8] Klosterman, S., Hufkens, K., and Richardson, A. D. Later springs green-up faster: the relation between onset and completion of green-up in deciduous forests of North America. *International Journal of Biometeorology* **62**, 1645–1655 May (2018).
- [9] Richardson, A. D., Keenan, T. F., Migliavacca, M., Ryu, Y., Sonnentag, O., and Toomey, M. Climate change, phenology, and phenological control of vegetation feedbacks to the climate system. Agricultural and Forest Meteorology 169, 156 – 173 (2013).