

## Frost resistance and ice nucleation in leaves of five woody timberline species measured in situ during shoot expansion

D. TASCHLER,<sup>1,2</sup> B. BEIKIRCHER<sup>1</sup> and G. NEUNER<sup>1</sup>

<sup>1</sup> Institute of Botany, University of Innsbruck, Sternwartestrasse 15, 6020 Innsbruck, Austria

<sup>2</sup> Corresponding author (Dunja.Taschler@uibk.ac.at)

Received April 9, 2003; accepted August 3, 2003; published online January 2, 2004

**Summary** Frost resistance and ice nucleation temperatures of leaves, from bud swelling until after full expansion, were measured in situ for five major woody timberline species with recently developed field freezing equipment. Frost resistance determined in situ on leaves of attached twigs was significantly higher than values determined on detached leaves in laboratory tests (e.g., the temperature at which incipient frost damage was observed ( $LT_i$ ) was 1.2 °C higher for detached leaves than for attached leaves of *Picea abies* (L.) Karst.). Frost resistance of leaves of all species changed significantly during shoot expansion (e.g., changes of 7.2 and 11 °C for *Rhododendron ferrugineum* L. and *Larix decidua* Mill., respectively). Expanding leaves (between 0 and 60% of full expansion) were the most sensitive to frost, with  $LT_i$  values ranging from –3.4 °C in *R. ferrugineum* to –6.3 °C in *L. decidua*. Among the studied species, *P. abies* and *R. ferrugineum* were the most frost sensitive throughout the shoot elongation period. In situ freezing patterns of leaves of attached twigs also differed from those of leaves of excised twigs. During leaf expansion, two distinct freezing exotherms were always registered in situ. The first freezing event (E1, high-temperature exotherm) was recorded at  $-1.5 \pm 0.2$  °C and reflected extracellular ice formation. Exposure of leaves to temperatures at which E1 occurred was, in all cases, noninjurious. The low-temperature exotherm (E2) mostly coincided with frost damage, except for some stages of leaf expansion in *R. ferrugineum* and *P. abies*, indicating that in situ freezing exotherms were not accurate estimators of frost damage in these species.

**Keywords:** field freezing equipment, freezing exotherms, *Larix decidua*, *Picea abies*, *Pinus cembra*, *Rhododendron ferrugineum*, *Sorbus aucuparia*.

### Introduction

Low temperature is an important environmental constraint on plant distribution (cf. Sakai and Larcher 1987). At the timberline ecotone, freezing stress affects the development and growth of adult conifers (Tranquillini 1979, Sakai and Larcher 1987, Gross et al. 1991) and particularly contributes to the distorted growth of timberline trees (Tranquillini 1979, Larcher 1985). Although frost resistance in timberline trees is gener-

ally sufficient during winter (Tranquillini 1979), frost resistance decreases in spring to values that are insufficient to protect newly developed tissues against episodic frosts in late spring and summer (Kerner 1869, Matuszkiewicz 1977, Christersson et al. 1987). Natural frost damage to new leaves of *Picea abies* (L.) Karst., *Rhododendron ferrugineum* L., and a range of other subalpine woody species has been repeatedly observed in June at timberline on Mt. Patscherkofel, near Innsbruck (Kerner 1869), including after night frosts of –2.6 °C in 1999 and –4.3 °C in 2001 (D. Taschler and G. Neuner, unpublished observations). Damage caused by late episodic frost events could be aggravated by climate warming. Rising global mean temperatures can induce earlier bud break (Murray et al. 1989, Hänninen 1991, 1996), and climate warming is also anticipated to increase the frequency of unpredictable, episodic frost events (Katz and Brown 1992).

Although winter frost resistance of leaves of timberline conifers has been extensively studied (cf. Bannister and Neuner 2000), much less is known about the summer frost resistance of young leaves during the period from bud swelling until full leaf expansion. New leaves of woody timberline species tolerate frosts between –2 and –10 °C (cf. Tranquillini 1979, Sakai and Larcher 1987).

In most studies of summer frost resistance, little distinction has been made between different developmental stages during shoot expansion, and investigations have been carried out in the laboratory. Laboratory investigations have several shortcomings and are often inaccurate predictors of field frost survival (Pellet et al. 1981). Often, seedlings and young plants are used, which may differ significantly in frost resistance from adults (Sakai and Larcher 1987). Similarly, results obtained with excised plant parts may be unreliable indicators of behavior in the field because ice nucleation temperatures of excised plant parts generally decrease as a result of artifactual supercooling (Ashworth et al. 1985, Sakai and Larcher 1987, Robberecht and Junttila 1992, Flinn and Ashworth 1994, Neuner et al. 1997, Pearce 2001). Furthermore, only frosts occurring in the plant's natural environment allow study of the recuperation and further development of frost-injured plants or plant parts under natural conditions. Because ice nucleation temperatures are affected in laboratory freezing tests, we hypothesized that ice nucleation temperature would also affect frost

resistance, particularly during shoot elongation. Therefore, we frosted plants at their natural growing site in the field using recently developed field freezing equipment that can measure changes in freezing patterns and frost resistance of attached shoots. Because little is known about changes in frost resistance during shoot expansion, we conducted our measurements from bud swelling until full leaf expansion to assess the developmental stage most susceptible to frost damage.

## Materials and methods

### Plant species and experimental site

In situ freezing experiments were conducted at the timberline ecotone (1950 m a.s.l.) on the northwest-facing slope of Mt. Patscherkofel near Innsbruck, Austria (47°12' N, 11°27' E). All major timberline tree species of the European Alps are common at this site. Investigations were carried out on three coniferous tree species, *Larix decidua* Mill., *Picea abies* and *Pinus cembra* L., and two woody angiosperm species frequently found in the subalpine belt: *Sorbus aucuparia* L., a deciduous tree, and *Rhododendron ferrugineum*, an evergreen dwarf shrub. Frost treatments were conducted in situ on twigs of at least four adult trees of each species from May until August over two successive growing periods (2001 and 2002). These periods covered the complete development of the current year's growth.

For comparative purposes, the development of current-year growth from buds to fully expanded leaves was classified in eight developmental stages. Measurements were started on resting buds (s1) and continued during bud swelling (s2) and bud break (s3). During leaf expansion, three stages were identified: the onset of expansion marked by the loss of bud scales (0–30% of full leaf expansion; s4), the main expansion phase (30–60% of full leaf expansion; s5) and full expansion (60–100% of full leaf expansion; s6). Additional measurements were conducted from July 23 to 30 (s7) and from August 1 to 10 (s8) after full leaf expansion had been reached.

### In situ frost treatment

Attached shoots were exposed to controlled freezing temperatures with a recently developed field portable freezing system (MCC-6, BK-Elektronik, Natters, Austria; [http://www.bk-elektronik.com/pages/e\\_develop.htm](http://www.bk-elektronik.com/pages/e_develop.htm)). The freezing system consists of a control unit and six freezing chambers that can each be programmed independently. In contrast with other field frost equipment (cf. Neuner et al. 1997), our system measures frost resistance in situ by exposing plants to six different freezing temperatures in parallel. In each freezing chamber, air temperature is measured at high frequency (8 Hz) with an NTC temperature sensor (SEMI 833 ET, Hygrotec, Titisee-Neustadt, Germany) and recorded with a programmable data logger (CR10X micrologger, Campbell Scientific, Logan, UT). The data logger acts as a control unit and compares air temperature in the freezing chamber with the set-point temperature of a user-defined freezing program. Depending on the result, the switched control port of the data logger either sends a

signal to the relay units or no signal is sent. As a result, a Peltier cooling unit (MAA 050T-12, Melcor, Trenton, NJ) mounted on top of each freezing chamber is turned on or off. The system can be powered by either a 12 V DC or a 230 V AC supply. The maximum temperature difference that can be maintained between ambient air and that inside the freezing chamber is –28 °C.

Each freezing chamber (interior diameter: 11 × 11 × 15 cm) is made of aluminum sheets insulated with 3-cm-thick Styrodur. The base of the chamber opens to permit insertion of a shoot while the shoot remains attached to the plant. A lid equipped with a slot that shuts allows twigs to be inserted and seals the freezing chamber during measurements. To prevent stratification of air inside the freezing chamber, a small ventilator is mounted on the cold side of the Peltier cooling unit.

Controlled in situ freezing programs followed constant cooling and thawing rates of 2 °C h<sup>-1</sup> and 4-h exposure times (Figure 1). Deviations from the preprogrammed set-point temperatures were less than ± 0.2 °C. Rates of 2 °C h<sup>-1</sup> were chosen because they approach naturally occurring freezing rates in subalpine environments and are in accordance with standard laboratory freezing procedures (Sakai and Larcher 1987). The exposure temperatures were selected such that the highest test temperature would cause no damage (LT<sub>0</sub>) and the lowest temperature would kill all leaves (LT<sub>100</sub>). The difference between adjacent target temperatures was less than 1.5 °C.

### Laboratory freezing test

For comparative purposes, frost resistance during shoot expansion was determined on leaves of excised twigs of *P. abies*. Twigs (20 cm) were collected at the site and transported in a moistened and temperature-insulated container to the Institute of Botany in Innsbruck within 1 h of detachment. Shoots were placed on wet paper towels inside polythene bags and exposed to six target freezing temperatures in computer-controlled commercial freezers (Huber, Innsbruck, Austria). Freezing




Figure 1. Controlled in situ freezing treatment (cooling and thawing rates = 2 °C h<sup>-1</sup>; exposure time = 4 h) of expanding *R. ferrugineum* leaves (s5) at 1950 m a.s.l. during the night of July 5, 2002. Two distinct exotherms (E1, E2) were recorded during shoot expansion.

programs were identical to those used for the in situ frost treatments.

#### Viability assay

Frost damage was assessed in situ 3 days after the end of each frost treatment. In summer, during shoot expansion, we observed that 100% frost damage to leaves coincided with 100% frost damage to the whole current-year shoot. The assessment of frost damage therefore focused on leaves. Leaves of all species showed either no frost damage, initial damage, or were completely killed. Partial frost damage was rarely observed. Therefore, the percentage of the leaf area damaged was easily evaluated visually at the field site. Damage ratings were then plotted against treatment leaf temperatures. A classic logistic function was fitted to the data with P-Fit software (Biosoft, Durham, NC). Values of  $LT_{50}$ , i.e., the temperature causing 50% frost damage, were read directly from the fitted curve. The  $LT_0$  value represents the lowest treatment temperature sustained without frost damage,  $LT_i$  denotes the temperature at which incipient frost damage was observed, and  $LT_{100}$  denotes the highest temperature causing 100% tissue death.

#### Ice nucleation temperatures

Concurrently with the in situ frost treatment and the laboratory freezing test, ice nucleation temperatures were measured with type T copper constantan fine-wire thermocouples (welding spot diameter: 0.127 mm). Temperatures were recorded every 12 s with a Campbell Scientific CR10X micrologger. Thermocouples were fixed to the leaves with lightweight, thermally insulated leaf clips during the in situ freezing treatment. During progressive lowering of leaf temperature ( $2\text{ }^{\circ}\text{C h}^{-1}$ ), one to several exotherms were recorded. Ice nucleation temperatures were determined graphically from the temperature record (see Figure 1). During shoot expansion, two different exotherms were usually recorded, a high-temperature exotherm (E1) and a low-temperature exotherm (E2).

#### Statistical data analysis

Frost resistance and ice nucleation temperatures were determined on leaves of randomly chosen shoots of at least three in-

dividuals per species. During each field freezing test, 36 ice nucleation temperatures were recorded concurrently, and six samples were taken for the estimation of frost resistance. For each developmental stage (s1–s8), mean values from at least 20 measurements were calculated. After the data had passed the Kolmogorov-Smirnov test, the significance of differences between frost resistance and ice nucleation temperatures at different stages in each species and between species were determined by analysis of variance (ANOVA) and the Bonferroni test ( $P < 0.01$ ) using SPSS software (SPSS, Chicago, IL).

## Results

#### Attached versus excised shoots

Attached shoots of *P. abies* were significantly more frost resistant than excised shoots from bud break until the end of the summer period (i.e.,  $LT_i$  was  $1.2\text{ }^{\circ}\text{C}$  higher for detached leaves than for attached leaves) (Table 1), indicating a potential underestimation of frost resistance in laboratory freezing tests. Freezing patterns also differed between attached shoots and excised shoots, with a single exotherm recorded for leaves of excised shoots and two exotherms (E1, E2) usually recorded for attached *P. abies* shoots.

#### Changes in frost resistance during shoot elongation

Significant differences ( $P < 0.01$ ) in frost resistance measured in situ were observed between stages of development of the current year's growth for the species tested (Figure 2). The observed changes were species-specific. The most frost-susceptible developmental stages were encountered after the loss of bud scales (s4; *R. ferrugineum*, *S. aucuparia*, *P. cembra*) and during the main expansion phase (s5; *L. decidua*, *P. abies*). Initial frost damage ( $LT_i$ ) occurred between  $-3.8$  and  $-6.5\text{ }^{\circ}\text{C}$  depending on the species. A further drop in temperature of usually  $2\text{ }^{\circ}\text{C}$  killed the new shoot completely. After full leaf expansion, frost resistance increased again, except in *R. ferrugineum* and *P. abies*. These two species were, at nearly all stages, the most frost-susceptible of the species examined. *Larix decidua* had a higher (usually significantly higher) frost

Table 1. Comparison of frost resistance and ice nucleation temperatures ( $^{\circ}\text{C}$ ) measured during shoot expansion in leaves of *P. abies* on excised twigs in a laboratory freezing test and on attached twigs in an in situ freezing test. Values are means of at least  $n = 30$  observations. Significance of differences between mean values of ice nucleation temperatures was tested by ANOVA and the Bonferroni test ( $P < 0.01$ ). Significant differences are indicated by different letters. The significance of differences between mean values of frost resistance was tested with Students *t*-test ( $P < 0.01$ ) and is indicated by an asterisk. Abbreviations: E1 = high-temperature exotherm; E2 = low-temperature exotherm;  $LT_0$  and  $LT_{50}$  = temperatures causing 0 and 50% frost damage, respectively; and  $LT_i$  = temperature at which incipient frost damage was observed.

Developmental stage	Attached twigs (in situ)				Excised twigs		
	Ice nucleation		Frost resistance		Ice nucleation	Frost resistance	
	E1	E2	$LT_i$	$LT_{50}$		$LT_i$	$LT_{50}$
Resting bud	-4.6 a	-12.5 b	-12.8	-13.4	-13.9 c	-12.5	-13.6
Bud swelling	-4.1 a	-11.6 b	-11.7	-12.0	-12.9 c	-11.4	-12.0
Bud breaking	-2.2 a	-6.8 b	-6.9	-7.3 *	-10.1 c	-7.3	-8.2 *
Leaf expansion	-1.8 a	-5.5 b	-5.2 *	-5.6 *	-6.1 b	-4.0 *	-4.9 *




Figure 2. Changes in frost resistance measured in situ during the development of the current year's growth in five woody timberline species during two successive growing periods (2001 and 2002) at 1950 m a.s.l. The  $LT_0$ ,  $LT_{50}$  and  $LT_{100}$  values are the temperatures causing 0, 50 and 100% frost damage, respectively. The  $LT_i$  value is the temperature at which incipient frost damage was observed. The dotted line indicates the absolute minimum air temperature recorded over a 30-year period at Mt. Patscherkofel at 2045 m a.s.l. in May (s1), June (s2–s6), July (s7) and August (s8). The mean sample size per species for each developmental stage was 45.

resistance than other species throughout its development. All species could potentially be frost damaged at the investigation site because absolute air temperature minima (30-year temperature record) are lower than frost resistance during the most susceptible stages of each species.

#### Freezing patterns and ice nucleation temperatures during shoot elongation

Two freezing exotherms were registered for attached shoots in all species at most stages of shoot development (Figure 3A). High-temperature exotherms (E1) were triggered at mean leaf temperatures between  $-1.1$  and  $-2.3$  °C (mean  $\pm$  SE;  $-1.5$  °C  $\pm$  0.2 °C) for all species except *S. aucuparia* in s4 ( $-3.5 \pm 0.6$  °C) (Figure 4B). The low-temperature exotherm (E2) occurred below  $-4.4$  °C and, except for *P. abies* and *L. decidua*, remained constant throughout shoot elongation. Expanding buds of *L. decidua* displayed so-called discontinuous freezing with multiple E2s (Sakai and Larcher 1987), i.e., one E1 at about  $-4.7$  °C followed by a series of small E2s at much lower temperatures (around  $-16.5$  °C). Each E2 marked the killing of one leaf in the bud. Thin layers of basal leaf tissue responsible for leaf dropping (Napp-Zinn 1966) may act as ice barriers. During bud break, the range of E2s diminished and drifted to higher temperatures (data not shown) until, in s4, a single E2 at  $-8.4$  °C marked  $LT_{100}$  for the whole shoot ( $-7.8$  °C). In *S. aucuparia* and *P. cembra*, E2s were observed only during bud break and the expansion period.




Figure 3. Freezing patterns (A) and ice nucleation temperatures (B) measured in situ at 1950 m a.s.l. in five woody plant species during the development of the current year's growth over two successive growing periods (2001 and 2002). (A) Different freezing patterns are indicated as follows: white = freezing with one exotherm; wide-hatched = freezing with two exotherms (one high-temperature exotherm (E1) and one low-temperature exotherm (E2)); and narrow-hatched = freezing with multiple exotherms (one E1 and multiple E2s). (B) The E1 (open symbol) is connected to the E2 (closed symbol) with a solid line. The dotted lines indicate the progressive disappearance of E2s after full expansion of leaves of *S. aucuparia* and *P. cembra*. The mean sample size per species for each developmental stage was 40.




Figure 4. Frost damage was determined by plotting percentage damage against frost treatment temperature. The  $LT_{50}$  values, i.e., the temperature causing 50% frost damage, can be read directly from the fitted curve. The  $LT_0$  value is the lowest treatment temperature sustained without frost damage,  $LT_i$  is the temperature where incipient frost damage was observed, and  $LT_{100}$  is the highest temperature causing 100% tissue death.



### Relationship between freezing exotherms and frost damage

For all species at all developmental stages, E1 occurred at a significantly higher leaf temperature than initial frost damage (see Figure 5 for *R. ferrugineum*). Immediate removal of samples at E1 temperatures revealed extracellular freezing, as tissues were stiffly frozen and were infiltrated after thawing but were never frost damaged. Thus, all developmental stages of all species investigated tolerated extracellular ice formation during summer.

For *R. ferrugineum*, from bud swelling onward, a single E2 was recorded at a mean of  $-6.2^{\circ}\text{C}$ . In contrast, significant changes in frost resistance were measured. In several developmental stages, E2 coincided with frost damage, but during the expansion period, E2 occurred significantly below ( $1.7^{\circ}\text{C}$  lower) the lethal temperature ( $\text{LT}_{100}$ ). Therefore, in these stages (s4 and s5), there was no causal relationship between E2 and frost damage.

In the other species, the relationship between E2 and frost damage was somewhat different (Figure 6). As in *R. ferrugineum*, some expanding *P. abies* needles (s5) were killed at temperatures higher than the E2. In *S. aucuparia*, E2s corresponded with  $\text{LT}_i$  or  $\text{LT}_{100}$ , and in *P. cembra*, with temperatures causing initial frost damage. In *L. decidua*, the E2 matched killing temperatures at all developmental stages except after full expansion when it corresponded with  $\text{LT}_0$  or  $\text{LT}_i$ .

## Discussion

### In situ frost resistance and freezing pattern

In our in situ freezing experiments, all conifers tested were significantly more frost resistant ( $0.5$ – $2.2^{\circ}\text{C}$ ) than indicated in earlier laboratory freezing experiments (Table 2). This finding

is probably associated with differences between in situ and laboratory freezing patterns. Our findings on detached shoots of *P. abies* corroborate results of earlier laboratory freezing experiments where usually only a single freezing exotherm at distinctly lower temperatures than E1 was recorded (Ashworth et al. 1985, Flinn and Ashworth 1994, Neuner and Bannister 1995, Neuner et al. 1997, Fuller and Wisniewski 1998). In contrast, two pronounced freezing exotherms were recorded in our field freezing experiments for all plant species investigated, at most stages of development.

### First freezing event

After bud break, E1s were recorded in leaves of all species at mean temperatures of between  $-1.1$  and  $-2.3^{\circ}\text{C}$  ( $-1.5 \pm 0.2^{\circ}\text{C}$ ), except for *S. aucuparia* in s4 ( $-3.5 \pm 0.6^{\circ}\text{C}$ ). These values correspond well with observations in earlier field freezing experiments where E1s were registered between  $-0.6$  and  $-2.6^{\circ}\text{C}$  (for review see Pearce 2001). This first freezing exotherm reflects extracellular ice formation. In woody plants, the presence of intrinsic nucleators has been confirmed by infrared video thermography, and intrinsic nucleators have been shown to be active near subzero temperatures (Wisniewski et al. 1997, Fuller and Wisniewski 1998, Workmaster et al. 1999, Carter et al. 2001, Pearce and Fuller 2001). Exposure of leaves to temperatures at which these first freezing exotherms were registered was in all cases non-injurious. Thus, leaves of all investigated species tolerated at least some extracellular ice and subsequent cellular dehydration because cell viability was fully retained despite the presence of extracellular ice at E1. This is in accordance with earlier findings (Pearce 1988) that extracellular freezing is not only a phenomenon of freezing-tolerant plants but also occurs in non-acclimated leaves such as barley and in leaves with no capacity for cold acclimation (Pearce 2001). In freezing-sensitive dicotyledonous species and freezing-tolerant cereals, all tissues within leaves are




Figure 5. Relationship between freezing exotherms ( $\nabla$  = E1 and  $\blacktriangle$  = E2) and frost damage (bars:  $\text{LT}_0$ ,  $\text{LT}_i$ ,  $\text{LT}_{50}$  and  $\text{LT}_{100}$ ) in leaves of *R. ferrugineum* measured in situ at 1950 m a.s.l. during the development of the current year's growth over two successive growing periods (2001 and 2002). The mean sample size for each developmental stage was 50.




Figure 6. Relationship between low-temperature exotherm (E2) and the extent of frost damage. Symbols: wide-hatched, E2 is not significantly different from  $\text{LT}_{50}$  or  $\text{LT}_{100}$ ; black, E2 is not significantly different from  $\text{LT}_0$  or  $\text{LT}_i$ ; narrow-hatched, E2 is significantly lower than  $\text{LT}_{100}$ . White fields indicate the absence of E2s. The mean sample size per species for each developmental stage was 40.

Table 2. Frost resistance of immature leaves of woody timberline species measured by different authors compared with the minimum frost resistance determined in our field freezing experiments. Abbreviations: LT<sub>0</sub> and LT<sub>50</sub> = temperatures causing 0 and 50% frost damage, respectively.

Species	Degree of damage	Frost resistance data from other authors	Frost resistance measured in situ
<i>Rhododendron ferrugineum</i>	LT <sub>50</sub>	−3.8 <sup>1</sup> , −4.0 <sup>2</sup>	−3.6
<i>Larix decidua</i>	LT <sub>50</sub>	−5.0 <sup>3</sup>	−6.4
<i>Picea abies</i>	LT <sub>50</sub>	−3.0 <sup>3</sup> , −3.5 <sup>4</sup>	−4.8
<i>Pinus cembra</i>	LT <sub>0</sub>	−2.0 <sup>5</sup> , −3.5 <sup>6</sup>	−4.1

<sup>1</sup> Neuner et al. 1999; <sup>2</sup> Schwarz 1970; <sup>3</sup> Christersson et al. 1987; <sup>4</sup> Repo 1992; <sup>5</sup> Tranquillini 1979; <sup>6</sup> Pisek and Schiessl 1947.

freeze-dehydrated (Pearce 1988, Pearce and Ashworth 1992, Ashworth and Pearce 2002). However, there are also species, such as maize, where certain leaf tissues, such as epidermal and bundle sheath cells, can supercool (Ashworth and Pearce 2002). Although we found an E1 reflecting extracellular ice formation in all leaves, suggesting tolerance of freeze dehydration, supercooling cannot be completely excluded, at least for certain leaf tissues.

Second freezing event and frost damage

All developmental stages showed initial frost damage significantly below E1, indicating that the main mechanism of frost damage, with the possible exception of some supercooling tissues, must be freezing-induced cell dehydration and its consequences. Most E2s measured in situ during shoot expansion matched frost damage (LT<sub>i</sub> to LT<sub>100</sub>) temperatures. These E2s indicate a lethal freezing process. Low-temperature exotherms usually mark the onset of intracellular ice nucleation caused by rupture of cell membranes (Pearce and Willison 1985) or homogeneous ice nucleation after supercooling (Olien 1978, 1981). Although cell membranes are a target for frost damage in freezing-tolerant species, the subcellular target in sensitive species (or stages) has not been identified (Pearce 2001). In expanding shoots of *R. ferrugineum* (s4–s5) and some *P. abies* (s5) shoots, where killing occurred at freezing temperatures higher than E2, the injurious event does not appear to be cell membrane damage because this should have immediately triggered an E2, whereas in *R. ferrugineum* during s4, E2s were not recorded until 2 °C below LT<sub>100</sub>. These exceptions indicate that freezing exotherms are not a reliable estimator of frost damage.

The difference in frost resistance of up to 2.2 °C that we observed for the conifers between the in situ and laboratory tests can be a crucial point for survival of episodic late spring or summer frosts at the timberline ecotone. After loss of current-year growth because of frost damage, which has been repeatedly observed (Kerner 1869; D. Taschler and G. Neuner, personal observations), conifers do not foliate again until the next growing season. Consequently, frost damage contributes to the distorted growth of trees at the tree line, although it does not appear to be a real threat to survival (Tranquillini 1979, Larcher 1985). In *R. ferrugineum*, frost damage to current-year growth can disorder sexual reproduction and flower development for years even though it does not threaten tree sur-

vival (J. Wagner, University of Innsbruck, Austria, personal communication).

Climate warming potentially increases the frequency of unpredictable, episodic frost events (Katz and Brown 1992) and concomitantly promotes earlier bud break (Cannel and Smith 1986, Murray et al. 1989, Hänninen 1991, 1996, Guak et al. 1998). Bud burst is an irreversible process and new shoots may frost-harden to a limited extent only in response to low temperatures. Thus, the risk of frost damage to sprouting leaves may increase with climate warming, as shown for *Pinus sylvestris* L. (Repo et al. 1996). Additionally, elevated atmospheric CO<sub>2</sub> concentration (Lutze et al. 1998, Beerling et al. 2001, Obrist et al. 2001) increases ice nucleation temperatures in some species. Thus, in terms of frost survival, global changes in climate are likely to have a major impact on plant performance at the alpine timberline ecotone.

Acknowledgments

This study was funded by the FWF (Project P14524-Bot) and the “Verein zur Förderung von Südtirolern an der Landesuniversität Innsbruck.” We thank O. Buchner for his guidance on the design and development of the field freezing system and P. Schröcksnadel for permitting free transportation on the Patscherkofel cable car to the Alpengarten laboratory.

References

Ashworth, E.N. and R.S. Pearce. 2002. Extracellular freezing in leaves of freezing sensitive species. *Planta* 214:798–805.

Ashworth, E.N., G.A. Davis and J.A. Anderson. 1985. Factors affecting ice nucleation in plant tissues. *Plant Physiol.* 79:1033–1037.

Bannister, P. and G. Neuner. 2000. Frost resistance and the distribution of conifers. In *Conifer Cold Hardiness*. Eds. F.J. Bigras and S.J. Colombo. Kluwer Academic Publishers, Dordrecht, pp 3–23.

Beerling, D.J., A.C. Terry, P.L. Mitchell, T.V. Callaghan, D. Gwynn-Jones and J.A. Lee. 2001. Time to chill: effects of simulated global change on leaf ice nucleation temperatures of subarctic vegetation. *Am. J. Bot.* 88:628–633.

Cannell, M.G.R. and R.I. Smith. 1986. Climatic warming, spring bud burst, and frost damage on trees. *J. Appl. Ecol.* 23:177–191.

Carter, J., R. Brennan, M. Wisniewski and R. Brennan. 2001. Patterns of ice formation and movement in blackcurrant. *HortScience* 366: 855–859.

Christersson, L., H. von Fircks and Y. Sihe. 1987. Damage to conifer seedlings by summer frost and winter drought. In *Plant Cold Hardiness and Freezing Stress*. Vol. 1. Ed. P.H. Li. Alan R. Liss, New York, pp 203–210.

- Flinn, C.L. and E.N. Ashworth. 1994. Blueberry flower-bud hardiness is not estimated by differential thermal analysis. *J. Am. Soc. Hortic. Sci.* 119:295–298.
- Fuller, M.P. and M. Wisniewski. 1998. The use of infrared thermal imaging in the study of ice nucleation and freezing of plants. *J. Therm. Biol.* 23:81–89.
- Gross, M., I. Rainer and W. Tranquillini. 1991. Über die Frostresistenz der Fichte mit besonderer Berücksichtigung der Zahl der Gefrierzyklen und der Geschwindigkeit der Temperaturänderung beim Frieren und Auftauen. *Forstw. Cbl.* 110:207–217.
- Guak, S., D.M. Olszyk, L.H. Fuchigami and D.T. Tingey. 1998. Effects of elevated CO<sub>2</sub> and temperature on cold hardiness and spring bud burst and growth in Douglas-fir (*Pseudotsuga menziesii*). *Tree Physiol.* 18:671–679.
- Hänninen, H. 1991. Does climate warming increase the risk of frost damage in Northern trees? *Plant Cell Environ.* 14:449–454.
- Hänninen, H. 1996. Effects of climatic warming on northern trees: testing the frost damage hypothesis with meteorological data from provenance transfer experiments. *Scand. J. Res.* 11:17–25.
- Katz, R.W. and B.G. Brown. 1992. Extreme events in a changing climate: variability is more important than averages. *Clim. Change* 21:289–302.
- Kerner, A. 1869. Die Abhängigkeit der Pflanzengestalt von Klima und Boden. *Festschrift der 43. Jahresversammlung Deutscher Naturforscher und Ärzte*, Wagner, Innsbruck 6:29–45.
- Larcher, W. 1985. Kälte und Frost. In *Handbuch der Pflanzenkrankheiten*. Ed. P. Sorauer. Parey, Berlin, pp 107–326.
- Lutze, J.L., J.S. Roden, C.J. Holly, J. Wolfe, J.J.G. Egerton and M.C. Ball. 1998. Elevated atmospheric [CO<sub>2</sub>] promotes frost damage in evergreen tree seedlings. *Plant Cell Environ.* 21:631–635.
- Matuszkiewicz, W. 1977. Spät- und Frühföste als standortsökologischer Faktor in den Waldgesellschaften des Białowieża Nationalparks (Polen). In *Vegetation und Klima*. Ed. H. Dierschke. Cramer, Vaduz, pp 195–233.
- Murray, M.B., M.G.R. Cannell and R.T. Smith. 1989. Date of bud burst of fifteen tree species in Britain following climatic warming. *J. Appl. Ecol.* 26:693–700.
- Napp-Zinn, K. 1966. Anatomie des Blattes. I. Blattanatomie der Gymnospermen. In *Handbuch der Pflanzenanatomie*. Ed. K. Linsbauer. Bornträger, Berlin, pp 1–369.
- Neuner, G. and P. Bannister. 1995. Frost resistance and susceptibility to ice formation during natural hardening in relation to leaf anatomy of three evergreen tree species from New Zealand. *Tree Physiol.* 15:371–377.
- Neuner, G., P. Bannister and W. Larcher. 1997. Ice formation and foliar frost resistance in attached and excised shoots from seedlings and adult trees of *Nothofagus menziesii*. *N.Z. J. Bot.* 35:221–227.
- Neuner, G., D. Ambach and O. Buchner. 1999. Readiness to frost harden during the dehardening period measured in situ in leaves of *Rhododendron ferrugineum* L. at the alpine timberline. *Flora* 194: 289–296.
- Obrist, D., J.A. Arone and C. Körner. 2001. In situ effects of elevated CO<sub>2</sub> on leaf freezing resistance and carbohydrates in a native temperate grassland. *Ann. Bot.* 87:839–844.
- Olien, C.R. 1978. Analyses of freezing stresses and plant responses. In *Plant Cold Hardiness and Freezing Stress*. Vol. 1. Eds. P.H. Li and A. Sakai. Academic Press, New York, pp 317–348.
- Olien, C.R. 1981. Analysis of midwinter freezing stress. In *Analysis and Improvement of Plant Cold Hardiness*. Eds. C.R. Olien and M.N. Smith. CRC Press, Boca Raton, FL, pp 35–59.
- Pearce, R.S. 1988. Extracellular ice and cell shape in frost-stressed cereal leaves: a low temperature scanning electron microscopy study. *Planta* 175:313–324.
- Pearce, R.S. 2001. Plant freezing and damage. *Ann. Bot.* 87: 417–424.
- Pearce, R.S. and E.N. Ashworth. 1992. Cell shape and localization of ice in leaves of overwintering wheat during frost stress in the field. *Planta* 188:324–331.
- Pearce, R.S. and M.P. Fuller. 2001. Freezing of barley studied by infrared video thermography. *Plant Physiol.* 125:227–240.
- Pearce, R.S. and J.H.M. Willison. 1985. A freeze-etch study of the effect of extracellular freezing on the cellular membranes of wheat. *Planta* 163:304–316.
- Pellet, H., M. Gearhart and M. Dir. 1981. Cold hardiness capability of woody ornamental plant taxa. *J. Am. Soc. Hortic. Sci.* 106: 239–243.
- Pisek, A. and R. Schiessl. 1947. Die Temperaturbeeinflussbarkeit der Frosthärte von Nadelhölzern und Zwergsträuchern an der alpinen Waldgrenze. *Ber. Naturwiss.-Med. Ver. Innsbr.* 47:33–52.
- Repo, T. 1992. Seasonal changes of frost hardiness in *P. abies* and *P. sylvestris* in Finland. *Can. J. For. Res.* 22:1949–1957.
- Repo, T., H. Hänninen and S. Kellomäki. 1996. The effects of long-term elevation of air temperature and CO<sub>2</sub> on frost hardiness of Scots pine. *Plant Cell Environ.* 19:209–216.
- Robberecht, R. and O. Junttila. 1992. The freezing response of an arctic cushion plant *S. caespitosa* L. Acclimatization, freezing tolerance and ice nucleation. *Ann. Bot.* 70:129–135.
- Sakai, A. and W. Larcher. 1987. Frost survival of plants. Responses and adaptation to freezing stress. *Ecol. Stud.* 62, Springer-Verlag, Berlin, 321 p.
- Schwarz, W. 1970. Der Einfluß der Photoperiode auf das Austreiben, die Frosthärte und die Hitzeresistenz von Zirben und Alpenrosen. *Flora* 159:258–285.
- Tranquillini, W. 1979. Physiological ecology of the alpine timberline. Tree existence at high altitude with special reference to the European Alps. *Ecol. Stud.* Vol. 31. Springer-Verlag, Berlin, 137 p.
- Wisniewski, M., S.E. Lindow and E.N. Ashworth. 1997. Observation of ice nucleation and propagation in plants using infrared video thermography. *Plant Physiol.* 113:327–334.
- Workmaster, B.A.A., J.P. Palta and M. Wisniewski. 1999. Ice nucleation and propagation in cranberry uprights and fruit using infrared video thermography. *J. Am. Soc. Hortic. Sci.* 124:619–625.

