

## Review

# Responses of forest trees to single and multiple environmental stresses from seedlings to mature plants: Past stress history, stress interactions, tolerance and acclimation

Ülo Niinemets\*

*Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, Kreutzwaldi 1, Tartu 51014, Estonia*

## ARTICLE INFO

### Article history:

Received 23 February 2010

Received in revised form 9 June 2010

Accepted 13 July 2010

### Keywords:

Single and interacting stresses

Non-structural carbon pools

Ontogeny

Photosynthesis

Plasticity

Stress diagnosis

Volatile organic compounds

## ABSTRACT

Forest trees are exposed to a myriad of single and combined stresses with varying strength and duration throughout their lifetime, and many of the simultaneous and successive stress factors strongly interact. While much progress has been achieved in understanding the effects of single stresses on tree performance, multiple interacting stress effects cannot be adequately assessed from combination of single factor analyses. In particular, global change brings about novel combinations of severity and timing of different stresses, the effects of which on tree performance are currently hard to predict. Furthermore, the combinations of stresses commonly sustained by trees change during tree ontogeny. In addition, tree photosynthesis and growth rates decline with increasing tree age and size, while support biomass in roots, stem and branches accumulates and the concentrations of non-structural carbohydrates increase, collectively resulting in an enhancement of non-structural carbon pools. In this review, tree physiological responses to key environmental stress factors and their combinations are analyzed from seedlings to mature trees. The key conclusions of this analysis are that combined stresses can influence survival of large trees even more than chronic exposure to a single predictable stress such as drought. In addition, tree tolerance to many environmental stresses increases throughout the ontogeny as the result of accumulation of non-structural carbon pools, implying major change in sensing, response and acclimation to single and multiple stresses in trees of different size and age.

© 2010 Elsevier B.V. All rights reserved.

## Contents

1. Introduction.....	1624
2. Stress in forests.....	1624
2.1. Occurrence of single and multiple stresses.....	1624
2.2. Stress combinations, interactions and implications for reaction norms.....	1626
3. Stress responsiveness of trees in the current and future climates: tolerance, acclimation and stress history.....	1626
3.1. Tolerance of single and multiple limitations.....	1626
3.2. Influence of acclimation, stress priming and stress sequence on the resistance to multiple stress episodes.....	1627
3.3. Global change and multiple stresses: CO <sub>2</sub> , temperature and drought interactions.....	1627
3.4. Multiple stresses in trees: use of storage carbon to gauge and predict stress responsiveness.....	1628
4. Stress in forest trees: from seedlings to mature plants.....	1628
4.1. Variation in stress combinations throughout tree ontogeny.....	1628
4.2. Effects of ontogeny on stress sensitivity.....	1629
4.3. Determinate vs. indeterminate growth as an explanation for variations in stress tolerance with ontogeny.....	1630
5. Towards adequate diagnosis of stress in forest trees: from single plant to ecosystem responses.....	1631
5.1. Complexity of assessing stress in trees.....	1631
5.2. Traditional methods to diagnose stress.....	1632
5.3. Monitoring volatile organic compounds: a promising method for stress studies.....	1633

\* Tel.: +372 53457189.

E-mail addresses: [ylo.niinemets@emu.ee](mailto:ylo.niinemets@emu.ee), [ylo@emu.ee](mailto:ylo@emu.ee).

6. Conclusions .....	1634
Acknowledgements .....	1634
References .....	1634

## 1. Introduction

Forest environments are characterized by strongly fluctuating conditions for tree growth and development. Sustained deviation of any environmental condition beyond the optimum range reduces plant potential productivity, constituting a stress for the plant. Stressful conditions are part of “normal” plant life in the field, and therefore, understanding the limits of stress tolerance and acclimation to stresses is of great fundamental and practical value in predicting the potential and realized limits of plant productivity (Lichtenthaler, 1998; Isebrands et al., 2000). A key environmental limitation in forest understory is light availability, especially for developing seedlings and saplings (Valladares and Niinemets, 2008). In addition to light, excess water availability and drought, either sustained or periodic, are common stress factors of different types of forests (Franco and Lüttge, 2002; Irvine et al., 2002; Dale and Ware, 2004; Niinemets and Valladares, 2008). Limited nutrient availability is another key environmental limitation significantly curbing growth at many forested areas (Sampson and Allen, 1998; Bergh et al., 1999), whereas often several nutrients such as nitrogen and phosphorus may be simultaneously limiting for plant growth (Niinemets and Kull, 2005; Ågren, 2008). In field environments, often various chronic stresses occur in correlated and interactive manner, for instance, low light in forest understory combined with low soil nutrient availability can be limiting in old growth forests (Jozsa and Brix, 1989; Walters and Reich, 1997; Bergh et al., 1999). Predicting stress responses is especially relevant under current globally changing environmental conditions, i.e., elevated atmospheric CO<sub>2</sub> concentration and air temperature and enhanced probability of drought and stress interactions in world ecosystems (Hobbs et al., 2006), already now being characterized by major shifts in forested biomes (Peñuelas and Boada, 2003; Dobbartin et al., 2005; Grund et al., 2005; Walther et al., 2005; Peñuelas et al., 2007).

Given the relevance of stress on plant performance, extensive work is currently being conducted to understand plant responses, tolerance and acclimation to key environmental stresses (e.g., Mittler, 2006; Günthardt-Goerg and Vollenweider, 2007; Jones and Schofield, 2008; Jackson et al., 2009; Paoletti et al., 2010). Analyzing the influence of stress on plant performance in the field is complicated by occurrence of multiple successive and/or simultaneous stresses, often influencing plant performance interactively, either amplifying or ameliorating the plant responses (Pell et al., 1994; Valladares and Pearcy, 1997; Isebrands et al., 2000; Alonso et al., 2001; Niinemets and Valladares, 2004; Valladares et al., 2005; Mittler, 2006; Rennenberg et al., 2006). Given that global change involves modification of a series of environmental factors concurrently and changes in the severity of different stress factors, knowledge of how plants acclimate to multiple successive or multiple combined stresses is of key significance in understanding the effects of future climates on vegetation.

A further complication in understanding stress responses of forests is that naturally regenerating forest stands typically consist of multiple tree age classes ranging from seedlings to large trees. Numerous tree structural and physiological characteristics change throughout tree ontogeny, including reductions in net assimilation rate and stem hydraulic conductance (Ryan and Yoder, 1997; Bond, 2000; Niinemets, 2002; Mullin et al., 2009), increases in leaf dry mass per unit area (Niinemets and Kull, 1995; Sack and Grubb, 2001; Niinemets, 2002) as well as accumulation of support woody

biomass and non-structural storage pools of carbon (Hoch et al., 2003; Sala and Hoch, 2009). In addition to profound structural and physiological changes, the combinations of stresses commonly sustained by forest trees change during tree ontogeny. For example, establishing trees in the understory gradually intercept more light as they increase in size and reach the canopy, and thus, the limitation of carbon gain by low light is gradually relieved (Givnish, 1988). Yet, enhanced radiation interception together with reduced hydraulic conductance leads to greater water limitations of carbon gain in large trees (Ambrose et al., 2009; Mullin et al., 2009).


The way trees sense stress also varies throughout the ontogeny. In particular, there is evidence of overall greater resistance to drought and ozone stress and biotic stress in large non-senescent trees compared with seedlings and saplings (Fredericksen et al., 1995, 1996a; Kolb et al., 1997; Fleischmann et al., 2005, 2009; Nunn et al., 2005a). In fact, large trees are infrequently killed by even long-term single stress events such as drought (e.g., Bréda et al., 2006) that result in massive mortality of undergrowth (Frazer and Davis, 1988; Donovan et al., 1993; Poorter and Hayashida-Oliver, 2000). Massive decline of large forest trees is commonly associated with multiple sustained stresses (Wargo, 1996; Walters and McCarthy, 1997; Adams et al., 2009a; Allen et al., 2010; Paoletti et al., 2010), highlighting the importance of multiple stresses in understanding the mortality of mature trees.

So far, most of the work on tree stress resistance has been carried out with young plants, most often with first-year seedlings (e.g., Walters and Reich, 1999). Given the changes in stress combinations and overall stress resistance with increasing tree size, the lack of stress studies on older plants is an important deficiency. This synthesis article first provides an overview of the complex environmental setting in forests, effects of single and multiple stresses on plant performance, and the interactions of past and present stresses and tolerance and acclimation to stress combinations under current and predicted future conditions. This paper further analyzes the modifications in stress responsiveness from seedlings to mature trees, and finally provides a short summary of currently used methods and novel promising tools to characterize stress in forest trees of varying size, and to scale the responses to entire trees and forest ecosystems. This analysis collectively suggests that the severity of various single stresses and stress combinations varies throughout tree ontogeny. Furthermore, future climates may be characterized by novel combinations of severity of various stresses. Understanding tree responses to such ontogeny- and global change-driven modifications in stress combinations is needed to predict forest stress responsiveness in current and future climates.

## 2. Stress in forests

### 2.1. Occurrence of single and multiple stresses

According to classical stress concepts, every environmental factor deviating from the optimum constitutes a stress to the plants. While every factor can be individually controlled in lab experiments, a plethora of abiotic and biotic factors fluctuate simultaneously in field environments. The severity and timing of different stresses varies among different ecosystems and during the growing season as outlined in Fig. 1 for temperate bogs supporting low stature conifer tree cover, temperate conifer forests with high stature forest cover and Mediterranean savannas supporting



**Fig. 1.** Generalized illustration of occurrence of severe frequent stresses (solid vertical lines), and mild or sporadic stresses (dotted lines) in three contrasting Northern hemisphere temperate ecosystems dominated by evergreen tree species. Plants mainly sustain stress during the growing period and are less susceptible to stress during the winter frost period when the plants are dormant, especially in cool temperate ecosystems. However, frost stress in cool temperate species can be associated with premature dehardening in response to warmer periods in the middle of winter, resulting in major winter dieback (Ögren, 1996, 1997). The importance of various stress factors varies throughout the growing season in each ecosystem in a characteristic manner as explained in detail in Section 2.1. Sample images of a temperate bog (Männikjärve bog, Estonia, 58° 52' N, 26° 15' E) and a temperate forest (Ihaste Forest, Tartu, Estonia, 58° 21' N, 26° 47' E) both dominated by evergreen conifer *Pinus sylvestris*, and a Mediterranean savanna (montado) dominated by evergreen sclerophylls *Quercus suber* and *Q. ilex* (Mitra, Évora, Portugal, 38° 34' N, 7° 54' W) are also shown.

sparse evergreen tree cover. Temperate bogs are characterized by chronic nutrient deficiency throughout the growing season (Small, 1972; Damman, 1990; Niinemets et al., 2001). Differently from the bogs, nutrient limitation is not always a chronic severe stress in temperate and Mediterranean forests. Severe nutrient limitations can still occur in the beginning of the growing season during new foliage formation (Weikert et al., 1989; Marschner et al., 1997; Saur et al., 2000), and moderate nutrient deficiency can be present during most of the season, especially in forested areas on soils not considered suitable for agricultural crops (Linder, 1987; Carreira et al., 1994; Rambal, 2001; Maroco et al., 2002).


The other key stress in the bogs is anoxia (waterlogging) during a major part of the growing season (Fig. 1). However, as the bulk of the roots occupies the upper 5–15 cm soil layer, mid-seasonal reduction of water table depth can also result in severe drought in bogs (Portsmouth et al., 2005; Portsmouth and Niinemets unpublished observations, Fig. 1) and the reduction in water availability can lead to photoinhibition (Fig. 1, Murray et al., 1989, 1993; Hájek et al., 2008). In temperate forests, drought and photoinhibition can significantly limit productivity, although the importance of these stresses strongly varies between the years in unpredictable manner (Linder, 1987; Rosenqvist et al., 1991; Niinemets et al., 1999b; Niinemets and Kull, 2001). In the Mediterranean environments, prolonged drought and photoinhibition are among the most severe stress factors (Flexas et al., 2001; Werner et al., 2001; Chaves et al., 2009; Unger et al., 2009). In ecosystems with dense canopy, different stresses can be prevalent in the upper (e.g., photoinhibition) and the lower canopy (shading) (Valladares et al., 2005, 2008; Valladares and Niinemets, 2008).

In addition to these stress factors, sub-optimal temperatures constitute a major environmental limitation in the cool temperate ecosystems (Troeng and Linder, 1982), while supraoptimal temperatures occur sporadically (Niinemets et al., 1999a). In contrast, heat stress is a major environmental limitation in the Mediterranean ecosystems (Gratani, 2000; Valladares and Niinemets, 2007;

García-Plazaola et al., 2008), although sub-optimal temperatures can also be limiting for the productivity during winter months (Beyschlag et al., 1987; Harley et al., 1987), and limit the plant survival especially at the northern margin of the Mediterranean evergreens (Larcher, 1981, 2000).

In all these three ecosystems analyzed, the plants commonly sustain multiple stresses, episodically even three or more environmental factors may be beyond the plant optimum. For example, in the bog, nutrient deficiency, waterlogging and photoinhibition, and nutrient deficiency, drought and photoinhibition can co-occur (Fig. 1, Murray et al., 1989, 1993; Tømmervik et al., 1994; Hájek et al., 2008), while the key co-occurring stresses in the Mediterranean ecosystems are drought, heat and photoinhibition (Fig. 1, Valladares and Percy, 1997, 2002; Valladares et al., 2005). Apart from the multiple co-occurring stresses, plants often also sustain multiple successive stresses such as cycles of drought followed by heat spells or waterlogging (Fig. 1, Liu and Dickmann, 1993; Fan et al., 1994; Conrath et al., 2006). For such successive stresses, order, duration and severity of different stresses can significantly vary, resulting in unique stress patterns. In addition to abiotic stresses, various biological stresses such as insect and pathogens attacks are further often superimposed on these key environmental limitations in all forested ecosystems (Schoeneweiss, 1981b, 1983; Kurz et al., 2008). As with the abiotic stress factors, the severity of different biotic stresses can also vary among different ecosystems (Walter, 1992; Kittredge and Ashton, 1995; Zavala et al., 2000; Anthony et al., 2002; Baier et al., 2002) further diversifying the stress patterns.

This evidence collectively demonstrates that forest trees are frequently exposed to a variety of stresses, making the prediction of plant physiological responsiveness to variations in environmental and biotic drivers a formidable task. As the result of common stress factors such as lack of nutrients, water and sub- or supraoptimal temperatures, the realized carbon gain of many forests has for long been recognized to be much less than the potential productivity (Jarvis, 1985; Landsberg, 1986; Linder, 1987).



**Fig. 2.** Illustration of the concept of additive and interactive stresses. Classical stress studies have demonstrated that the rate of plant physiological processes (e.g., photosynthesis, growth) depends on resource availability, e.g., on nutrient availability, according to a bell shaped curve. Under optimum conditions, the level of a given resource is not limiting for plant performance, while super- and supraoptimal resource levels result in reduced plant performance. Typically, any additional stress, e.g., drought, further reduces plant performance. In the case of an additive effect (dashed line), the plant performance is reduced at any given nutrient availability to a similar degree. In the case of an interactive effect (dotted line), the effect of the second stress varies in dependence on the level of the first stress factor. For nutrients, the interactive effect typically implies changes in the form of the response curve, i.e., alteration in the shape of the reaction norm, visible in the shifted optimum nutrient availability ( $N_{opt,1}$  for the responses without an additional stress and with an additive stress and  $N_{opt,2}$  for the interactive stress).

## 2.2. Stress combinations, interactions and implications for reaction norms

Multiple stresses can occur simultaneously or successively. In the case of simultaneous stresses, seldom one encounters a situation with two or more limitations occurring at exactly the same time. More frequently, an episodic stress factor is superimposed on an already existing stressor. For instance, heat stress events are superimposed on drought-stressed plants, or shaded plants in understory experience seasonal drought. The difference between the two multiple stress situations is that previous exposure to the first stress can result in acclimation responses to this first stressor that, in turn, can alter the plant response to the superimposed stress, while such acclimation effects are absent for the overlaying stresses. Thus, tolerance and acclimation to the underlying stress can importantly alter the overall plant response to the combined superimposed stress (Section 3). The circumstances are similar with successive stresses, where acclimation to the first stress can occur during the recovery periods intervening the different stresses.

Due to the acclimation responses and variations in tolerance to different environmental limitations (Section 3) the influence of superimposed and successive environmental stresses is often not additive, but different stresses can interact (Fig. 2). Interactions among the stress factors mean that any additional stress factor alters the reaction norm for any given environmental driver (Fig. 2). The interactions can be either negative, implying amplification of the plant response to the given stressor by an additional stress, or positive, leading to the reduced plant responsiveness to the given stress factor by an additional stress. As an example of negative interactions, in some studies, drought has been found to be more severe under shading (i.e., a negative drought-shade interaction, Valladares and Pearcy, 2002), while in other studies, plants under excess light have been shown to be more vulnerable to drought (a negative drought-high light interaction, Valladares and Pearcy, 1997; Aranda et al., 2001). As another example, waterlogging affected tree saplings more strongly under high light than under low light (Wagner and Dreyer, 1997). Negative interactions are also common for combined abiotic and biotic stresses. Trees exposed to a certain environmental stress factor, e.g., to severe drought or frost, are much more vulnerable to pathogen attacks than plants

not exposed to severe environmental stress (Schoeneweiss, 1981a, 1983; Appel, 1984).

As examples of positive interactions, studies demonstrate that drought-stressed plants are often more resistant to heat stress than plants under optimum water availability, reflecting the positive effects of enhanced content of sugars and other osmotically active compounds in drought-stressed plants in preserving membrane integrity under heat stress (Seemann et al., 1986; Hüve et al., 2006). Drought also leads to stomatal closure, and reduces the uptake of pollutants via stomata, thereby ameliorating the effects of gaseous pollutants, e.g., ozone (Chen et al., 1994; Ollinger et al., 1997; Löw et al., 2006). As a further positive interaction, nitrogen fertilization can reduce the impact of several stresses, e.g., waterlogging stress (Liu and Dickmann, 1993).


## 3. Stress responsiveness of trees in the current and future climates: tolerance, acclimation and stress history

### 3.1. Tolerance of single and multiple limitations

Tree stress responses depend on the plant resistance to any given stress and to any combination of stresses. Stress resistance depends on the tolerance of the specific stress, and on the degree of acclimation to the specific stress. Tolerance is a relative term that characterizes the species capacity to tolerate a given environmental limitation relative to the other species of the same ontogenetic stage (Niinemets and Valladares, 2008). In the past, tolerance was commonly characterized by a qualitative ranking, e.g., very tolerant, tolerant, mid-tolerant, intolerant and very intolerant (e.g. Baker, 1949 for qualitative shade tolerance rankings). Nowadays, quantitative numerical rankings based on combining and cross-calibrating the qualitative rankings or based on species abundance along environmental gradients are becoming increasingly more used (Ellenberg et al., 1991; Hill et al., 1999; Niinemets and Valladares, 2006).

Tree species widely differ in the tolerance to different environmental factors (Niinemets and Valladares, 2006 for tolerance rankings of Northern hemisphere species for light, flooding and drought). While the structural and physiological traits responsible for single factor tolerance are becoming increasingly well understood (Eissenstat, 2000; Lusk and Del, 2002; Sánchez-Gómez et al., 2006; Hallik et al., 2009), we currently poorly understand the factors responsible for the tolerance to multiple environmental drivers. However, there is evidence that multifactor tolerance is rare among the woody plants. In a study investigating the tolerance to shade, drought and waterlogging in Northern hemisphere temperate species, at most 10% of species were tolerant at least to two environmental factors, and ca. 0.4% of species were moderately tolerant to all three environmental factors. In fact, negative correlations between shade and drought tolerance (Fig. 3), shade and waterlogging tolerance and drought and waterlogging tolerance have been observed, suggesting that polytolerance is inherently constrained in temperate woody species (Niinemets and Valladares, 2006). Such negative correlations between tolerances to different stresses likely reflect different combinations of structural and physiological traits required to tolerate given environmental limitation (Smith and Huston, 1989; Hallik et al., 2009).

Although the tolerance of several chronic stresses is rare, the capacity to tolerate short-term biotic and abiotic stress events superimposed on a chronic environmental stress such as shade, can be critical to survive under the main chronic environmental limitation. For instance, fungal pathogen attacks have been shown to limit recruitment of tree seedlings in shade (O'Hanlon-Manners and Kotanen, 2004). The capacity to survive short-term additional biotic and abiotic stresses under shade has been linked to the size



**Fig. 3.** Relationships between shade and drought tolerance (standardized major axis regression lines) among the Northern hemisphere temperate tree species from Europe ( $n=262$ , solid line), North America ( $n=364$ , dotted line) and East Asia ( $n=211$ , dashed line). The inset shows the comparisons among the slopes according to the maximum-likelihood ratio test of Warton and Weber (Falster et al., 2003; Warton et al., 2006). Both the shade and drought tolerance are defined as numerical relative rankings between 1 (lowest tolerance) to 5 (highest tolerance). Modified from Niinemets and Valladares (2006).

of the storage pool of non-structural carbohydrates (Myers, 2005; Myers and Kitajima, 2007; Poorter and Kitajima, 2007). Greater pool of non-structural carbohydrates can contribute to the survival of the episodes of additional stresses both by directly covering plant metabolic requirements as well as by fuelling structural and metabolic acclimation to additional stresses (Amthor and McCree, 1990).

### 3.2. Influence of acclimation, stress priming and stress sequence on the resistance to multiple stress episodes

Stress acclimation is defined as full or partial recovery of plant functioning during environmental stress and lower stress responsiveness compared to non-acclimated plants. The degree of plant acclimation to the given stress depends on plant phenotypic plasticity and the time available for stress hardening. Therefore, past stress history can importantly alter plant responsiveness to environmental stresses. Exposure to mild stress typically results in induction of acclimation responses leading to greater resistance to a given type of stress (Conrath et al., 2006). For instance, a mild drought stress episode enhances the tree resistance to a subsequent more severe drought stress (Liu and Dickmann, 1993; Fan et al., 1994). Analogously, an early defoliation event results in greater contents of protective chemicals, thereby enhancing the resistance to subsequent herbivory episodes (Stevens and Lindroth, 2005). This so-called stress priming may enhance the resistance not only to a given stress, but to several subsequent stresses. For example, acquired resistance to pathogens can result in development of resistance to herbivory (Heil and Silva Bueno, 2007; Frost et al., 2008; Niinemets, 2010b) or even to abiotic stresses such as drought (Fujita et al., 2006).

For sequential multiple stresses, plant response may critically depend on what stress comes first and on the degree to which the plant has acclimated to this first stress factor. For example, the plants already acclimated to long-term low light conditions in the understory and having large leaf area and a low fraction of biomass in roots are expected to respond more strongly to drought (e.g., Valladares and Pearcy, 2002) than the plants recently overtopped by other competing plant individuals and not yet having substantially altered allocation of biomass to leaves at the expense of roots.

As a further example, plants exposed to long-term drought and having depleted the carbon reserves will have low resistance to pathogen attacks (Schoeneweiss, 1981a), while pathogen attacks preceding drought stress may activate defence pathways that can contribute to drought resistance as well (Fujita et al., 2006, see also below). In particular, accumulation of the stress hormone abscisic acid in pathogen-infected leaves can lead to stomatal closure and water conservation (Lee et al., 1999; Flors et al., 2005; Asselbergh et al., 2008). However, the patterns largely vary for different pathogens, and in some cases, pathogen attack can force stomatal opening, resulting in greater vulnerability to subsequent drought events (Allègre et al., 2007; Prats et al., 2007).

In addition to the stress sequence, stress severity and duration of stress events can also drive the overall plant response. Exposure to a severe stress in the past may reduce plant immediate capacity to cope with any subsequent event of the same type of stress or different stress. This is because a severe stress can deplete plant storage carbon reserves, leading to exhaustion of plant resistance to stress (Lichtenthaler, 1998). In most experimental stress studies, only one stress cycle has been applied. However, experiments with repeated stress cycles have demonstrated that multiple successive stress cycles impair plant performance more strongly than a single stress cycle (e.g., for drought stress cycles, Alcocer-Ruthling et al., 1989; Liu and Dickmann, 1993; Stewart et al., 1995). It takes increasingly more time to recover from the next stress cycle (Alcocer-Ruthling et al., 1989; Liu and Dickmann, 1993; Stewart et al., 1995), reflecting the exhaustion of plant resistance by recurrent stress episodes. This evidence suggests that apart from induced mechanisms of stress resistance, the availability of carbon reserves at the beginning of each stress episode plays a major role in determining the resistance against a further stress of the same type or an additional stress. Thus, repeated past stress events intervened by short periods of recovery insufficient to fill up the carbon reserves reduce the plant performance to any subsequent stress. However, when sufficient time is available between the stress episodes for recovery, priming responses may enhance plant resistance against further stresses. Thus, a delicate balance seems to exist between priming for stress resistance and depletion of carbon reserves, jointly determining the way the plant stress history alters the plant response to subsequent stresses. At any rate, these examples demonstrate that duration, severity and the order of different stresses importantly alter the overall effect of multiple stresses on plants.

### 3.3. Global change and multiple stresses: CO<sub>2</sub>, temperature and drought interactions

Predicted increases in temperature and changes in precipitation pattern (Meehl et al., 2007; Trenberth et al., 2007) are likely to lead to occurrence of novel combinations of stresses in ecosystems that have not encountered such stress combinations in the past as well as to novel combinations of severity of various stresses (Williams and Jackson, 2007). For example, enhanced temperatures can lead to occurrence of heat and waterlogging stress in bogs and more severe heat, drought and photoinhibition stress periods in temperate bog and forest ecosystems (Fig. 1). In future warmer climates, drought periods are generally expected to be longer and more severe (Breshears et al., 2005; Adams et al., 2009a; Allen et al., 2010).

Consideration of carbon reserves in plant stress resistance is particularly relevant in understanding the effects of key global change drivers on forest growth. This is because all global change associated modifications in environment occur at the background of CO<sub>2</sub> elevation, and typically, elevated CO<sub>2</sub> increases the non-structural carbon storage (Wullschlegel et al., 1992; Roberntz and Stockfors, 1998; Tognetti et al., 1998; Vu et al., 2002), and enhances the resistance to several stresses. Plants grown in elevated CO<sub>2</sub> atmospheres

have been shown to be less vulnerable to drought (Conroy et al., 1986; Le Thiec and Dixon, 1996; Roden and Ball, 1996; Anderson and Tomlinson, 1998; Tomlinson and Anderson, 1998; Schwanz and Polle, 2001; Polley et al., 2002; Vu et al., 2002), maintaining higher growth rate under drought than plants under lower CO<sub>2</sub> atmospheres (Tscharplinski et al., 1995; Polley et al., 2002). Elevated CO<sub>2</sub> also enhances plant resistance to heat and frost stresses (Faria et al., 1996; Wayne et al., 1998; Taub et al., 2000; Vu et al., 2002; Darbush et al., 2010), likely reflecting greater concentrations of membrane-stabilizing sugars in the tissues (for sugar effects on heat stability of photosynthetic apparatus see Hüve et al., 2006) (for sugar effects on frost stability see Rütten and Santarius, 1988; Santarius, 1990).

Apart from these positive effects, elevated CO<sub>2</sub> commonly induces greater nutrient deficiency (for reviews see Curtis, 1996; Saxe et al., 1998). This in turn can lead to accumulation of secondary carbon-rich chemicals such as tannins, improving plant resistance to herbivory attacks as observed in several studies (Roth et al., 1997; McDonald et al., 1999; Agrell et al., 2000; Mohan et al., 2006). On the other hand, if the content of secondary chemicals is not affected by elevated CO<sub>2</sub>, greater leaf sugar concentrations may promote herbivory attacks under elevated CO<sub>2</sub> (DeLucia et al., 2008).

Predicting stress responses in the future climates is further complicated by concomitant modifications in temperature. Warming and associated increase in plant respiratory requirements can result in depletion of carbon reserves (Ahmed et al., 1993; Tjoelker et al., 1998, 2008; Campbell et al., 2007). This may enhance the vulnerability of plants to additional stresses such as to ozone stress (Riikonen et al., 2009) and drought stress (Xu et al., 2008; Adams et al., 2009a; Allen et al., 2010). Depletion of carbon reserves has been suggested to be the primary reason for the drought-driven mortality at warmer temperatures, reflecting “carbon starvation” (McDowell et al., 2008; Adams et al., 2009a). However, so far the evidence of enhanced carbon limitations under combined drought and warming treatment is lacking (Adams et al., 2009b; Sala, 2009; Sala et al., 2010). Furthermore, projected enhancement of tree mortality in future climates due to warming-drought interaction (Adams et al., 2009a; Allen et al., 2010) has not addressed the concomitant modifications in atmospheric CO<sub>2</sub> concentration (Sala et al., 2010 for a discussion). In fact, the combined effects of elevated CO<sub>2</sub> and warming on sugars has been examined only in a few studies. These experiments have consistently observed higher carbohydrate contents in plants grown under elevated CO<sub>2</sub> at both normal and elevated temperatures (e.g., Tjoelker et al., 1998; Vu et al., 2002), suggesting that the effects of elevated CO<sub>2</sub> on carbon reserves are dominating. This suggestion is supported by greater tolerance of simultaneous drought and heat stress episodes in plants grown under elevated CO<sub>2</sub> (Hammerlynck et al., 2000).

Alternative to “carbon starvation” hypothesis, greater mortality of trees under combined warming and drought treatment may be associated with hydraulic failure (McDowell et al., 2008) or/and with impaired long-distance transport of non-structural carbon from storage tissues to sinks (Sala et al., 2010), especially if drought occurs rapidly (Leuzinger et al., 2009). Lack of non-structural carbon pool data in trees undergoing drought- and warming-induced death makes it currently impossible to discriminate among the different hypotheses (Sala, 2009). Clearly, additional research on storage carbohydrate pools is needed for quantitative prediction of plant stress responsiveness in future conditions.

#### 3.4. Multiple stresses in trees: use of storage carbon to gauge and predict stress responsiveness

Apart from the resilience to sustained chronic stresses mainly determined by species tolerance, tree response to environmental variability at any time during the growing season is also critically

driven by tree acclimation capacity and the past stress history. Tree responses may be completely different depending on whether the past stress induces acclimation responses, possible only when there is enough time for recovery and acclimation, or whether the past stress results in depletion of carbon reserves. Given the unpredictability of the period and length of recurrent stress cycles during the growing season, the effects of past stress history have been so far particularly difficult to consider. I suggest that the effects of multiple successive and superimposed stresses and past stress history effects are best included in predictive models at the level of storage carbon dynamics. As storage carbon pool is one of the key players affected by elevated CO<sub>2</sub>, dynamic simulation of plant carbon allocation and storage becomes especially relevant in predicting plant stress responsiveness under future climates (Constable and Friend, 2000 for review of models).


Measurements of storage carbon are not routinely conducted in tree studies, and there is an emerging consensus that more experimental research in tree carbon reserves is needed, especially under environmental stress (Adams et al., 2009b; Leuzinger et al., 2009; Sala, 2009; McDowell and Sevanto, 2010; Sala et al., 2010). Furthermore, only a few growth models consider the reserve carbon pool, and even in the models including the storage carbon, the dynamics of this pool is often considered in a limited manner, essentially as a “black box” without considering external and internal feedbacks (Le Roux et al., 2001). Even in fewer models, mediation of the stress effects at the level of carbon pools are considered (Chen, 1993; Chen et al., 1994; Weinstein and Yanai, 1994; Retzlaff et al., 1997). In addition to simulating the storage carbon pool, stress models should ideally also be able to simulate the investment in defence, for instance on the basis of carbon/nitrogen balance (Gayler and Priesack, 2007). Consideration of defence investments together with storage would make an important interface to consider plant resistance to biotic stresses as well.

## 4. Stress in forest trees: from seedlings to mature plants

### 4.1. Variation in stress combinations throughout tree ontogeny

The previous sections demonstrated the complexity of forest tree responses to various single and combined environmental and biotic stresses. Apart from the complex interplay between different stresses, important shifts in the severity of key stresses and modifications in tree stress sensitivity occur throughout the tree ontogeny. In particular, with increasing tree size, the depth of the tree root system commonly increases, resulting in access to deeper soil water supplies (Dawson, 1996; de Soyza et al., 1996; Drake et al., 2009). The water supply from these deeper water sources, in particular from groundwater reservoirs, is less affected by precipitation-free periods than the water availability in the upper soil layers; modifications of water availability in surface soil layers can already occur during short-term drought events (Dawson, 1996). This suggests that the water status of larger trees is less affected by lack of precipitation than the water status of smaller trees (Dawson, 1996). In fact, in open-grown woody plants, smaller plants have been shown to suffer from greater water deficiencies (Donovan and Ehleringer, 1992; de Soyza et al., 1996; Drake et al., 2009) and have greater drought-dependent mortality (Donovan et al., 1993) than larger plants.

However, the situation can be different in plants growing in closed forests. In closed stands, the foliage of larger trees receives gradually more light with increasing size. In addition to light, temperature is higher at greater heights in the forest canopy due to greater interception of solar radiation (Niinemets and Valladares, 2004 for a review). Higher temperatures in turn imply greater evaporative demands that can constrain water availability in the crowns




**Fig. 4.** Tree size-dependent increases in leaf dry mass per unit area ( $M_A$ , a) and decreases in photosynthetic capacity per unit dry mass ( $A_{max}$ , b) in two wide-spread European conifer species (modified from Niinemets, 2002). Photosynthetic capacity per area ( $A_{max}M_A$ ) scaled also negatively with tree height, although the variation was narrower, 3–4-fold across the height range due to the opposite trends in  $M_A$  and  $A_{max}$  (data not shown, see, Niinemets, 2002 for details).

of tall trees. In addition, gravitational component of water potential increases by  $0.01 \text{ MPa m}^{-1}$ . To compensate for this increase in water potential, and be able to maintain water supply to the top of the large plants, water potentials in the crowns of tall trees must be significantly lower than in smaller trees (Tobiessen et al., 1971; Koch et al., 2004). Furthermore, there is evidence that the hydraulic conductivity of larger trees is less than in smaller plants (Woodruff et al., 2007; Ambrose et al., 2009). Although the larger trees may have access to deeper soil water reservoirs, their lower water potentials, limited hydraulic conductivity, and greater evaporative demands can under certain conditions (e.g., under low atmospheric humidity) effectively uncouple the foliage at the top of the tree from the water in the soil, resulting in lower stomatal conductance in crowns of larger forest trees than in the smaller trees (Yoder et al., 1994; Fredericksen et al., 1996b; Bond, 2000). These reductions in stomatal conductance are partly responsible for lower rates of photosynthesis in larger trees (Fig. 4, Yoder et al., 1994; Bond, 2000).

Higher temperature and greater light itself can constitute a significant stress for larger trees, especially in conjunction with limited water availability at greater heights in the canopy. There is evidence that photoinhibitory damage to foliage photosynthetic apparatus is more severe under limited water availability and heat stress (Havaux, 1992; Flexas et al., 1999; Ishida et al., 1999), and significant photodamage in larger trees can occur in years with exceptionally severe additional stresses such as drought (García-Plazaola et al., 2008). On the other hand, extensive shading in the understory importantly curbs the carbon gain in tree seedlings and saplings, being the major limitation for forest regrowth (Walters and Reich, 2000; Valladares and Niinemets, 2008).

Among other environmental factors, nutrient availability has been considered limiting for the growth of larger trees in old-growth forests, where limiting nutrients can become largely fixed in standing biomass and soil organic matter (Gower et al., 1996; Murty et al., 1996). Thus, productivity of large trees in old-growth forests can be limited even by three chronic environmental




**Fig. 5.** Generalized responses of tree size-dependent alterations in photosynthesis, stem hydraulic conductance (based on Niinemets, 2002; Koch et al., 2004; Woodruff et al., 2007, 2009; Mullin et al., 2009; Steppe et al., 2011), overall plasticity in growth and biomass allocation to environmental modifications, and severity and responsiveness to different stresses and stress combinations. In forests, young seedlings and saplings typically grow under low limiting light availabilities, but light availability gradually improves as the trees gain in size. With increasing radiation input, evaporative demand also increases. This, combined with reduced hydraulic conductance, potentially leads to greater water stress (e.g., Hubbard et al., 1999; Bond, 2000), and also commonly results in greater leaf temperatures (Niinemets et al., 1999a). However, the situation can be different for open-grown seedlings and saplings that commonly have shallower root systems than mature trees, and may therefore sustain greater drought-dependent mortality (Donovan and Ehleringer, 1992; Donovan et al., 1993). In old-growth forests, accumulation of limiting nutrients such as nitrogen in litter, soil organic matter and standing biomass may also result in reductions in stand growth (Gower et al., 1996). Simultaneous occurrence of drought and photoinhibition and heat stresses is expected to increase with increasing tree size, while shading and drought stress interaction is expected to be more important in seedlings and saplings (for shade–drought interactions in understory and photoinhibition–drought interactions in overstory see Valladares and Pearcy, 1997, 2002; Niinemets and Valladares, 2004). Due to larger carbon reserves, the overall stress sensitivity of large mature non-senescent trees is expected to be less than in seedlings and saplings.

stresses: limited water and nutrient availabilities and excess light, while the productivity of undergrowth is limited by low light and nutrient availabilities.

This evidence collectively suggests that trees of varying size encounter different combinations of environmental limitations throughout their ontogeny (Fig. 5). Such ontogenetic modifications in the severity of various key stresses have important implications for transferring the knowledge of stress responses from the lab studies conducted with seedlings to the forest stands, as well as for prediction of forest tree growth and survival from seedlings to mature trees.

#### 4.2. Effects of ontogeny on stress sensitivity

In addition to changes in the stress combinations, the stress sensitivity also importantly varies from seedlings to mature trees. With increasing tree size, the fraction of plant biomass in stems and roots increases, while the fraction of biomass in leaves decreases (Niinemets, 1998, 2010a; Jokela and Martin, 2000; Messier and Nikinmaa, 2000; Claveau et al., 2005; Portsmouth et al., 2005). In addition, the concentration of non-structural carbon also increases with increasing tree size (Fig. 6), possibly reflecting greater inhibition of growth than photosynthesis with increasing size (Körner, 2003; Sala and Hoch, 2009). Thus, together with the accumulation



**Fig. 6.** Variations in stem sapwood, branch sapwood and 1-year-old needle non-structural carbohydrate concentrations (NSC, sum of sugars and starch) in different-sized trees of *Pinus ponderosa* (a) (dry site, Sala and Hoch, 2009), and in NSC in current-year needles of different-sized trees of *Picea abies* (Niinemets, 1997a,b). In *P. ponderosa*, the trend of increasing NSC with increasing tree height was significant for all three biomass fractions on the dry site shown here ( $P < 0.014$ ), whereas on moist site, the trend was significant for branch sapwood (Sala and Hoch, 2009). In *P. abies*, the difference between the groups is significant at  $P < 0.005$ . The error bars show +SE ( $n = 5$  for all tree size classes in *P. ponderosa* and for small trees of *P. abies* and  $n = 3$  for large trees of *P. abies*).

of structural carbon in the woody biomass, the reserves of non-structural carbon (starch and soluble sugars) in stems and roots (concentration times the mass of roots and stem) increase with increasing tree size (Hoch et al., 2003; Körner, 2003; Würth et al., 2005; Sala and Hoch, 2009). In large trees, the non-structural carbon stocks may exceed several-fold the amount of carbon needed to form the entire canopy leaf area (Hoch et al., 2003; Würth et al., 2005).

The variations in the size of non-structural carbon reserves bear important implications for understanding changes in stress resistance throughout tree ontogeny (Fig. 7). Typically, any stress results in the reduction of plant net carbon gain (Fig. 7). Influence of stress-driven reductions in net assimilation rate on plant survival apparently depends on the extent to which the total tree non-structural carbohydrate pool relies on immediate photosynthesis. In seedlings, the storage carbon pool size is relatively small, and a reduction in net assimilation rate leads to fast depletion of the storage reserves in plant maintenance processes (Fig. 7), further accentuating the stress effect on the plant. In contrast, in large trees, immediate carbon assimilation contributes only to a minor degree to the total tree non-structural carbon pools, and stress-dependent reduction in carbon uptake only moderately alters total tree carbon reserves (Fig. 7). As the result of these differences, seedlings that more strongly rely on immediate photosynthesis are expected to be more sensitive to prolonged stress events than large trees that are less strongly dependent on immediate photosynthetic production (Fig. 7).


Available evidence is in broad agreement with these predictions. The resilience of trees to fungal attacks increases significantly during the tree ontogeny (Fleischmann et al., 2005, 2009). Analogously, the responsiveness to ozone stress is higher (Fredericksen et al., 1995, 1996a; Kolb et al., 1997; Nunn et al., 2005a) and the overall antioxidative capacity (Wieser et al., 2003) is lower in seedlings than in mature trees. Tree water stress resistance also increases with increasing tree size, accompanying greater fractional investment of plant biomass in roots in larger plants (Anderson and Tomlinson, 1998; Tomlinson and Anderson, 1998). In fact, large trees can tolerate sustained drought periods extending over the entire growing season and even across multiple growing seasons (Epron and Dreyer, 1993), while such severe drought periods can lead to mortality of seedlings and saplings (for seedling physiological responses see Epron et al., 1993; Mendel et al., 1997; Mueller et al., 2005; Lopez and Kursar, 2007). The presence of large carbon reserves in mature plants can also allow for new shoot forma-

tion and resprouting (in resprouting species) following catastrophic events such as fire (Koskela, 2000; Langley et al., 2002) and breakage of branches or entire stem (Génard et al., 1998; Verdaguer and Ojeda, 2002; van Bloem et al., 2003; Wong et al., 2005; Vargas, 2009; Vargas et al., 2009). Thus, the availability of large reserves contributes to survival of and greater recovery after such devastating events. In contrast, lack of sufficient reserves to build a new stem and foliage following the death of above-ground plant part makes smaller plants highly vulnerable to biomass losses. In addition, larger plants have a greater capacity to tolerate sustained periods of combined stresses such as ozone and pathogen stress (Fleischmann et al., 2009). Due to greater antioxidative capacity under higher light in adult trees than in seedlings (Wieser et al., 2003), adult trees are expected to tolerate better the interactions of photoinhibition with other stresses such as ozone stress. In plants with limited content of antioxidants, such combined stresses have been shown to lead to extensive formation of reactive oxygen species (ROS) and extensive damage (Wise, 1995; Garcia-Plazaola and Becerril, 2001; Tausz et al., 2001; Wieser et al., 2001).

#### 4.3. Determinate vs. indeterminate growth as an explanation for variations in stress tolerance with ontogeny

The reason for higher carbon pools in larger forest trees is partly associated with their access to greater light availabilities. In fact, given the lower photosynthetic rates and greater fraction of non-photosynthetic support biomass, tolerance of low light conditions decreases with increasing tree size (Valladares and Niinemets, 2008 for a review). However, even if grown in higher light under optimum conditions, the seedlings or the saplings partition less carbon in storage than the mature trees. This is because seedlings and saplings are commonly characterized by indeterminate biomass growth or multiple growth cycles during the growing season, while mature trees, especially in seasonal environments, have a determinate growth pattern characterized by a distinct period of rapid growth in the beginning of the favourable season (e.g., Borchert, 1991). Beyond the rapid growth period, much of the carbon produced is stored in large trees (Hoch et al., 2003).

The capacity for indeterminate growth implies that younger trees can more plastically respond to biomass losses (Stevens and Lindroth, 2005) and modify biomass allocation and construct foliage that is acclimated to prevailing conditions. Greater growth plasticity can be particularly advantageous in the acclimation to short stress events as well as in priming for future stress events



**Fig. 7.** Comparison of stress responses in seedlings and trees: a conceptual model. Most stress factors result in reduction in plant net assimilation rate, reducing the newly assimilated carbon pool present in leaves immediately after photosynthesis and further translocated to growing organs (developing leaves, fine roots, and woody structures) and used for maintenance respiration throughout the plant body possessing living cells, and also used to build up the storage carbon reserves. Apart from the newly synthesized carbon, all woody plants have significant non-structural carbon (sugars and starch) reserves in leaves, stem and branches and in roots (Fig. 1, Hoch et al., 2003; Myers and Kitajima, 2007; Poorter and Kitajima, 2007; Sala and Hoch, 2009) that can be used to support plant maintenance requirements throughout the stress periods when photosynthesis rates are low. In seedlings, a larger part of total non-structural carbon (newly assimilated carbon plus storage carbon) relies on newly assimilated carbon. Thus, reductions in the net carbon assimilation rate are essentially immediately reflected in plant survival. In contrast, mature trees have huge carbon reserves, even corresponding to several years growth (Hoch et al., 2003; Würth et al., 2005), and newly assimilated carbon contributes a small fraction to total tree non-structural carbon. Thus, the survival of trees during stress is less dependent on reductions in immediate rate of carbon gain. Boxes are not to scale.

(Stevens and Lindroth, 2005). For instance, drought tolerance has been associated with enhancement of root growth under moderate drought in small-seeded seedlings of several Mediterranean species (Padilla et al., 2007). However, under severe prolonged stress that strongly inhibits growth, it is not always clear whether the variation in plastic modifications in plant function (stronger vs. smaller reduction in growth rate) is associated with fitness. In fact, plants capable of growing in low-fertility habitats with chronic drought and nutrient stress have commonly low growth plasticity (Grime, 1989; Lambers and Poorter, 1992). Survival under shade has also been found to be even negatively associated with the plasticity in growth (Walters and Reich, 1996; Kaelke et al., 2001).


This discussion demonstrates that tree stress responsiveness is critically driven by both the dimensions of time (stress length, stress prehistory) and size (storage pool, exposure to different stresses). In the forest ecosystems, mortality is a frequent event in smaller tree size classes (Lorimer et al., 2001; Palahi et al., 2003; van Mantgem et al., 2009), reflected in massive reduction in stand density in early stages of stand development (Lorimer et al., 2001; Palahi et al., 2003). With increasing tree size, mortality rate commonly decreases (Lorimer et al., 2001; Palahi et al., 2003). Even extended periods of severe single stress typically do not lead to mortality of large trees (Bréda et al., 2006; Bigler et al., 2007). How-

ever, stresses co-occurring with extreme environmental events such as pathogen or herbivory attacks in drought-stressed plants with significantly exhausted pools of storage carbon can result in major forest decline episodes (Schoeneweiss, 1981a, 1983; Mattson and Haack, 1987; McLaughlin, 1998; Manion, 2003; Bréda et al., 2006; Kurz et al., 2008). Such catastrophic events can be more frequent in old-growth stands, reflected in enhanced mortality of older trees (Lorimer et al., 2001).

## 5. Towards adequate diagnosis of stress in forest trees: from single plant to ecosystem responses

### 5.1. Complexity of assessing stress in trees

Understanding the stress effects in trees and forest stands critically depends on the capacity to diagnose stress. A plethora of methods have been developed to study stress in plants (Fig. 8) and commonly a series of tools are used simultaneously to characterize tree physiological status in the field (Nunn et al., 2005b). However, trees largely vary in size throughout the ontogeny from seedlings to mature trees. While most methods can be applied to seedlings, saplings and larger trees constitute a major challenge



**Fig. 8.** Overview of the methods used to diagnose stress in plants. The stress diagnosis methods can be divided between non-destructive (commonly also non-invasive) methods that do not require harvesting of the plant or a plant part and destructive (always invasive) methods that require harvesting of the entire plant or a plant part (typically leaves). A plethora of methods exists within each sub-division, and here only the examples of widespread methods are provided. The advantage of non-destructive methods is that the same plant or the plant part can be repeatedly sampled and evolution of stress can be monitored essentially continuously (the inset for a sample relationship between net assimilation rate and time of drought treatment that can be assessed in the same plant or leaf). In the case of invasive methods, either new plants or new plant parts must be sampled at different times during stress development. This is associated with inherent variability between the samples and also entails that continuous monitoring is impossible (the inset for a sample response of leaf water potential and duration of drought, where each data point necessarily corresponds to an individual leaf). However, typically non-destructive methods provide an integrated estimate of plant performance and provide limited information of sensing the stress and modifications in plant metabolism, with some exceptions. For example, studies on volatile compound emissions (volatile plant hormones and the compounds released as the result of oxidative stress in the plant) can provide information of the time-kinetics of elicitation of stress-signalling network (Pichersky and Gershenzon, 2002; Martin et al., 2003; Beauchamp et al., 2005; Niinemets, 2010b).

for stress studies. In particular, strong environmental gradients of light and temperature exist in the crowns of large trees (e.g., Niinemets and Valladares, 2004). These gradients are accompanied by several-fold variations in foliage photosynthetic capacities and pigment contents (for reviews see e.g., Nunn et al., 2005b; Niinemets, 2007) as well as in variations in foliage sensitivity to environmental stresses (Niinemets et al., 2003; Aasamaa et al., 2004; García-Plazaola et al., 2004). Due to this large within-crown environmental and physiological heterogeneity, gaining insight into environmental limitations in large trees using conventional leaf-level methods requires large sampling effort and also models to integrate from leaf-level observations to the whole plant response (Matyssek et al., 2008; Paoletti et al., 2010). Some methods are also available to directly diagnose stress at whole plant or at entire ecosystem scale. However, as shown below, many of the available methods, especially large-scale methods, are relatively unspecific and may not always be sensitive enough in detecting early stress stages or an overall mild stress. In the following, a case is made that more specific methods need to be developed to diagnose early stages of stress and monitor stress development and recovery.

## 5.2. Traditional methods to diagnose stress

The available stress diagnosing methods can be broadly divided between non-destructive (commonly also non-invasive) methods that allow for repeated monitoring of the same plants or plant organs, and destructive (always invasive) methods that require

destructive harvesting of either the entire plant or plant organ (Fig. 8). Among the non-destructive methods, the simplest methods are various visual indices based on foliage colour, degree of leaf damage (necrotic spots, fraction of area removed by herbivores) and degree of full crown defoliation (crown transparency) (Innes and Boswell, 1990; Schmid-Haas, 1993; Tausz et al., 1997; Solberg, 1999; Wood et al., 2000; Nunn et al., 2005b; Vollenweider and Günthardt-Goerg, 2005; Aldea et al., 2006). However, these visual indices demonstrate already the presence of a severe stress, and are not suitable for diagnosing a mild stress. They also do not allow for reliable monitoring of the stress evolution kinetics. As an extension of visual estimates, various spectral indices have been constructed to monitor stress (Buschmann et al., 1991; Samson et al., 2000; Styliniski et al., 2002; Coops et al., 2003; Richardson et al., 2004; Porder et al., 2005; Schaepman-Strub et al., 2006; Jones and Schofield, 2008; Delaieux et al., 2009; Hilker et al., 2009). These spectral tools can be applied at scales ranging from leaves to whole tree canopies and ecosystems and provide an important means to assess stress using remote sensing techniques. Compared to visual estimates, spectral indices provide higher resolution into stress development kinetics. Using specific wavebands, the spectral methods can be used to simultaneously study various processes such as changes in foliage water content, chlorophyll and carotenoid content, foliage temperature, and elicitation of xanthophyll cycle as well as changes in canopy transparency (Styliniski et al., 2002; Coops et al., 2003; Schaepman-Strub et al., 2006; Jones and Schofield, 2008; Delaieux et al., 2009). However, the sensitivity of these spectral tools may be limited by the opacity of thick leaves

and dense canopies of many forest species, leading to signal saturation (Gamon et al., 1995; Asner et al., 2000; Porder et al., 2005; Kumar et al., 2010).

Measurements of foliage photosynthesis rates either by monitoring carbon uptake or chlorophyll fluorescence provide an important tool to gaining insight into modifications in foliage physiological activity. The rates of photosynthesis have been frequently applied to diagnose stress in the field (Long et al., 1996; Chaerle and Van der Straeten, 2000; Maxwell and Johnson, 2000; Laisk et al., 2001; Long and Bernacchi, 2003; Nunn et al., 2005b), including mature trees (Nunn et al., 2005b). Detailed photosynthetic measurements separating between partial limitations of photosynthesis (e.g., Laisk et al., 2001) are possible only with single leaves, and replicate sampling coupled with models is needed to integrate from leaf-level measurements to whole plant (Jarvis, 1995; Infante et al., 1997; Ollinger et al., 1997; Karnosky et al., 2005; Matyssek et al., 2008). The sensitivity of photosynthesis to different stresses varies in dependence on plant tolerance, and this may make it difficult to relate the reduction in photosynthetic function directly to the severity of environmental limitation. For example, the degree of photoinhibitory reduction of photosynthesis is greater in more shade tolerant species (Krause et al., 2001; Valladares et al., 2002, 2005). A given soil water limitation also often reduces photosynthesis less in more drought tolerant species that can maintain stomata open at lower soil water potentials than the less tolerant species (Flexas and Medrano, 2002; Chaves et al., 2009).

For seedlings and saplings, whole plant measurements provide an alternative way to gain insight into the integrated alterations in plant photosynthetic process during stress (Klingeman et al., 2005; Baltzer and Thomas, 2007). Whole tree photosynthesis measurement systems enclosing 8–14 m<sup>2</sup> plant leaf area have been constructed (Corelli-Grappadelli and Magnanini, 1993; Lloyd et al., 1995; Wünsche and Palmer, 1997; Wallin et al., 2001; Medhurst et al., 2006). However, extensive cost, slow response time and alteration of ambient climatic conditions complicate the employment of such large systems to assess the tree carbon gain.

Alternatively, the physiological activity of large trees can be assessed by measuring sap-flow rate at the stem base using thermoelectric sensors (e.g., Granier et al., 1996; Köstner et al., 1998; James et al., 2002). Given the strong correlation between plant physiological activity and plant water use, these measurements can provide important insight into the development of stress in forest trees (e.g., Alsheimer et al., 1998; Pataki et al., 1998; Nadezhdina, 1999; Rothe et al., 1999; Tenhunen et al., 2001). From whole tree measurements of transpiration, estimates of canopy conductance can be derived (Ewers et al., 2001). These values of canopy conductance can be further used to estimate the uptake of gaseous pollutants such as ozone, making it possible to directly evaluate whole tree ozone stress (Matyssek et al., 2008).

At the ecosystem scale, stress-dependent alterations in the physiological activity of forest stands can be assessed by eddy covariance techniques (Baldocchi, 1997; Arneth et al., 1999; Law et al., 2001). However, as the latter method provides an integrated estimate of both the soil and the plant activities, separate measurements of the soil respiratory activity or modelling approaches are needed to separate the vegetation and soil contributions to whole ecosystem fluxes (e.g., Janssens et al., 2000). Growth studies using dendrometers (Breitsprecher and Bethel, 1990; Vogel et al., 1996; Schleppei et al., 1999; Nunn et al., 2005b) can also be used to gain insight into the environmental variations during tree growth period. Analyses of tree ring time series can further be employed to diagnose the biotic and abiotic conditions for growth during a specific year. However, the latter measurements have inherently low time resolution and are therefore hard to associate with climatic conditions during the given year. In addition, a significant part of fixed carbon is stored and not immediately used for tree

stem growth (e.g., Fig. 7). Especially in seasonal climates, intensive diameter growth occurs over a limited time period even under optimum conditions. Thus, the growth measurements alone are not always strongly linked to stress effects on carbon assimilation.

Most methods outlined above provide an integrated estimate of tree physiological status, and do not necessarily allow separation of different stresses as well as gaining insight into various physiological processes affected by stress. A variety of invasive methods are available to diagnose stress-related modifications in plant metabolism, including analyses of plant elemental composition, pigment contents, antioxidant production and secondary chemistry and gene expression (Fig. 8). Some of the invasive methods such as analyses of foliage elemental composition, pigment contents and water potential are routinely used together with non-invasive methods to characterize plant physiological status. In addition to these methods, studies on gene expression and proteomics are becoming increasingly more often used in trees. Such detailed analyses provide important insight into the stress-dependent elicitation of defence pathways, accumulation of stress-elicited protective compounds and acclimation to stress (Mittler et al., 2001; Pelloux et al., 2001; Andersson et al., 2004; Keskitalo et al., 2005; Fracheboud et al., 2009). However, as discussed above, large sampling effort is needed to gain insight into the whole tree stress status from these detailed measurements that are necessarily conducted at the leaf scale.

### 5.3. Monitoring volatile organic compounds: a promising method for stress studies

Environmental and biotic stresses elicit a cascade of events starting with the oxidative signalling and concluding with the modification in plant phenotype (acclimation) or with the programmed cell death (Doke, 1997; Dizengremel, 2001; Vranová et al., 2002; Baier et al., 2005). During signal transduction, a plethora of signal compounds are formed, many of which are volatile, including NO, methyljasmonate, methylsalicylate and ethylene (Arimura et al., 2002; Degenhardt and Lincoln, 2006; Filella et al., 2006; Gas et al., 2009). Such emissions of volatile stress hormones have been monitored in response to heat (Ibrahim et al., 2006), ozone (Beauchamp et al., 2005) and waterlogging (Copolovici and Niinemets, 2010) stress as well as in response to various biotic stresses (Filella et al., 2006).

In addition, several volatile markers of the onset of stress-dependent metabolic pathways are also released from the stressed plants; some of these compounds can also be involved in propagation of stress signals (Arimura et al., 2009). For instance, methanol is emitted from the stressed plants as the result of demethylation of cell wall pectins by pectin methylesterases (Micheli, 2001), while lipoxygenase pathway products (LOX products, volatile C<sub>6</sub> aldehydes, so-called green leaf volatiles) are emitted as the result of the release of free fatty acids from membranes and activation of various lipoxygenases (Porta and Rocha-Sosa, 2002; Matsui, 2006). As the result of expression of different terpenoid synthases, both biotic and abiotic stresses can lead to elicitation of emissions of mono- and sesquiterpenes (Loughrin et al., 1997; Staudt and Bertin, 1998; Mandujano-Chávez et al., 2000; Steindel et al., 2005; Opitz et al., 2008; Copolovici et al., 2009). The emissions of key stress volatiles, such as LOX products, have been demonstrated to be quantitatively linked to the severity of the stress (Beauchamp et al., 2005; Steindel et al., 2005; Loreto et al., 2006). In addition, different volatile compounds are elicited at different times during stress development (Beauchamp et al., 2005). Thus, the measurements of volatile emissions provide a promising way to diagnose both the severity of the stress and to assess the kinetics of stress development.

Availability of rapid sensors to gauge the emissions of volatile compound emissions such as the Proton Transfer Reaction Mass

Spectrometer (PTR-MS, Lindinger et al., 1998) has made it possible to diagnose stress at leaf to ecosystems scales in the field (Niinemets, 2010b for a review). So far, PTR-MS combined with eddy co-variance methodology has been used to measure the emissions of stress volatile methylsalicylate from forest stands (Karl et al., 2008). These emissions were quantitatively correlated with heat spells (Karl et al., 2008), demonstrating that diagnosing forest stress by monitoring volatile compound emissions is principally possible.

## 6. Conclusions

Due to interactions among environmental drivers, forest stands exposed to different combinations of light, water and nutrient availabilities can have unique responsiveness to environmental modifications. Global change is further predicted to bring about novel interactions among the severity and timing of environmental limitations, cardinally shifting the degree of stress in different communities (Hobbs et al., 2006). The difficulties with diagnosing multiple stresses and assessing the effects on plants are associated with overall lack of knowledge of how the reaction norms to given environmental factor change in response to any additional simultaneous or successive stress and how the reaction norms change with tree ontogeny. More experimental work is needed to characterize the effects of various abiotic and biotic stress combinations and gain more quantitative insight into how reaction norms change during tree ontogeny.

Currently the effects of various stresses cannot be deconvoluted in field environments, but storage carbon size likely provides an important point of convergence of various stress pathways. The predictions based on non-structural carbon pools suggest that maintenance of higher storage carbon pools increases the tolerance of sustained stress episodes. The existing evidence indicates an increase in storage carbon pools with increasing tree size, likely explaining the enhancement of single and combined stress resistance during tree ontogeny. Thus, general predictive forest tree growth models including the effects of multiple stresses, past stress history and ontogeny can be principally developed on the basis of the dynamics of storage carbon pools.

## Acknowledgements

The work of ÜN on plant stress tolerance and volatile organic compound emissions has been funded by the Estonian Science Foundation (Grant 7645), and the Estonian Ministry of Education and Science (grant SF109006507). I thank Fernando Valladares and an anonymous reviewer for thoughtful comments on the manuscript.

## References

- Aasamaa, K., Söber, A., Hartung, W., Niinemets, Ü., 2004. Drought acclimation of two deciduous tree species of different layers in a temperate forest canopy. *Trees* 18, 93–101.
- Adams, H.D., Guardiola-Claramonte, M., Barron-Gafford, G.A., Villegas, J.C., Breshears, D.D., Zou, C.B., Troch, P.A., Huxman, T.E., 2009a. Temperature sensitivity of drought-induced tree mortality portends increased regional die-off under global change-type drought. *Proc. Natl. Acad. Sci. U.S.A.* 106, 7063–7066.
- Adams, H.D., Guardiola-Claramonte, M., Barron-Gafford, G.A., Villegas, J.C., Breshears, D.D., Zou, C.B., Troch, P.A., Huxman, T.E., 2009b. Reply to Sala: temperature sensitivity in drought-induced tree mortality hastens the need to further resolve a physiological model of death. *Proc. Natl. Acad. Sci. U.S.A.* 106, E69.
- Agrell, J., McDonald, E.P., Lindroth, R.L., 2000. Effects of CO<sub>2</sub> and light on tree phytochemistry and insect performance. *Oikos* 88, 259–272.
- Ågren, G.I., 2008. Stoichiometry and nutrition of plant growth in natural communities. *Annu. Rev. Ecol. Syst.* 39, 153–170.
- Ahmed, F.E., Hall, A.E., Madore, M.A., 1993. Interactive effects of high temperature and elevated carbon dioxide concentration on cowpea [*Vigna unguiculata* (L.) Walp.]. *Plant Cell Environ.* 16, 835–842.
- Alcocer-Ruthling, M., Robberecht, R., Thill, D.C., 1989. The response of *Bouteloua scirpioides* to water stress at two phenological stages. *Bot. Gaz.* 150, 454–461.
- Aldea, M., Frank, T.D., DeLucia, E.H., 2006. A method for quantitative analysis of spatially variable physiological processes across leaf surfaces. *Photosynth. Res.* 90, 161–172.
- Allège, M., Daire, X., Héloir, M.-C., Trouvelot, S., Mercier, L., Adrian, M., Pugin, A., 2007. Stomatal deregulation in *Plasmopara viticola*-infected grapevine leaves. *New Phytol.* 173, 832–840.
- Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D.D., Hogg, E.H., Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Limp, J.-H., Allard, G., Running, S.W., Semerci, A., Cobb, N., 2010. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecol. Manage.* 259, 660–684.
- Alonso, R., Elvira, S., Castillo, F.J., Gimeno, B.S., 2001. Interactive effects of ozone and drought stress on pigments and activities of antioxidative enzymes in *Pinus halepensis*. *Plant Cell Environ.* 24, 905–916.
- Alsheimer, M., Köstner, B., Falge, E., Tenhunen, J.D., 1998. Temporal and spatial variation in transpiration of Norway spruce stands within a forested catchment of the Fichtelgebirge, Germany. *Ann. Sci. Forest.* 55, 103–123.
- Ambrose, A.R., Sillett, S.C., Dawson, T.E., 2009. Effects of tree height on branch hydraulics, leaf structure and gas exchange in California redwoods. *Plant Cell Environ.* 32, 743–757.
- Amthor, J.S., McCree, K.J., 1990. Carbon balance of stressed plants: a conceptual model for integrating research results. In: Alsheimer, R.G., Cumming, J.R. (Eds.), *Stress Responses in Plants: Adaptation and Acclimation Mechanisms*. Wiley-Liss, Inc., New York, pp. 1–15.
- Anderson, P.D., Tomlinson, P.T., 1998. Ontogeny affects response of northern red oak seedlings to elevated CO<sub>2</sub> and water stress. I. Carbon assimilation and biomass production. *New Phytol.* 140, 477–491.
- Andersson, A., Kesikitalo, J., Sjödin, A., Bhalerao, R., Sterky, F., Wissel, K., Tandre, K., Aspeborg, H., Moyle, R., Ohmiya, Y., Bhalerao, R., Brunner, A., Gustafsson, P., Karlsson, J., Lundberg, J., Nilsson, O., Sandberg, G., Strauss, S., Sundberg, B., Uhlen, M., Jansson, S., Nilsson, P., 2004. A transcriptional timetable of autumn senescence. *Genome Biol.* 5, R24.
- Anthony, P.A., Holtum, J.A.M., Jackes, B.R., 2002. Shade acclimation of rainforest leaves to colonization by lichens. *Funct. Ecol.* 16, 808–816.
- Appel, D.N., 1984. Canker expansion on water stressed pin oaks colonized by *Endothia gyrosa*. *Plant Dis.* 68, 851–853.
- Aranda, I., Gil, L., Pardos, J.A., 2001. Effects of thinning in a *Pinus sylvestris* L. stand on foliar water relations of *Fagus sylvatica* L. seedlings, planted within the pinewood. *Trees* 15, 358–364.
- Arimura, G.-I., Matsui, K., Takabayashi, J., 2009. Chemical and molecular ecology of herbivore-induced plant volatiles: proximate factors and their ultimate functions. *Plant Cell Physiol.* 50, 911–923.
- Arimura, G.I., Ozawa, R., Nishioka, T., Boland, W., Koch, T., Kühnemann, F., Takabayashi, J., 2002. Herbivore-induced volatiles induce the emission of ethylene in neighbouring lima bean plants. *Plant J.* 29, 87–98.
- Arneth, A., Kelliher, F.M., McSeveny, T.M., Byers, A.N., 1999. Assessment of annual carbon exchange in a water-stressed *Pinus radiata* plantation: an analysis based on eddy covariance measurements and an integrated biophysical model. *Glob. Change Biol.* 5, 531–545.
- Asner, G.P., Wessman, C.A., Bateson, A.C., Privette, J.L., 2000. Impact of tissue, canopy, and landscape factors on the hyperspectral reflectance variability of arid ecosystems. *Remote Sens. Environ.* 74, 69–84.
- Asselbergh, B., De Vleeschauwer, D., Höfte, M., 2008. Global switches and fine-tuning—ABA modulates plant pathogen defense. *Mol. Plant: Microbe Interact.* 21, 709–719.
- Baier, M., Kandlbinder, A., Gollack, D., Dietz, K.-J., 2005. Oxidative stress and ozone: perception, signalling and response. *Plant Cell Environ.* 28, 1012–1020.
- Baier, P., Führer, E., Kirsitis, T., Rosner, S., 2002. Defence reactions of Norway spruce against bark beetles and the associated fungus *Ceratocystis polonica* in secondary pure and mixed species stands. *Forest Ecol. Manage.* 159, 73–86.
- Baker, F.S., 1949. A revised tolerance table. *J. Forest.* 47, 179–181.
- Baldocchi, D., 1997. Measuring and modelling carbon dioxide and water vapour exchange over a temperate broad-leaved forest during the 1995 summer drought. *Plant Cell Environ.* 20, 1108–1122.
- Baltzer, J.L., Thomas, S.C., 2007. Physiological and morphological correlates of whole-plant light compensation point in temperate deciduous tree seedlings. *Oecologia* 153, 209–223.
- Beauchamp, J., Wisthaler, A., Hansel, A., Kleist, E., Miebach, M., Niinemets, Ü., Schurr, U., Wildt, J., 2005. Ozone induced emissions of biogenic VOC from tobacco: relations between ozone uptake and emission of LOX products. *Plant Cell Environ.* 28, 1334–1343.
- Bergh, J., Linder, S., Lundmark, T., Elfving, B., 1999. The effect of water and nutrient availability on the productivity of Norway spruce in northern and southern Sweden. *Forest Ecol. Manage.* 119, 51–62.
- Beyschlag, W., Lange, O.L., Tenhunen, J.D., 1987. Photosynthese und Wasserhaushalt der immergrünen mediterranen Hartlaubpflanze *Arbutus unedo* L. im Jahreslauf am Freilandstandort in Portugal. II. Charakteristische Gaswechselformparameter von CO<sub>2</sub>-Aufnahme und Transpiration. *Flora* 179, 399–420.
- Bigler, C., Gavin, D.G., Gunning, C., Veblen, T.T., 2007. Drought induces lagged tree mortality in a subalpine forest in the Rocky Mountains. *Oikos* 116, 1983–1994.
- Bond, B.J., 2000. Age-related changes in photosynthesis of woody plants. *Trends Plant Sci.* 5, 349–353.

- Borchert, R., 1991. Growth periodicity and dormancy. In: Raghavendra, A.S. (Ed.), *Physiology of Trees*. John Wiley & Sons Inc., New York, pp. 221–245.
- Bréda, N., Huc, R., Granier, A., Dreyer, E., 2006. Temperate forest trees and stands under severe drought: a review of ecophysiological responses, adaptation processes and long-term consequences. *Ann. Forest Sci.* 63, 625–644.
- Breitsprecher, A., Bethel, J.S., 1990. Stem-growth periodicity of trees in a tropical wet forest of Costa Rica. *Ecology* 71, 1156–1164.
- Breshears, D.D., Cobb, N.S., Rich, P.M., Price, K.P., Allen, C.D., Balice, R.G., Romme, W.H., Kastens, J.H., Floyd, M.L., Belnap, J., Anderson, J.J., Myers, O.B., Meyer, C.W., 2005. Regional vegetation die-off in response to global-change-type drought. *Proc. Natl. Acad. Sci. U.S.A.* 102, 15144–15148.
- Buschmann, C., Rinderle, U., Lichtenthaler, H.K., 1991. Detection of stress in coniferous forest trees with the VIRAF spectrometer. *IEEE Trans. Geosci. Remote Sens.* 29, 96–100.
- Campbell, C., Atkinson, L., Zaragoza-Castells, J., Lundmark, M., Atkin, O., Hurry, V., 2007. Acclimation of photosynthesis and respiration is asynchronous in response to changes in temperature regardless of plant functional group. *New Phytol.* 176, 375–389.
- Carreira, J.A., Niell, F.X., Lajtha, K., 1994. Soil nitrogen availability and nitrification in Mediterranean shrublands of varying fire history and successional stage. *Biogeochemistry* 26, 189–209.
- Chaerle, L., Van der Straeten, D., 2000. Imaging techniques and the early detection of plant stress. *Trends Plant Sci.* 5, 495–501.
- Chaves, M.M., Flexas, J., Pinheiro, C., 2009. Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Ann. Bot.* 103, 551–560.
- Chen, C.W., 1993. The response of plants to interacting stresses: PGSM version 1.3 model documentation. In: Technical Report EPRI-TR-101880. Electric Power Research Inst., Palo Alto, CA, USA.
- Chen, C.W., Tsai, W.T., Gomez, L.E., 1994. Modeling responses of ponderosa pine to interacting stresses of ozone and drought. *Forest Sci.* 40, 267–288.
- Claveau, Y., Messier, C., Comeau, P.G., 2005. Interacting influence of light and size on aboveground biomass distribution in sub-boreal conifer saplings with contrasting shade tolerance. *Tree Physiol.* 25, 373–384.
- Conrath, U., Beckers, G.J.M., Flors, V., García-Agustín, P., Jakab, G., Mauch, F., Newman, M.-A., Pieterse, C.M.J., Poinssot, B., Pozo, M.J., Pugin, A., Schaffrath, U., Ton, J., Wendehenne, D., Zimmerli, L., Mauch-Mani, B., 2006. Priming: getting ready for battle. *Mol. Plant: Microbe Interact.* 19, 1062–1071.
- Conroy, J.P., Smillie, R.M., Küppers, M., Bevege, D.I., Barlow, E.W., 1986. Chlorophyll a fluorescence and photosynthetic and growth responses of *Pinus radiata* to phosphorus deficiency, drought stress, and high CO<sub>2</sub>. *Plant Physiol.* 81, 423–429.
- Constable, J.V.H., Friend, A.L., 2000. Suitability of process-based tree growth models for addressing tree response to climate change. *Environ. Pollut.* 110, 47–59.
- Coops, N.C., Stone, C., Culvenor, D.S., Chisholm, L.A., Merton, R.N., 2003. Chlorophyll content in eucalypt vegetation at the leaf and canopy scales as derived from high resolution spectral data. *Tree Physiol.* 23, 23–31.
- Copolovici, L., Kännaste, A., Niinemets, Ü., 2009. Gas chromatography–mass spectrometry method for determination of monoterpene and sesquiterpene emissions from stressed plants. *Studia Universitatis Babeş-Bolyai* 54, 329–339.
- Copolovici, L., Niinemets, Ü., 2010. Flooding induced emissions of volatile signalling compounds in three tree species with differing waterlogging tolerance. *Plant Cell Environ.* 33, 1582–1594.
- Corelli-Grappadelli, L., Magnanini, E., 1993. A whole-tree system for gas-exchange studies. *Hortscience* 28, 41–45.
- Curtis, P.S., 1996. A meta-analysis of leaf gas exchange and nitrogen in trees grown under elevated carbon dioxide. *Plant Cell Environ.* 19, 127–137.
- Dale Jr., E.E., Ware, S., 2004. Distribution of wetland tree species in relation to a flooding gradient and backwater versus streamside location in Arkansas, U.S.A. *J. Torrey Bot. Soc.* 131, 177–186.
- Damman, A.W.H., 1990. Nutrient status of ombrotrophic peat bogs. *Aquilo Ser. Bot.* 28, 5–14.
- Darbah, J.N.T., Sharkey, T.D., Calfapietra, C., Karnosky, D.F., 2010. Differential response of aspen and birch trees to heat stress under elevated carbon dioxide. *Environ. Pollut.* 158, 1008–1014.
- Dawson, T.E., 1996. Determining water use by trees and forests from isotopic, energy balance and transpiration analyses: the roles of tree size and hydraulic lift. *Tree Physiol.* 16, 263–272.
- de Soyza, A.G., Franco, A.C., Virginia, R.A., Reynolds, J.F., Whitford, W.G., 1996. Effects of plant size on photosynthesis and water relations in the desert shrub *Prosopis glandulosa* (Fabaceae). *Am. J. Bot.* 83, 99–105.
- Degenhardt, D.C., Lincoln, D.E., 2006. Volatile emissions from an odorous plant in response to herbivory and methyl jasmonate exposure. *J. Chem. Ecol.* 32, 725–743.
- Delaieux, S., Somers, B., Verstraeten, W.W., Van Aardt, J.A.N., Keulemans, W., Coppin, P., 2009. Hyperspectral indices to diagnose leaf biotic stress of apple plants, considering leaf phenology. *Int. J. Remote Sens.* 30, 1887–1992.
- DeLucia, E.H., Casteel, C.L., Nabity, P.D., O'Neill, B.F., 2008. Insects take a bigger bite out of plants in a warmer, higher carbon dioxide world. *Proc. Natl. Acad. Sci. U.S.A.* 105, 1781–1782.
- Dizengremel, P., 2001. Effects of ozone on the carbon metabolism of forest trees. *Plant Physiol. Biochem.* 39, 729–742.
- Dobbertin, M., Hilker, N., Rebetez, M., Zimmermann, N.E., Wohlgemuth, T., Rigling, A., 2005. The upward shift in altitude of pine mistletoe (*Viscum album* ssp. *auriacum*) in Switzerland—the result of climate warming? *Int. J. Biometeorol.* 50, 40–47.
- Doke, N., 1997. The oxidative burst: roles in signal transduction and plant stress. In: Scandalios, J.G. (Ed.), *Oxidative Stress and the Molecular Biology of Antioxidant Defenses*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 785–813.
- Donovan, L.A., Ehleringer, J.R., 1992. Contrasting water-use patterns among size and life-history classes of a semi-arid shrub. *Funct. Ecol.* 6, 482–488.
- Donovan, L.A., Mausberg, J., Ehleringer, J.R., 1993. Seedling size and survival for *Chrysothamnus nauseosus*. *Great Basin Nat.* 53, 237–245.
- Drake, P.L., Mendham, D.S., White, D.A., Ogden, G.N., 2009. A comparison of growth, photosynthetic capacity and water stress in *Eucalyptus globulus* coppice regrowth and seedlings during early development. *Tree Physiol.* 29, 663–674.
- Eissenstat, D., 2000. Root structure and function in an ecological context. *New Phytol.* 148, 353–354.
- Ellenberg, H., Weber, H.E., Düll, R., Wirth, V., Werner, W., Paulißen, D., 1991. *Zeigerwerte von Pflanzen in Mitteleuropa*. Verlag Erich Goltze KG, Göttingen.
- Epron, D., Dreyer, E., 1993. Long-term effects of drought on photosynthesis of adult oak trees [*Quercus petraea* (Matt.) Liebl. and *Quercus robur* L.] in a natural stand. *New Phytol.* 125, 381–389.
- Epron, D., Dreyer, E., Aussenac, G., 1993. A comparison of photosynthetic responses to water stress in seedlings from 3 oak species: *Quercus petraea* (Matt.) Liebl., *Q. rubra* L. and *Q. cerris* L. *Ann. Sci. Forest.* 50, 48–60.
- Ewers, B.E., Oren, R., Phillips, N., Stromgren, M., Linder, S., 2001. Mean canopy stomatal conductance responses to water and nutrient availabilities in *Picea abies* and *Pinus taeda*. *Tree Physiol.* 21, 841–850.
- Falster, D.S., Warton, D.I., Wright, I.J., 2003. (S)MATR: Standardized Major Axis Tests and Routines Version 1.0. Department of Biological Sciences, Macquarie University, Sydney, Australia.
- Fan, S., Blake, T.J., Blumwald, E., 1994. The relative contribution of elastic and osmotic adjustments to turgor maintenance of woody species. *Physiol. Plant.* 90, 408–413.
- Faria, T., Wilkins, D., Besford, R.T., Vaz, M., Pereira, J.S., Chaves, M.M., 1996. Growth at elevated CO<sub>2</sub> leads to down-regulation of photosynthesis and altered response to high temperature in *Quercus suber* L. seedlings. *J. Exp. Bot.* 47, 1755–1761.
- Filella, I., Peñuelas, J., Llusià, J., 2006. Dynamics of the enhanced emissions of monoterpenes and methyl salicylate, and decreased uptake of formaldehyde, by *Quercus ilex* leaves after application of jasmonic acid. *New Phytol.* 169, 135–144.
- Fleischmann, F., Koehl, J., Portz, R., Beltrame, A.B., Oßwald, W., 2005. Physiological change of *Fagus sylvatica* seedlings infected with *Phytophthora citricola* and the contribution of its elicitor “Citricolin” to pathogenesis. *Plant Biol.* 7, 650–658.
- Fleischmann, F., Winkler, J.B., Oßwald, W., 2009. Effects of ozone and *Phytophthora citricola* on non-structural carbohydrates of European beech (*Fagus sylvatica*) saplings. *Plant Soil* 323, 75–84.
- Flexas, J., Badger, M., Chow, W.S., Medrano, H., Osmond, C.B., 1999. Analysis of the relative increase in photosynthetic O<sub>2</sub> uptake when photosynthesis in grapevine leaves is inhibited following low night temperatures and/or water stress. *Plant Physiol.* 121, 675–684.
- Flexas, J., Guliás, J., Jonasson, S., Medrano, H., Mus, M., 2001. Seasonal patterns and control of gas exchange in local populations of the Mediterranean evergreen shrub *Pistacia lentiscus* L. *Acta Oecol.* 22, 33–43.
- Flexas, J., Medrano, H., 2002. Drought-inhibition of photosynthesis in C<sub>3</sub> plants: stomatal and non-stomatal limitations revisited. *Ann. Bot.* 89, 183–189.
- Flors, V., Ton, J., Jakab, G., Mauch-Mani, B., 2005. Abscissic acid and callose: team players in defence against pathogens? *J. Phytopathol.* 153, 377–383.
- Fracheboud, Y., Luquez, V., Björkén, L., Sjödin, A., Tuominen, H., Jansson, S., 2009. The control of autumn senescence in European aspen. *Plant Physiol.* 149, 1982–1991.
- Franco, A.C., Lüttge, U., 2002. Midday depression in savanna trees: coordinated adjustments in photochemical efficiency, photorespiration, CO<sub>2</sub> assimilation and water use efficiency. *Oecologia* 131, 356–365.
- Frazer, J.M., Davis, S.D., 1988. Differential survival of chaparral seedlings during the first summer drought after wildfire. *Oecologia* 76, 215–221.
- Fredericksen, T.S., Joyce, B.J., Skelly, J.M., Steiner, K.C., Kolb, T.E., Kouterick, K.B., Savage, J.E., Snyder, K.R., 1995. Physiology, morphology, and ozone uptake of leaves of black cherry seedlings, saplings, and canopy trees. *Environ. Pollut.* 89, 273–283.
- Fredericksen, T.S., Skelly, J.M., Steiner, K.C., Kolb, T.E., Kouterick, K.B., 1996a. Size-mediated foliar response to ozone in black cherry trees. *Environ. Pollut.* 91, 53–63.
- Fredericksen, T.S., Steiner, K.C., Skelly, J.M., Joyce, B.J., Kolb, T.E., Kouterick, K.B., Ferdinand, J.A., 1996b. Diel and seasonal patterns of leaf gas exchange and xylem water potentials of different-sized *Prunus serotina* Ehrh. trees. *Forest Sci.* 42, 359–365.
- Frost, C.J., Mescher, M.C., Carlson, J.E., De Moraes, C.M., 2008. Plant defense priming against herbivores: getting ready for a different battle. *Plant Physiol.* 146, 818–824.
- Fujita, M., Fujita, Y., Noutoshi, Y., Takahashi, F., Narusaka, Y., Yamaguchi-Shinozaki, K., Shinozaki, K., 2006. Crosstalk between abiotic and biotic stress responses: a current view from the points of convergence in the stress signaling networks. *Curr. Opin. Plant Biol.* 9, 436–442.
- Gamon, J.A., Field, C.B., Goulden, M.L., Griffin, K.L., Hartley, A.E., Joel, G., Peñuelas, J., Valentini, R., 1995. Relationships between NDVI, canopy structure, and photosynthesis in three Californian vegetation types. *Ecol. Appl.* 5, 28–41.
- García-Plazaola, J.I., Becerril, J.M., 2001. Seasonal changes in photosynthetic pigments and antioxidants in beech (*Fagus sylvatica*) in a Mediterranean climate: implications for tree decline diagnosis. *Aust. J. Plant Physiol.* 28, 225–232.

- García-Plazaola, J.I., Becerril, J.M., Hernández, A., Niinemets, Ü., Kollist, H., 2004. Acclimation of antioxidant pools to the light environment in a natural forest canopy. *New Phytol.* 163, 87–97.
- García-Plazaola, J.I., Esteban, R., Hormaetxe, K., Fernández-Marín, B., Becerril, J.M., 2008. Photoprotective responses of Mediterranean and Atlantic trees to the extreme heat-wave of summer 2003 in Southwestern Europe. *Trees* 22, 385–392.
- Gas, E., Flores-Pérez, U., Sauret-Güeto, S., Rodríguez-Concepción, M., 2009. Hunting for plant nitric oxide synthase provides new evidence of a central role for plastids in nitric oxide metabolism. *Plant Cell* 21, 18–23.
- Gayler, S., Priesack, E., 2007. PLATHO—a dynamic plant growth model considering competition between individuals and allocation to carbon-based secondary compounds. In: Fourcaud, T., Zhang, X. (Eds.), *Plant Growth Modeling and Applications*. Proceedings PMA06: The Second International Symposium on Plant Growth Modeling, Simulation, Visualization and Applications. Beijing, China, November 13–17, 2006. IEEE Computer Society, Los Alamitos, pp. 109–116.
- Génard, M., Pagès, L.C., Kervella, J., 1998. A carbon balance model of peach tree growth and development for studying the pruning response. *Tree Physiol.* 18, 351–362.
- Givnish, T.J., 1988. Adaptation to sun and shade: a whole-plant perspective. *Aust. J. Plant Physiol.* 15, 63–92.
- Gower, S.T., McMurtrie, R.E., Murty, D., 1996. Aboveground net primary production decline with stand age: potential causes. *TREE* 11, 378–382.
- Granier, A., Biron, P., Bréda, N., Pontailleur, J.Y., Saugier, B., 1996. Transpiration of trees and forest stands: short and long-term monitoring using sapflow methods. *Glob. Change Biol.* 2, 265–274.
- Gratani, L., 2000. Leaf temperature effects on gas exchange in *Quercus ilex* L. growing under field conditions. *Plant Biosyst.* 134, 19–24.
- Grime, P.J., 1989. The C-S-R model of primary plant strategies—origins, implications and tests. In: Gottlieb, L.D., Tain, S.K. (Eds.), *Plant Evolutionary Biology*. Chapman and Hall, London, New York, pp. 371–393.
- Grund, K., Conedera, M., Schröder, H., Walther, G.-R., 2005. The role of fire in the invasion process of evergreen broad-leaved species. *Basic Appl. Ecol.* 6, 47–56.
- Günthardt-Goerg, M.S., Vollenweider, P., 2007. Linking stress with macroscopic and microscopic leaf response in trees: new diagnostic perspectives. *Environ. Pollut.* 147, 467–488.
- Hájek, T., Tuittila, E.-S., Ilomets, M., Laiho, R., 2008. Light responses of mire mosses—a key to survival after water-level drawdown? *Oikos* 118, 240–250.
- Hallik, L., Niinemets, Ü., Wright, I.J., 2009. Are species shade and drought tolerance reflected in leaf-level structural and functional differentiation in Northern Hemisphere temperate woody flora? *New Phytol.* 184, 257–274.
- Hamerlynck, E.P., Huxman, T.E., Loik, M.E., Smith, S.D., 2000. Effects of extreme high temperature, drought and elevated CO<sub>2</sub> on photosynthesis of the Mojave Desert evergreen shrub, *Larrea tridentata*. *Plant Ecol.* 148, 183–193.
- Harley, P.C., Tenhunen, J.D., Beyschlag, W., Lange, O.L., 1987. Seasonal changes in net photosynthesis rates and photosynthetic capacity in leaves of *Cistus salvifolius*, a European mediterranean semi-deciduous shrub. *Oecologia* 74, 380–388.
- Havaux, M., 1992. Stress tolerance of photosystem II *in vivo*. Antagonistic effects of water, heat and photoinhibition stress. *Plant Physiol.* 100, 424–432.
- Heil, M., Silva Bueno, J.C., 2007. Within-plant signaling by volatiles leads to induction and priming of an indirect plant defense in nature. *Proc. Natl. Acad. Sci. U.S.A.* 104, 5467–5472.
- Hilker, T., Coops, N.C., Coggins, S.B., Wulder, M.A., Brown, M., Black, T.A., Nesic, Z., Lessard, D., 2009. Detection of foliage conditions and disturbance from multi-angular high spectral resolution remote sensing. *Remote Sens. Environ.* 113, 421–434.
- Hill, M.O., Mountford, J.O., Roy, D.B., Bunce, R.G.H., 1999. Ellenberg's Indicator Values for British Plants. Centre for Ecology & Hydrology, Huntingdon, Cambs.
- Hobbs, R.J., Arico, S., Aronson, J., Baron, J.S., Bridgewater, P., Cramer, V.A., Epstein, P.R., Ewel, J.J., Klink, C.A., Lugo, A.E., Norton, D., Ojima, D., Richardson, D.M., Sanders, E.W., Valladares, F., Vilà, M., Zamora, R., Zobel, M., 2006. Novel ecosystems: theoretical and management aspects of the new ecological world order. *Global Ecol. Biogeogr.* 15, 1–7.
- Hoch, G., Richter, A., Körner, C., 2003. Non-structural carbon compounds in temperate forest trees. *Plant Cell Environ.* 26, 1067–1081.
- Hubbard, R.M., Bond, B.J., Ryan, M.G., 1999. Evidence that hydraulic conductance limits photosynthesis in old *Pinus ponderosa* trees. *Tree Physiol.* 19, 165–172.
- Hüve, K., Bichele, I., Tobias, M., Niinemets, Ü., 2006. Heat sensitivity of photosynthetic electron transport varies during the day due to changes in sugars and osmotic potential. *Plant Cell Environ.* 29, 212–228.
- Ibrahim, M.A., Nissinen, A., Prozherina, N., Oksanen, E.J., Holopainen, J.K., 2006. The influence of exogenous monoterpene treatment and elevated temperature on growth, physiology, chemical content and headspace volatiles of two carrot cultivars (*Daucus carota* L.). *Environ. Exp. Bot.* 56, 95–107.
- Infante, J.M., Rambal, S., Joffre, R., 1997. Modelling transpiration in holm-oak savannah: scaling up from the leaf to the tree scale. *Agric. Forest Meteorol.* 87, 273–289.
- Innes, J.L., Boswell, R.C., 1990. Reliability, presentation, and relationships among data from inventories of forest condition. *Can. J. Forest Res.* 20, 790–799.
- Irvine, J., Law, B.E., Anthoni, P.M., Meinzer, F.C., 2002. Water limitations to carbon exchange in old-growth and young ponderosa pine stands. *Tree Physiol.* 22, 189–196.
- Isebrands, J.G., Dickson, R.E., Rebbeck, J., Karnosky, D.F., 2000. Interacting effects of multiple stresses on growth and physiological processes in northern forest trees. In: Mickler, R.A., Birdsey, R.A., Hom, J. (Eds.), *Responses of Northern U.S. Forests to Environmental Change*. Springer Verlag, New York, pp. 149–180.
- Ishida, A., Toma, T., Marjenah, 1999. Limitation of leaf carbon gain by stomatal and photochemical processes in the top canopy of *Macaranga confiera*, a tropical pioneer tree. *Tree Physiol.* 19, 467–473.
- Jackson, M.B., Ishizawa, K., Ito, O., 2009. Evolution and mechanisms of plant tolerance to flooding stress. *Ann. Bot.* 103, 137–142.
- James, S.A., Clearwater, M.J., Meinzer, F.C., Goldstein, G., 2002. Heat dissipation sensors of variable length for the measurement of sap flow in trees with deep sapwood. *Tree Physiol.* 22, 277–283.
- Janssens, I.A., Kowalski, A.S., Longdoz, B., Ceulemans, R., 2000. Assessing forest soil CO<sub>2</sub> efflux: an *in situ* comparison of four techniques. *Tree Physiol.* 20, 23–32.
- Jarvis, P.G., 1985. Increasing forest productivity and value of temperate coniferous forest by manipulating site water balance. In: *Forest Potentials*. Weyerhaeuser Science Symposium. Weyerhaeuser Co., Centralia, WA, pp. 39–74.
- Jarvis, P.G., 1995. Scaling processes and problems. *Plant Cell Environ.* 18, 1079–1089.
- Jokela, E.J., Martin, T.A., 2000. Effects of ontogeny and soil nutrient supply on production, allocation, and leaf area efficiency in loblolly and slash pine stands. *Can. J. Forest Res.* 30, 1511–1524.
- Jones, H.G., Schofield, P., 2008. Thermal and other remote sensing of plant stress. *Gen. Appl. Plant Physiol.* 34, 19–32.
- Jozsa, L.A., Brix, H., 1989. The effects of fertilization and thinning on wood quality of a 24-year-old Douglas-fir stand. *Can. J. Forest Res.* 19, 1137–1145.
- Kaelke, C.M., Kruger, E.L., Reich, P.B., 2001. Trade-offs in seedling survival, growth, and physiology among hardwood species of contrasting successional status along a light-availability gradient. *Can. J. Forest Res.* 31, 1602–1616.
- Karl, T., Guenther, A., Turnipseed, A., Patton, E.G., Jardine, K., 2008. Chemical sensing of plant stress at the ecosystem scale. *Biogeosciences* 5, 1287–1294.
- Karnosky, D.F., Pregitzer, K.S., Zak, D.R., Kubiske, M.E., Hendrey, G.R., Weinstein, D., Nosal, M., Percy, K.E., 2005. Scaling ozone responses of forest trees to the ecosystem level in a changing climate. *Plant Cell Environ.* 28, 965–981.
- Keskitalo, J., Bergquist, G., Gardestrom, P., Jansson, S., 2005. A cellular timetable of autumn senescence. *Plant Physiol.* 139, 1635–1648.
- Kittredge, D.B., Ashton, P.M.S., 1995. Impact of deer browsing on regeneration in mixed stands in southern New England. *North. J. Appl. Forest.* 12, 115–120.
- Klingeman, W.E., van Iersel, M.W., Kang, J.G., Augé, R.M., Moore, J.L., Flanagan, P.C., 2005. Whole-plant gas exchange measurements of mycorrhizal 'Iceberg' roses exposed to cyclic drought. *Crop Prot.* 24, 309–317.
- Koch, G.W., Sillett, S.C., Jennings, G.M., Davis, S.D., 2004. The limits to tree height. *Nature* 428, 851–854.
- Kolb, T.E., Fredericksen, T.S., Steiner, K.C., Skelly, J.M., 1997. Issues in scaling tree size and age responses to ozone: a review. *Environ. Pollut.* 98, 195–208.
- Körner, C., 2003. Carbon limitation in trees. *J. Ecol.* 91, 4–17.
- Koskela, J., 2000. A process-based growth model for the grass stage pine seedlings. *Silva Fenn.* 34, 3–20.
- Köstner, B., Granier, A., Chermák, J., 1998. Sapflow measurements in forest stands: methods and uncertainties. *Ann. Sci. Forest.* 55, 13–27.
- Krause, G.H., Koroleva, O.Y., Dalling, J.W., Winter, K., 2001. Acclimation of tropical tree seedlings to excessive light in simulated tree-fall gaps. *Plant Cell Environ.* 24, 1345–1352.
- Kumar, L., Skidmore, A.K., Mutanga, O., 2010. Leaf level experiments to discriminate between *Eucalyptus* species using high spectral resolution reflectance data: use of derivatives, ratios and vegetation indices. *Geocarto Int.* 25, iFirst.
- Kurz, W.A., Dymond, C.C., Stinson, G., Rampley, G.J., Neilson, E.T., Carroll, A.L., Ebata, T., Safranyik, L., 2008. Mountain pine beetle and forest carbon feedback to climate change. *Nature* 452, 987–990.
- Laisk, A., Oja, V., Rasulov, B., Rämama, H., Eichmann, H., Kasparova, I., Pettai, H., Padu, E., Vapaavuori, E., 2001. A computer-operated routine of gas exchange and optical measurements to diagnose photosynthetic apparatus in leaves. *Plant Cell Environ.* 25, 923–943.
- Labbers, H., Poorter, H., 1992. Inherent variation in growth rate between higher plants: a search for physiological causes and ecological consequences. In: Begon, M., Fitter, A.H. (Eds.), *Adv. Ecol. Res. Academic Press*, London, pp. 187–261.
- Landsberg, J.J., 1986. *Physiological Ecology of Forest Production*. Academic Press, London.
- Langley, J.A., Drake, B.G., Hungate, B.A., 2002. Extensive belowground carbon storage supports roots and mycorrhizae in regenerating scrub oaks. *Oecologia* 131, 542–548.
- Larcher, W., 1981. Low temperature effects on Mediterranean sclerophylls: an unconventional viewpoint. In: Margaris, N.S., Mooney, H.A. (Eds.), *Components of Productivity of Mediterranean-climate Regions—Basic and Applied Aspects*. Dr W. Junk Publishers, The Hague/Boston/London, pp. 259–266.
- Larcher, W., 2000. Temperature stress and survival ability of Mediterranean sclerophyllous plants. *Plant Biosyst.* 134, 279–295.
- Law, B.E., Goldstein, A.H., Anthoni, P.M., Unsworth, M.H., Panek, J.A., Bauer, M.R., Fracheboud, J.M., Hultman, N., 2001. Carbon dioxide and water vapor exchange by young and old ponderosa pine ecosystems during a dry summer. *Tree Physiol.* 21, 299–308.
- Le Roux, X., Lacointe, A., Escobar-Gutiérrez, A., Le Dizès, S., 2001. Carbon-based models of individual tree growth: a critical appraisal. *Ann. Forest Sci.* 58, 469–506.
- Le Thiec, D., Dixon, M., 1996. Acclimation of photosynthesis in Norway spruce and red oak grown in open-top chambers and subjected to natural drought and to elevated CO<sub>2</sub>. *Can. J. Forest Res.* 26, 87–94.
- Lee, S., Choi, H., Suh, S., Doo, I.-S., Oh, K.-Y., Choi, E.J., Schroeder Taylor, A.T., Low, P.S., Lee, Y., 1999. Oligogalacturonic acid and chitosan reduce stomatal aperture by inducing the evolution of reactive oxygen species from guard cells of tomato and *Commelina communis*. *Plant Physiol.* 121, 147–152.

- Leuzinger, S., Bigler, C., Wolf, A., Körner, C., 2009. Poor methodology for predicting large-scale tree die-off. *Proc. Natl. Acad. Sci. U.S.A.* 106, E106.
- Lichtenthaler, H.K., 1998. The stress concept in plants: an introduction. *Ann. N. Y. Acad. Sci.* 851, 187–198.
- Linder, S., 1987. Responses to water and nutrients in coniferous ecosystems. In: Schulze, E.D., Zwölfer, H., Golley, F., Lange, O.L., Olson, J.S., Remmert, H. (Eds.), *Potentials and Limitations of Ecosystem Analysis*. Springer-Verlag, Berlin/Heidelberg/New York/London/Paris/Tokyo, pp. 180–202.
- Lindinger, W., Hansel, A., Jordan, A., 1998. On-line monitoring of volatile organic compounds at pptv levels by means of proton-transfer-reaction mass spectrometry (PTR-MS): medical applications, food control and environmental research. *Int. J. Mass Spectrom. Ion Proc.* 173, 191–241.
- Liu, Z., Dickmann, D.I., 1993. Responses of two hybrid *Populus* clones to flooding, drought, and nitrogen availability. II. Gas exchange and water relations. *Can. J. Bot.* 71, 927–938.
- Lloyd, J., Wong, S.C., Styles, J.M., Batten, D., Priddle, R., Turnbull, C., McConchie, C.A., 1995. Measuring and modelling whole-tree gas exchange. *Aust. J. Plant Physiol.* 22, 987–1000.
- Long, S.P., Bernacchi, C.J., 2003. Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. *J. Exp. Bot.* 54, 2393–2401.
- Long, S.P., Farage, P.K., Garcia, R.L., 1996. Measurement of leaf and canopy photosynthetic CO<sub>2</sub> exchange in the field. *J. Exp. Bot.* 47, 1629–1642.
- Lopez, O.R., Kursar, T.A., 2007. Interannual variation in rainfall, drought stress and seedling mortality may mediate monodominance in tropical flooded forests. *Oecologia* 154, 35–43.
- Loreto, F., Barta, C., Brilli, F., Nogues, I., 2006. On the induction of volatile organic compound emissions by plants as consequence of wounding or fluctuations of light and temperature. *Plant Cell Environ.* 29, 1820–1828.
- Lorimer, C.G., Dahir, S.E., Nordheim, E.V., 2001. Tree mortality rates and longevity in mature and old-growth hemlock-hardwood forests. *J. Ecol.* 89, 960–971.
- Loughrin, J.H., Potter, D.A., Hamilton-Kemp, T., Byers, M.E., 1997. Diurnal emission of volatile compounds by Japanese beetle-damaged grape leaves. *Phytochemistry* 45, 919–923.
- Löw, M., Herbinger, K., Nunn, A.J., Häberle, K.-H., Leuchner, M., Heerdt, C., Werner, H., Wipfler, P., Pretzsch, H., Tausz, M., Matyssek, R., 2006. Extraordinary drought of 2003 overrules ozone impact on adult beech trees (*Fagus sylvatica*). *Trees* 20, 539–548.
- Lusk, C.H., Del, P.A., 2002. Survival and growth of seedlings of 12 Chilean rainforest trees in two light environments: gas exchange and biomass distribution correlates. *Austral Ecol.* 27, 173–182.
- Mandujano-Chávez, A., Schoenbeck, M.A., Ralston, L.F., Lozoya-Gloria, E., Chappell, J., 2000. Differential induction of sesquiterpene metabolism in tobacco cell suspension cultures by methyl jasmonate and fungal elicitor. *Arch. Biochem. Biophys.* 381, 285–294.
- Manion, P.D., 2003. Evolution of concepts in forest pathology. *Phytopathology* 93, 1052–1055.
- Maroco, J.P., Breia, E., Faria, T., Pereira, J.S., Chaves, M.M., 2002. Effects of long-term exposure to elevated CO<sub>2</sub> and N fertilization on the development of photosynthetic capacity and biomass accumulation in *Quercus suber* L. *Plant Cell Environ.* 25, 105–113.
- Marschner, H., Kirkby, E.A., Engels, C., 1997. Importance of cycling and recycling of mineral nutrients within plants for growth and development. *Bot. Acta* 110, 265–273.
- Martin, D., Gershenzon, J., Bohlmann, J., 2003. Induction of volatile terpene biosynthesis and diurnal emission by methyl jasmonate in foliage of Norway spruce. *Plant Physiol.* 132, 1586–1599.
- Matsui, K., 2006. Green leaf volatiles: hydroperoxide lyase pathway of oxylipin metabolism. *Curr. Opin. Plant Biol.* 9, 274–280.
- Mattson, W.J., Haack, R.A., 1987. The role of drought in outbreaks of plant-eating insects. *BioScience* 37, 110–118.
- Matyssek, R., Sandermann, H., Wieser, G., Booker, F., Cieslik, S., Musselman, R., Ernst, D., 2008. The challenge of making ozone risk assessment for forest trees more mechanistic. *Environ. Pollut.* 156, 567–582.
- Maxwell, K., Johnson, G.N., 2000. Chlorophyll fluorescence—a practical guide. *J. Exp. Bot.* 51, 659–668.
- McDonald, E.P., Agrell, J., Lindroth, R.L., 1999. CO<sub>2</sub> and light effects on deciduous trees: growth, foliar chemistry, and insect performance. *Oecologia* 119, 389–399.
- McDowell, N., Pockman, W.T., Allen, C.D., Breshears, D.D., Cobb, N., Kolb, T., Plaut, J., Sperry, J., West, A., Williams, D.G., Yepez, E.A., 2008. Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought? *New Phytol.* 178, 719–739.
- McDowell, N.G., Sevanto, S., 2010. The mechanisms of carbon starvation: how, when, or does it even occur at all? *New Phytol.* 186, 264–266.
- McLaughlin, D., 1998. A decade of forest tree monitoring in Canada: evidence of air pollution effects. *Environ. Rev.* 6, 151–171.
- Medhurst, J., Parsby, J., Linder, S., Wallin, G., Ceschia, E., Slaney, M., 2006. A whole-tree chamber system for examining tree-level physiological responses of field-grown trees to environmental variation and climate change. *Plant Cell Environ.* 29, 1853–1869.
- Meehl, G.A., Stocker, T.F., Collins, W.D., Friedlingstein, P., Gaye, A.T., Gregory, J.M., Kitoh, A., Knutti, R., Murphy, J.M., Noda, A., Raper, S.C.B., Watterson, I.G., Weaver, A.J., Zhao, Z.-C., 2007. Global climate projections. In: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (Eds.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, pp. 747–845.
- Mendel, Z., Assaell, F., Saphir, N., Zehavi, A., Nestel, D., Schiller, G., 1997. Seedling mortality in regeneration of Aleppo pine following fire and attack by the scale insect *Matsucoccus josephi*. *Int. J. Wildland Fire* 7, 327–333.
- Messier, C., Nikinmaa, E., 2000. Effects of light availability and sapling size on the growth, biomass allocation, and crown morphology of understory sugar maple, yellow birch, and beech. *Coscience* 7, 345–356.
- Micheli, F., 2001. Pectin methylesterases: cell wall enzymes with important roles in plant physiology. *Trends Plant Sci.* 6, 414–419.
- Mittler, R., 2006. Abiotic stress, the field environment and stress combination. *Trends Plant Sci.* 11, 15–19.
- Mittler, R., Merquiol, E., Hallak-Herr, E., Rachmilevitch, S., Kaplan, A., Cohen, M., 2001. Living under a 'dormant' canopy: a molecular acclimation mechanism of the desert plant *Retama retam*. *Plant J.* 25, 407–416.
- Mohan, J.E., Ziska, L.H., Schlesinger, W.H., Thomas, R.B., Sicher, R.C., George, K., Clark, J.S., 2006. Biomass and toxicity responses of poison ivy (*Toxicodendron radicans*) to elevated atmospheric CO<sub>2</sub>. *Proc. Natl. Acad. Sci. U.S.A.* 103, 9086–9089.
- Mueller, R.C., Scudder, C.M., Porter, M.E., Trotter III, R.T., Gehring, C.A., Whitham, T.G., 2005. Differential tree mortality in response to severe drought: evidence for long-term vegetation shifts. *J. Ecol.* 93, 1085–1093.
- Mullin, L.P., Sillett, S.C., Koch, G.W., Tu, K.P., Antoine, M.E., 2009. Physiological consequences of height-related morphological variation in *Sequoia sempervirens* foliage. *Tree Physiol.* 29, 999–1010.
- Murray, K.J., Tenhunen, J.D., Kummerow, J., 1989. Limitations on *Sphagnum* growth and net primary production in the foothills of the Philip Smith Mountains, Alaska. *Oecologia* 80, 256–262.
- Murray, K.J., Tenhunen, J.D., Nowak, R.S., 1993. Photoinhibition as a control on photosynthesis and production of *Sphagnum* mosses. *Oecologia* 96, 200–207.
- Murty, D., McMurtrie, R.E., Ryan, M.G., 1996. Declining forest productivity in aging forest stands: a modeling analysis of alternative hypotheses. *Tree Physiol.* 16, 187–200.
- Myers, J.A., 2005. Seedling Carbohydrate Storage, Survival and Stress Tolerance in a Neotropical Forest. MSc. Thesis Department of Botany, University of Florida, Gainesville, FL.
- Myers, J.A., Kitajima, K., 2007. Carbohydrate storage enhances seedling shade and stress tolerance in a neotropical forest. *J. Ecol.* 95, 383–395.
- Nadezhkina, N., 1999. Sap flow index as an indicator of plant water status. *Tree Physiol.* 19, 885–891.
- Niinemets, Ü., 1997a. Distribution patterns of foliar carbon and nitrogen as affected by tree dimensions and relative light conditions in the canopy of *Picea abies*. *Trees* 11, 144–154.
- Niinemets, Ü., 1997b. Energy requirement for foliage construction depends on tree size in young *Picea abies* trees. *Trees* 11, 420–431.
- Niinemets, Ü., 1998. Growth of young trees of *Acer platanoides* and *Quercus robur* along a gap–understory continuum: interrelationships between allometry, biomass partitioning, nitrogen, and shade-tolerance. *Int. J. Plant Sci.* 159, 318–330.
- Niinemets, Ü., 2002. Stomatal conductance alone does not explain the decline in foliar photosynthetic rates with increasing tree age and size in *Picea abies* and *Pinus sylvestris*. *Tree Physiol.* 22, 515–535.
- Niinemets, Ü., 2007. Photosynthesis and resource distribution through plant canopies. *Plant Cell Environ.* 30, 1052–1071.
- Niinemets, Ü., 2010a. A review of light interception in plant stands from leaf to canopy in different plant functional types and in species with varying shade tolerance. *Ecol. Res.* 25, 693–714.
- Niinemets, Ü., 2010b. Mild versus severe stress and BVOCs: thresholds, priming and consequences. *Trends Plant Sci.* 15, 145–153.
- Niinemets, Ü., Ellsworth, D.S., Lukjanova, A., Tobias, M., 2001. Site fertility and the morphological and photosynthetic acclimation of *Pinus sylvestris* needles to light. *Tree Physiol.* 21, 1231–1244.
- Niinemets, Ü., Kollist, H., García-Plazaola, J.I., Hernández, A., Becerril, J.M., 2003. Do the capacity and kinetics for modification of xanthophyll cycle pool size depend on growth irradiance in temperate trees? *Plant Cell Environ.* 26, 1787–1801.
- Niinemets, Ü., Kull, K., 2005. Co-limitation of plant primary productivity by nitrogen and phosphorus in a species-rich wooded meadow on calcareous soils. *Acta Oecol.* 28, 345–356.
- Niinemets, Ü., Kull, O., 1995. Effects of light availability and tree size on the architecture of assimilative surface in the canopy of *Picea abies*: variation in needle morphology. *Tree Physiol.* 15, 307–315.
- Niinemets, Ü., Kull, O., 2001. Sensitivity to photoinhibition of photosynthetic electron transport in a temperate deciduous forest canopy: photosystem II centre openness, non-radiative energy dissipation and excess irradiance under field conditions. *Tree Physiol.* 21, 899–914.
- Niinemets, Ü., Oja, V., Kull, O., 1999a. Shape of leaf photosynthetic electron transport versus temperature response curve is not constant along canopy light gradients in temperate deciduous trees. *Plant Cell Environ.* 22, 1497–1514.
- Niinemets, Ü., Söber, A., Kull, O., Hartung, W., Tenhunen, J.D., 1999b. Apparent controls on leaf conductance by soil water availability and via light-acclimation of foliage structural and physiological properties in a mixed deciduous, temperate forest. *Int. J. Plant Sci.* 160, 707–721.
- Niinemets, Ü., Valladares, F., 2004. Photosynthetic acclimation to simultaneous and interacting environmental stresses along natural light gradients: optimality and constraints. *Plant Biol.* 6, 254–268.

- Niinemets, Ü., Valladares, F., 2006. Tolerance to shade, drought and waterlogging in the temperate dendroflora of the Northern hemisphere: tradeoffs, phylogenetic signal and implications for niche differentiation. *Ecol. Monogr.* 76, 521–547.
- Niinemets, Ü., Valladares, F., 2008. Environmental tolerance. In: Jørgensen, S.E., Fath, B.D. (Eds.), *Encyclopedia of Ecology*. Elsevier, Oxford, pp. 1370–1376.
- Nunn, A.J., Kozovits, A.R., Reiter, I.M., Heerdt, C., Leuchner, M., Lütz, C., Liu, X., Löw, M., Winkler, J.B., Grams, T.E.E., Häberle, K.-H., Werner, H., Fabian, P., Rennenberg, H., Matyssek, R., 2005a. Comparison of ozone uptake and sensitivity between a phytotron study with young beech and a field experiment with adult beech (*Fagus sylvatica*). *Environ. Pollut.* 137, 494–506.
- Nunn, A.J., Reiter, I.M., Häberle, K.-H., Langebartsels, C., Bahnweg, G., Pretzsch, H., Sandermann, H., Matyssek, R., 2005b. Response patterns in adult forest trees to chronic ozone stress: identification of variations and consistencies. *Environ. Pollut.* 136, 365–369.
- O'Hanlon-Manners, D.L., Kotanen, P.M., 2004. Evidence that fungal pathogens inhibit recruitment of a shade-intolerant tree, white birch (*Betula papyrifera*), in understory habitats. *Oecologia* 140, 650–653.
- Ögren, E., 1996. Premature dehardening in *Vaccinium myrtillus* during a mild winter: a cause for winter dieback? *Funct. Ecol.* 10, 724–732.
- Ögren, E., 1997. Relationship between temperature, respiratory loss of sugar and premature dehardening in dormant Scots pine seedlings. *Tree Physiol.* 17, 47–51.
- Ollinger, S.V., Aber, J.D., Reich, P.B., 1997. Simulating ozone effects on forest productivity: interactions among leaf-, canopy-, and stand-level processes. *Ecol. Appl.* 7, 1237–1251.
- Opitz, S., Kunert, G., Gershenzon, J., 2008. Increased terpenoid accumulation in cotton (*Gossypium hirsutum*) foliage is a general wound response. *J. Chem. Ecol.* 34, 508–522.
- Padilla, F.M., de Dios Miranda, J., Pugnaire, F.I., 2007. Early root growth plasticity in seedlings of three Mediterranean woody species. *Plant Soil* 296, 103–113.
- Palahi, M., Pukkala, T., Miina, J., Montero, G., 2003. Individual-tree growth and mortality models for Scots pine (*Pinus sylvestris* L.) in north-east Spain. *Ann. Forest Sci.* 60, 1–10.
- Paoletti, E., Schaub, M., Matyssek, R., Wieser, G., Augustaitis, A., Bastrup-Birk, A.M., Bytnerowicz, A., Günthardt-Goerg, M.S., Müller-Starck, G., Serengil, Y., 2010. Advances of air pollution science: from forest decline to multiple-stress effects on forest ecosystem services. *Environ. Pollut.* 158, 1986–1989.
- Pataki, D.E., Oren, R., Katul, G., Sigmon, J., 1998. Canopy conductance of *Pinus taeda*, *Liquidambar styraciflua* and *Quercus phellos* under varying atmospheric and soil water conditions. *Tree Physiol.* 18, 307–315.
- Pell, E.J., Temple, P.J., Friend, A.L., Mooney, H.A., Winner, W.E., 1994. Compensation as a plant response to ozone and associated stresses: an analysis of ROPIS experiments. *J. Environ. Qual.* 23, 429–436.
- Pelloux, J., Jolivet, Y., Fontaine, V., Banvoy, J., Dizengremel, P., 2001. Changes in Rubisco and Rubisco activase gene expression and polypeptide content in *Pinus halepensis* M. subjected to ozone and drought. *Plant Cell Environ.* 24, 123–131.
- Peñuelas, J., Boada, M., 2003. A global change-induced biome shift in the Motesny mountains (NE Spain). *Glob. Change Biol.* 9, 131–140.
- Peñuelas, J., Ogaya, R., Boada, M., Jump, A.S., 2007. Migration, invasion and decline: changes in recruitment and forest structure in a warming-linked shift of European beech forest in Catalonia (NE Spain). *Ecography* 30, 829–837.
- Pichersky, E., Gershenzon, J., 2002. The formation and function of plant volatiles: perfumes for pollinator attraction and defense. *Curr. Opin. Plant Biol.* 5, 237–243.
- Polley, H.W., Tischler, C.R., Johnson, H.B., Derner, J.D., 2002. Growth rate and survivorship of drought: CO<sub>2</sub> effects on the presumed tradeoff in seedlings of five woody legumes. *Tree Physiol.* 22, 383–391.
- Poorter, L., Hayashida-Oliver, Y., 2000. Effects of seasonal drought on gap and understorey seedlings in a Bolivian moist forest. *J. Trop. Ecol.* 16, 481–498.
- Poorter, L., Kitajima, K., 2007. Carbohydrate storage and light requirements of tropical moist and dry forest species. *Ecology* 88, 1000–1011.
- Porder, S., Asner, G.P., Vitousek, P.M., 2005. Ground-based and remotely sensed nutrient availability across a tropical landscape. *Proc. Natl. Acad. Sci. U.S.A.* 102, 10909–10912.
- Porta, H., Rocha-Sosa, M., 2002. Plant lipoxygenases. Physiological and molecular features. *Plant Physiol.* 130, 15–21.
- Portsmouth, A., Niinemets, Ü., Truus, L., Pensa, M., 2005. Biomass allocation and growth rates in *Pinus sylvestris* are interactively modified by nitrogen and phosphorus availabilities and by tree size and age. *Can. J. Forest Res.* 35, 2346–2359.
- Prats, E., Carver, T.L.W., Gay, A.P., Mur, L.A.J., 2007. Enemy at the gates. Interaction-specific stomatal responses to pathogenic challenge. *Plant Signal. Behav.* 2, 275–277.
- Rambal, S., 2001. Productivity of Mediterranean-type ecosystems. In: Mooney, H.A., Saugier, B., Roy, J. (Eds.), *Terrestrial Global Productivity: Past, Present, and Future*. Academic Press, Inc., San Diego, pp. 315–344.
- Rennenberg, H., Loreto, F., Polle, A., Brilli, F., Fares, S., Beniwall, R.S., Gessler, A., 2006. Physiological responses of forest trees to heat and drought. *Plant Biol.* 8, 556–571.
- Retzlaff, W.A., Weinstein, D.A., Laurence, J.A., Gollands, B., 1997. Simulating the growth of a 160-year-old sugar maple (*Acer saccharum*) tree with and without ozone exposure using the TREGRO model. *Can. J. Forest Res.* 27, 783–789.
- Richardson, A.D., Aikens, M., Berlyn, G.P., Marshall, P., 2004. Drought stress and paper birch (*Betula papyrifera*) seedlings: effects of an organic biostimulant on plant health and stress tolerance, and detection of stress effects with instrument-based, noninvasive methods. *J. Arboric.* 30, 52–61.
- Riikonen, J., Mäenpää, M., Alavillamo, M., Silfver, T., Oksanen, E., 2009. Interactive effect of elevated temperature and O<sub>3</sub> on antioxidant capacity and gas exchange in *Betula pendula* saplings. *Planta* 230, 419–427.
- Robertz, P., Stockfors, J., 1998. Effects of elevated CO<sub>2</sub> concentration and nutrition on net photosynthesis, stomatal conductance and needle respiration of field-grown Norway spruce trees. *Tree Physiol.* 18, 233–241.
- Roden, J.S., Ball, M.C., 1996. The effect of elevated [CO<sub>2</sub>] on growth and photosynthesis of two *Eucalyptus* species exposed to high temperatures and water deficits. *Plant Physiol.* 111, 909–919.
- Rosenqvist, E., Wingsle, G., Ögren, E., 1991. Photoinhibition of photosynthesis in intact willow leaves in response to moderate changes in light and temperature. *Physiol. Plant.* 83, 390–396.
- Roth, S., McDonald, E.P., Lindroth, R.L., 1997. Atmospheric CO<sub>2</sub> and soil water availability: consequences for tree-insect interactions. *Can. J. Forest Res.* 27, 1281–1290.
- Rothe, M., Krabel, D., Roloff, A., 1999. Responses of Norway spruce (*Picea abies* [L.] Karst.) to water deficiency. *Phyton-Ann. Rei Bot.* 39, 183–190.
- Rütten, D., Santarius, K.A., 1988. Cold acclimation of *Ilex aquifolium* under natural conditions with special regard to the photosynthetic apparatus. *Physiol. Plant.* 72, 807–815.
- Ryan, M.G., Yoder, B.J., 1997. Hydraulic limits to tree height and tree growth. What keeps trees from growing beyond a certain height? *BioScience* 47, 235–242.
- Sack, L., Grubb, P.J., 2001. Why do species of woody seedlings change rank in relative growth rate between low and high irradiance? *Funct. Ecol.* 15, 145–154.
- Sala, A., 2009. Lack of direct evidence for the carbon-starvation hypothesis to explain drought-induced mortality in trees. *Proc. Natl. Acad. Sci. U.S.A.* 106, E68.
- Sala, A., Hoch, G., 2009. Height-related growth declines in ponderosa pine are not due to carbon limitation. *Plant Cell Environ.* 32, 22–30.
- Sala, A., Piper, F., Hoch, G., 2010. Physiological mechanisms of drought-induced tree mortality are far from being resolved. *New Phytol.* 186, 274–281.
- Sampson, D.A., Allen, H.L., 1998. Light attenuation in a 14-year-old loblolly pine stand as influenced by fertilization and irrigation. *Trees* 13, 80–87.
- Samson, G., Tremblay, N., Dudelzak, A.E., Babichenko, S.M., Dextraze, L., Wollring, J., 2000. Nutrient stress of corn plants: early detection and discrimination using a compact multiwavelength fluorescent LIDAR. *EARSeL eProceedings* 1, 214–223.
- Sánchez-Gómez, D., Valladares, F., Zavala, M.A., 2006. Functional traits and plasticity underlying shade tolerance in seedlings of four Iberian forest tree species. *Tree Physiol.* 26, 1425–1433.
- Santarius, K.A., 1990. Freezing of isolated thylakoid membranes in complex media. V. Inactivation and protection of electron transport reactions. *Photosynth. Res.* 23, 49–58.
- Saur, E., Nambiar, E.K.S., Fife, D.N., 2000. Foliar nutrient retranslocation in *Eucalyptus globulus*. *Tree Physiol.* 20, 1105–1112.
- Saxe, H., Ellsworth, D.S., Heath, J., 1998. Tansley review no. 98. Tree and forest functioning in an enriched CO<sub>2</sub> atmosphere. *New Phytol.* 139, 395–436.
- Schaeppman-Strub, G., Schaeppman, M.E., Painter, T.H., Dangel, S., Martonchik, J.V., 2006. Reflectance quantities in optical remote sensing—definitions and case studies. *Remote Sens. Environ.* 103, 27–42.
- Schleppi, P., Muller, N., Edwards, P.J., Bucher, J.B., 1999. Three years of increased nitrogen deposition do not affect the vegetation of a montane forest ecosystem. *Phyton-Ann. Rei Bot.* 39, 197–204.
- Schmid-Haas, P., 1993. Crown transparency and mortality rates in spruce, fir and beech. *Forstwiss. Centralbl.* 112, 325–333.
- Schoeneweiss, D.F., 1981a. Infectious diseases of trees associated with water and freezing stress. *J. Arboric.* 7, 13–18.
- Schoeneweiss, D.F., 1981b. The role of environmental stress in diseases of woody plants. *Plant Dis.* 56, 308–314.
- Schoeneweiss, D.F., 1983. Drought predisposition to *Cytospora* canker in blue spruce. *Plant Dis.* 67, 383–385.
- Schwanz, P., Polle, A., 2001. Differential stress responses of antioxidative systems to drought in pendunculate oak (*Quercus robur*) and maritime pine (*Pinus pinaster*) grown under high CO<sub>2</sub> concentrations. *J. Exp. Bot.* 52, 133–143.
- Seemann, J.R., Downton, W.J.S., Berry, J.A., 1986. Temperature and leaf osmotic potential as factors in the acclimation of photosynthesis to high temperature in desert plants. *Plant Physiol.* 80, 926–930.
- Small, E., 1972. Photosynthetic rates in relation to nitrogen recycling as an adaptation to nutrient deficiency in peat bog plants. *Can. J. Bot.* 50, 2227–2233.
- Smith, T., Huston, M., 1989. A theory of the spatial and temporal dynamics of plant communities. *Vegetatio* 83, 49–69.
- Solberg, S., 1999. Crown condition and growth relationships within stands of *Picea abies*. *Scand. J. Forest Res.* 14, 320–327.
- Staudt, M., Bertin, N., 1998. Light and temperature dependence of the emission of cyclic and acyclic monoterpenes from holm oak (*Quercus ilex* L.) leaves. *Plant Cell Environ.* 21, 385–395.
- Steindel, F., Beauchamp, J., Hansel, A., Kesselmeier, J., Kleist, E., Kuhn, U., Wisthaler, A., Wildt, J., 2005. Stress induced VOC emissions from mildew infested oak. *Geophys. Res. Abstr.* 7, EGU05-A-03010.
- Steppe, K., Niinemets, Ü., Teskey, R.O., 2011. Tree size- and age-related changes in leaf physiology and their influence on carbon gain. In: Meinzer, F.C., Dawson, T., Lachenbruch, B. (Eds.), *Size- and Age-related Changes in Tree Structure and Function*. Springer, Berlin, pp. xx–xx.
- Stevens, M.T., Lindroth, R.L., 2005. Induced resistance in the indeterminate growth of aspen (*Populus tremuloides*). *Oecologia* 145, 298–306.
- Stewart, J.D., Zine El-Abidine, A., Bernier, P.Y., 1995. Stomatal and mesophyll limitations of photosynthesis in black spruce seedlings during multiple cycles of drought. *Tree Physiol.* 15, 57–64.
- Stylinski, C.D., Gamon, J.A., Oechel, W.C., 2002. Seasonal patterns of reflectance indices, carotenoid pigments and photosynthesis of evergreen chaparral species. *Oecologia* 131, 366–374.

- Taub, D.R., Seemann, J.R., Coleman, J.S., 2000. Growth in elevated CO<sub>2</sub> protects photosynthesis against high-temperature damage. *Plant Cell Environ.* 23, 649–656.
- Tausz, M., Bytnerowicz, A., Arbaugh, M.J., Wonisch, A., Grill, D., 2001. Multivariate patterns of biochemical responses of *Pinus ponderosa* trees at field plots in the San Bernardino Mountains, southern California. *Tree Physiol.* 21, 329–336.
- Tausz, M., Grill, D., Stabentheiner, E., Kromp-Kolb, H., Stohl, A., Führer, E., Mitterböck, F., 1997. Comparative assessment of forest condition by visual criteria and physiological stress-indicators as affected by ambient air pollution and meteorology. *Acta Biol.* 19, 1–19.
- Tenhunen, J.D., Falge, E., Ryel, R., Manderscheid, B., Peters, K., Ostendorf, B., Joss, U., Lischheid, G., 2001. Modelling of fluxes in a spruce forest catchment of the Fichtelgebirge. In: *Ecosystem Approaches to Landscape Management in Central Europe*. Springer-Verlag Berlin, Berlin, pp. 417–462.
- Tjoelker, M.G., Oleksyn, J., Lorenc-Plucinska, G., Reich, P.B., 2008. Acclimation of respiratory temperature responses in northern and southern populations of *Pinus banksiana*. *New Phytol.* 181, 218–229.
- Tjoelker, M.G., Oleksyn, J., Reich, P.B., 1998. Seedlings of five boreal tree species differ in acclimation of net photosynthesis to elevated CO<sub>2</sub> and temperature. *Tree Physiol.* 18, 715–726.
- Tobiessen, P., Rundel, P.W., Stecker, R.E., 1971. Water potential gradient in a tall *Sequoiadendron*. *Plant Physiol.* 48, 303–304.
- Tognetti, R., Johnson, J.D., Michelozzi, M., Raschi, A., 1998. Response of foliar metabolism in mature trees of *Quercus pubescens* and *Quercus ilex* to long-term elevated CO<sub>2</sub>. *Environ. Exp. Bot.* 39, 233–245.
- Tomlinson, P.T., Anderson, P.D., 1998. Ontogeny affects response of northern red oak seedlings to elevated CO<sub>2</sub> and water stress. II. Recent photosynthate distribution and growth. *New Phytol.* 140, 493–504.
- Tømmervik, H., Odaas, A.M., Skevik, K., 1994. Measurements of chlorophyll fluorescence in several plant species in different air pollution affected sites in Varanger, Northern Norway, compared with remote sensing data. In: *Geoscience and Remote Sensing Symposium, 1994. IGARSS '94. Surface and Atmospheric Remote Sensing: Technologies, Data Analysis and Interpretation*, p. 990–992.
- Trenberth, K.E., Jones, P.D., Ambenje, P., Bojariu, R., Easterling, D., Klein Tank, A., Parker, D., Rahimzadeh, F., Renwick, J.A., Rusticucci, M., Soden, B., Zhai, P., 2007. Observations: surface and atmospheric climate change. In: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (Eds.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, pp. 235–336.
- Troeng, E., Linder, S., 1982. Gas exchange in a 20-year-old stand of Scots pine. I. Net photosynthesis of current and one-year-old shoots within and between seasons. *Physiol. Plant.* 54, 7–14.
- Tschaplinski, T.J., Stewart, D.B., Norby, R.J., 1995. Interactions between drought and elevated CO<sub>2</sub> on osmotic adjustment and solute concentrations of tree seedlings. *New Phytol.* 131, 169–177.
- Unger, S., Máguas, C., Pereira, J.S., Aires, L.M., David, T.S., Werner, C., 2009. Partitioning carbon fluxes in a Mediterranean oak forest to disentangle changes in ecosystem sink strength during drought. *Agric. Forest Meteorol.* 149, 949–961.
- Valladares, F., Chico, J.M., Aranda, I., Balaguer, L., Dizengremel, P., Manrique, E., Dreyer, E., 2002. The greater seedling high-light tolerance of *Quercus robur* over *Fagus sylvatica* is linked to a greater physiological plasticity. *Trees* 16, 395–403.
- Valladares, F., Dobarro, I., Sánchez-Gómez, D., Pearcy, R.W., 2005. Photoinhibition and drought in Mediterranean woody saplings: scaling effects and interactions in sun and shade phenotypes. *J. Exp. Bot.* 56, 483–494.
- Valladares, F., Niinemets, Ü., 2007. The architecture of plant crowns: from design rules to light capture and performance. In: Pugnaire, F.I., Valladares, F. (Eds.), *Handbook of Functional Plant Ecology*. CRC Press, Boca Raton, pp. 101–149.
- Valladares, F., Niinemets, Ü., 2008. Shade tolerance, a key plant feature of complex nature and consequences. *Annu. Rev. Ecol. Evol. Syst.* 39, 237–257.
- Valladares, F., Pearcy, R.W., 1997. Interactions between water stress, sun-shade acclimation, heat tolerance and photoinhibition in the sclerophyll *Heteromeles arbutifolia*. *Plant Cell Environ.* 20, 25–36.
- Valladares, F., Pearcy, R.W., 2002. Drought can be more critical in the shade than in the sun: a field study of carbon gain and photo-inhibition in a Californian shrub during a dry El Niño year. *Plant Cell Environ.* 25, 749–759.
- Valladares, F., Zaragoza-Castells, J., Sánchez-Gómez, D., Matesanz, S., Alonso, B., Portsmouth, A., Delgado, A., Atkin, O.K., 2008. Is shade beneficial for Mediterranean shrubs experiencing periods of extreme drought and late-winter frosts? *Ann. Bot.* 102, 923–933.
- van Bloem, S.J., Murphy, P.G., Lugo, A.E., 2003. Subtropical dry forest trees with no apparent damage sprout following a hurricane. *Trop. Ecol.* 44, 137–145.
- van Mantgem, P.J., Stephenson, N.L., Byrne, J.C., Daniels, L.D., Franklin, J.F., Fulé, P.Z., Harmon, M.E., Larson, A.J., Smith, J.M., Taylor, A.H., Veblen, T.T., 2009. Widespread increase of tree mortality rates in the western United States. *Science* 323, 521–524.
- Vargas, R., 2009. On the fate of old stored carbon after large-infrequent disturbances in plants. *Plant Signal. Behav.* 4, 617–619.
- Vargas, R., Trumbore, S.E., Allen, M.F., 2009. Evidence of old carbon used to grow new fine roots in a tropical forest. *New Phytol.* 182, 710–718.
- Verdaguer, D., Ojeda, F., 2002. Root starch storage and allocation patterns in seeder and resprouter seedlings of two Cape *Erica* (Ericaceae) species. *Am. J. Bot.* 89, 1189–1196.
- Vogel, M., Peschke, G., Rulicke, A., 1996. Calculation and interpretation of stem storage changes of a 100-year-old spruce. *Allg. Forst Jagdztg.* 167, 162–167.
- Vollenweider, P., Günthardt-Goerg, M.S., 2005. Diagnosis of abiotic and biotic stress factors using the visible symptoms in foliage. *Environ. Pollut.* 137, 455–465.
- Vranová, E., Inzé, D., Van Breusegem, F., 2002. Signal transduction during oxidative stress. *J. Exp. Bot.* 53, 1227–1236.
- Vu, J.C.V., Newman, Y.C., Allen Jr., L.H., Gallo-Meagher, M., Zhang, M.-Q., 2002. Photosynthetic acclimation of young sweet orange trees to elevated growth CO<sub>2</sub> and temperature. *J. Plant Physiol.* 159, 147–157.
- Wagner, P.A., Dreyer, E., 1997. Interactive effects of waterlogging and irradiance on the photosynthetic performance of seedlings from three oak species displaying different sensitivities (*Quercus robur*, *Q. petraea* and *Q. rubra*). *Ann. Sci. Forest.* 54, 409–429.
- Wallin, G., Linder, S., Lindroth, A., Rantfors, M., Flemberg, S., Grelle, A., 2001. Carbon dioxide exchange in Norway spruce at the shoot, tree and ecosystem scale. *Tree Physiol.* 21, 969–976.
- Walter, D.E., 1992. Leaf surface structure and the distribution of *Phytoseius* mites (*Acarina: Phytoseiidae*) in south-eastern Australian forests. *Aust. J. Zool.* 40, 593–603.
- Walters, G.M., McCarthy, B.C., 1997. Forest decline and tree mortality in a South-eastern Ohio oak-hickory forest. *Ohio J. Sci.* 97, 5–9.
- Walters, M.B., Reich, P.B., 1996. Are shade tolerance, survival, and growth linked? Low light and nitrogen effects on hardwood seedlings. *Ecology* 77, 841–853.
- Walters, M.B., Reich, P.B., 1997. Growth of *Acer saccharum* seedlings in deeply shaded understories of northern Wisconsin: effects of nitrogen and water availability. *Can. J. Forest Res.* 27, 237–247.
- Walters, M.B., Reich, P.B., 1999. Research review. Low-light carbon balance and shade tolerance in the seedlings of woody plants: do winter deciduous and broad-leaved evergreen species differ? *New Phytol.* 143, 143–154.
- Walters, M.B., Reich, P.B., 2000. Trade-offs in low-light CO<sub>2</sub> exchange: a component of variation in shade tolerance among cold temperate tree seedlings. *Funct. Ecol.* 14, 155–165.
- Walther, G.-R., Berger, S., Sykes, M.T., 2005. An ecological 'footprint' of climate change. *Proc. R. Soc. Lond. Ser. B* 272, 1427–1432.
- Wargo, P.M., 1996. Consequences of environmental stress on oak: predisposition to pathogens. *Ann. Sci. Forest.* 53, 359–368.
- Warton, D.I., Wright, I.J., Falster, D.S., Westoby, M., 2006. Bivariate line-fitting methods for allometry. *Biol. Rev. Camb. Philos. Soc.* 81, 259–291.
- Wayne, P.M., Reekie, E.G., Bazzaz, F.A., 1998. Elevated CO<sub>2</sub> ameliorates birch response to high temperature and frost stress: implications for modeling climate-induced geographic range shifts. *Oecologia* 114, 335–342.
- Weikert, R.M., Wedler, M., Lippert, M., Schramel, P., Lange, O.L., 1989. Photosynthetic performance, chloroplast pigments, and mineral content of various needle age classes of spruce (*Picea abies*) with and without the new flush: an experimental approach for analysing forest decline phenomena. *Trees* 3, 161–172.
- Weinstein, D.A., Yanai, R.D., 1994. Integrating the effects of simultaneous multiple stresses on plants using the simulation model TREGRO. *J. Environ. Qual.* 23, 418–428.
- Werner, C., Ryel, R.J., Correia, O., Beyschlag, W., 2001. Effects of photoinhibition on whole-plant carbon gain assessed with a photosynthesis model. *Plant Cell Environ.* 24, 27–40.
- Wieser, G., Hecke, K., Tausz, M., Häberle, K.-H., Grams, T.E.E., Matussek, R., 2003. The influence of microclimate and tree age on the defense capacity of European beech (*Fagus sylvatica* L.) against oxidative stress. *Ann. Forest Sci.* 60, 131–135.
- Wieser, G., Tausz, M., Wonisch, A., Havranek, W.M., 2001. Free radical scavengers and photosynthetic pigments in *Pinus cembra* L. needles as affected by ozone exposure. *Biol. Plant.* 44, 225–232.
- Williams, J.W., Jackson, S.T., 2007. Novel climates, no-analog communities, and ecological surprises. *Front. Ecol. Environ.* 5, 475–482.
- Wise, R.R., 1995. Chilling-enhanced photooxidation: the production, action and study of reactive oxygen species produced during chilling in the light. *Photosynth. Res.* 45, 79–97.
- Wong, B.L., Staats, L.J., Burfeind, A.S., Baggett, K.L., Rye, A.H., 2005. Carbohydrate reserves in *Acer saccharum* trees damaged during the January 1998 ice storm in northern New York. *Can. J. Bot.* 83, 668–677.
- Wood, B.W., Reilly, J.C., Cottrell, T., Tedders, W.L., Yates, I., 2000. Bronzing of pecan foliage. *J. Am. Soc. Hort. Sci.* 125, 41–46.
- Woodruff, D.R., McCulloh, K.A., Warren, J.M., Meinzer, F.C., Lachenbruch, B., 2007. Impacts of tree height on leaf hydraulic architecture and stomatal control in Douglas-fir. *Plant Cell Environ.* 30, 559–569.
- Woodruff, D.R., Meinzer, F.C., Lachenbruch, B., Johnson, D.M., 2009. Coordination of leaf structure and gas exchange along a height gradient in a tall conifer. *Tree Physiol.* 29, 261–272.
- Wullschlegel, S.D., Norby, R.J., Hendrix, D.L., 1992. Carbon exchange rates, chlorophyll content, and carbohydrate status of two forest tree species exposed to carbon dioxide enrichment. *Tree Physiol.* 10, 21–31.
- Wünsche, J.N., Palmer, J.W., 1997. Portable through-flow cuvette system for measuring whole-canopy gas exchange of apple trees in the field. *Hortscience* 32, 653–658.
- Würth, M.K.R., Peláez-Riedl, S., Wright, S.J., Körner, C., 2005. Non-structural carbohydrate pools in a tropical forest. *Oecologia* 143, 11–24.
- Xu, X., Yang, F., Xiao, X., Zhang, S., Korpelainen, H., Li, C., 2008. Sex-specific responses of *Populus cathayana* to drought and elevated temperatures. *Plant Cell Environ.* 31, 850–860.
- Yoder, B.J., Ryan, M.G., Waring, R.H., Schoettle, A.W., Kaufmann, M.R., 1994. Evidence of reduced photosynthetic rates in old trees. *Forest Sci.* 40, 513–527.
- Zavala, M.A., Espelta, J.M., Retana, J., 2000. Constraints and trade-offs in Mediterranean plant communities: the case of holm oak-Aleppo pine forests. *Bot. Rev.* 66, 119–149.