

# Rethinking False Spring Risk

Authors:

C. J. Chamberlain <sup>1,2</sup>, B. I. Cook <sup>3</sup>, I. Garcia de Cortazar Atauri <sup>4</sup> & E. M. Wolkovich <sup>1,2,5</sup>

*Author affiliations:*

<sup>1</sup>Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

<sup>2</sup>Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, USA;

<sup>3</sup>NASA Goddard Institute for Space Studies, New York, New York, USA;

<sup>4</sup>French National Institute for Agricultural Research, INRA, US1116 AgroClim, F-84914 Avignon, France

<sup>5</sup>Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia, 2424 Main Mall, Vancouver, BC V6T 1Z4

\*Corresponding author: 248.953.0189; cchamberlain@g.harvard.edu

*Keywords:* false spring, phenology, freezing tolerance, climate change, forest communities

*Paper type:* Opinion

## Abstract

Temperate plants are at risk of being exposed to late spring freezes — often called false springs — which can be damaging ecologically and economically. As climate change may alter the prevalence and severity of false springs, our ability to forecast such events has become more critical. Currently, many false spring studies simplify the ecological and physiological information needed for accurate predictions of the level of plant damage from late spring freezes. Here we review the complexity of factors driving a plant’s false spring risk. We highlight how life stage and functional group, species differences in morphology and phenology, and regional climatic differences contribute to the damage potential of false springs. Integrating these complexities could rapidly advance forecasting of false spring events in climate change and ecological studies.

## Introduction

Plants from temperate environments time their growth each spring to follow rising temperatures alongside increasing light and soil resource availability. While tracking spring resource availability, individuals that budburst before the last freeze date are at risk of leaf loss, damaged wood tissue, and slowed canopy development (Gu *et al.*, 2008; Hufkens *et al.*, 2012). These damaging late spring freezes are also known as false springs, and are widely documented to result in adverse ecological and economic consequences (Ault *et al.*, 2013; Knudson, 2012).

Climate change is expected to cause an increase in damage from false spring events due to earlier spring onset and potentially greater fluctuations in temperature in some regions (Inouye, 2008; Martin *et al.*, 2010). Already, multiple studies have documented false springs in recent years (Augsburger, 2009, 2013; Gu *et al.*, 2008; Menzel *et al.*, 2015) and some have linked these events to climate change (Allstadt *et al.*, 2015; Ault *et al.*, 2013; Muffler *et al.*, 2016; Vitra *et al.*, 2017; Xin, 2016). This interest in false springs has led to a growing body of research investigating the effects on temperate forests. For this research to produce accurate predictions, however, researchers need methods that properly evaluate the effects of false springs across diverse species and climate regimes.

Current metrics for estimating false springs events are generally simple, often requiring an estimate for the start of biological ‘spring’ (i.e., budburst) and whether temperatures below a particular threshold occurred in the following week. Such estimates inherently assume consistency of damage across species, functional group,

life stages, and other climatic regimes, ignoring that such factors can greatly impact plants' false spring risk. As a result, such indices may lead to inaccurate estimates and predictions.

In this paper we highlight the complexity of factors driving a plant's false spring risk and provide a road map for improved metrics. We show how location within a forest or canopy (Augspurger, 2013), interspecific variation in avoidance and tolerance strategies (Martin *et al.*, 2010; Muffler *et al.*, 2016), freeze temperature thresholds (Lenz *et al.*, 2013), and regional effects (Muffler *et al.*, 2016) unhinge simple metrics of false spring. We argue that a new approach that integrates these and other crucial factors would help accurately determine current false spring damage and improve predictions of spring freeze risk under a changing climate — while potentially providing novel insights to how plants respond to and are shaped by spring frost.

## Defining False Spring: An example in one temperate plant community

Temperate forest plants experience elevated risk of frost damage during the spring due to the stochastic timing of frosts. Freezing temperatures following a warm spell can result in plant damage or even death (Ludlum, 1968; Mock *et al.*, 2007). Many temperate species exhibit flexible spring phenologies, which help them minimize spring freezing risk, but freeze damage can still occur. Flowering and fruiting are generally more sensitive to freezing temperatures than vegetative phases (Inouye, 2000; Augspurger, 2009; CaraDonna, Paul J and Bain, Justin A, 2016; Lenz *et al.*, 2013), but false spring events that occur during the vegetative growth phenophases may impose the greatest freezing threat to deciduous plant species. It can take 16-38 days for trees to refoliate after a spring freeze (Gu *et al.*, 2008; Augspurger, 2009, 2013; Menzel *et al.*, 2015), which can detrimentally affect crucial processes such as carbon uptake and nutrient cycling (Hufkens *et al.*, 2012; Richardson *et al.*, 2013; Klosterman *et al.*, 2018). For this reason, we will focus primarily on spring freeze risk for the vegetative phases.

Once buds exit the dormancy phase, they are less freeze tolerant and less resistant to ice formation in contrast to mature tissues (Lenz *et al.*, 2013; Taschler *et al.*, 2004; Vitasse *et al.*, 2014b). Intracellular ice formation from false spring events often results in severe leaf and stem damage (Burke *et al.*, 1976; Sakai & Larcher, 1987). Ice formation can also occur indirectly (i.e., extracellularly), which results in freezing dehydration and mimics drought conditions (Beck *et al.*, 2004; Hofmann & Bruehlheide, 2015; Pearce, 2001). Both forms of ice formation can cause defoliation and crown dieback (Gu *et al.*, 2008). An effective and consistent definition of

false spring would accurately determine the amount and type of ice formation to evaluate the level of damage that could occur.

Currently there are several ways to define a false spring. A common definition describes a false spring as having two phases: rapid vegetative growth prior to a freeze and a post-freeze setback (Gu *et al.*, 2008). Other definitions instill more precise temporal parameters, specific to certain regions (e.g., in Augspurger, 2013, false spring for the Midwestern United States is defined as a warmer than average March, a freezing April, and enough growing degree days between budburst and the last freeze date). A widely used definition integrates a mathematical equation to quantify a false spring event. This equation, known as a False Spring Index (FSI), signifies the likelihood of damage to occur from a late spring freeze. Currently, FSI is evaluated annually by the day of budburst and the day of last spring freeze (often calculated at  $-2.2^{\circ}\text{C}$ , Schwartz, 1993) through the simple equation (Marino *et al.*, 2011):

$$FSI = \text{Day of Year}(\textit{LastSpringFreeze}) - \text{Day of Year}(\textit{Budburst}) \quad (1)$$

Negative values indicate no-risk situations, whereas a damaging FSI is currently defined to be seven or more days between budburst and the last freeze date (Equation 1) (Peterson & Abatzoglou, 2014). This seven-day threshold captures the reality that leaf tissue is at high risk of damage from frost in the period after budburst, with later vegetative phases (e.g., after full leafout) being more resistant to such damage.

To demonstrate how the FSI definition works, we applied it to data from the Harvard Forest Long-term Ecological Research program in Massachusetts. We used three separate methodologies to calculate spring onset: long-term ground observational data (O’Keefe, 2014), PhenoCam data from Harvard Forest (Richardson, 2015), and USA National Phenology Network’s (USA-NPN) Extended Spring Index (SI-x) data (USA-NPN, 2016). These spring onset values were then inputted into the FSI equation (Equation 1) to calculate FSI from 2008 to 2014 (Figure 1).

Each methodology rendered different FSI values, suggesting different false spring damage for the same site and same year. For most years, the observational FSI and PhenoCam FSI are about 10-15 days lower than the SI-x data. This is especially important for 2008, when the SI-x data indicates a false spring year, whereas the other two datasets do not. In 2012, the observational data and PhenoCam data diverge slightly and the PhenoCam FSI is over 30 days less than the SI-x value.

The reason for these discrepancies is that each method evaluates spring onset by integrating different at-

tributes such as age, species or functional group. Spring phenology in temperate forests typically progresses by functional group: understory species and young trees tend to initiate budburst first, whereas larger canopy species start later in the season (Richardson & O’Keefe, 2009; Xin, 2016). The different FSI values determined in Figure 1 exemplify the differences in functional group spring onset dates and illustrate variations in forest demography and phenology. While the SI-x data (based on observations of early-active shrub species, including lilac, *Syringa vulgaris*) may best capture understory dynamics, the PhenoCam and observational FSI data integrate over larger canopy species and are more site-specific. Such differences are visible each year, as the canopy-related metrics show lower risk, but are especially apparent in 2012. In 2012, a false spring event was reported through many regions of the US due to warm temperatures occurring in March (Ault *et al.*, 2015). These high temperatures would most likely have been too early for larger canopy species to initiate budburst but they would have affected smaller understory species, as is seen by the high risk of the SI-x FSI in Figure 1.




Figure 1: False Spring Index (FSI) values from 2008 to 2014 vary across methodologies. To calculate spring onset, we used the USA-NPN Extended Spring Index tool for the USA-NPN FSI values, which are in green (USA-NPN, 2016), long-term ground observational data for the observed FSI values, which are in red (O’Keefe, 2014), and near-surface remote-sensing canopy data for the PhenoCam FSI values, which are in blue (Richardson, 2015). The solid line at FSI=0 indicates a boundary between a likely false spring event or not, with positive numbers indicating a false spring likely occurred and negative numbers indicating a false spring most likely did not occur. The dotted line at FSI=7 indicates the seven-day threshold frequently used in false spring definitions, which suggests years with FSI values greater than seven very likely had false spring events.

Yet, in contrast to our three metrics of spring onset for one site, most FSI work currently ignores variation across functional groups — instead using one metric of spring onset and assuming it applies to the whole community of plants (Allstadt *et al.*, 2015; Marino *et al.*, 2011; Mehdipoor & Zurita-Milla, 2017; Peterson & Abatzoglou, 2014). The risk of a false spring varies across habitats and with species composition since spring onset is not consistent across functional groups (Martin *et al.*, 2010). Therefore, one spring onset date cannot be used as an effective proxy for all species. False spring studies should first assess the forest demographics and functional groups relevant to the study question in order to estimate the date of spring onset. However, as we outline below, considering different functional groups is unlikely to be enough for robust predictions. It will also be important to integrate species differences within functional groups and to consider the various interspecific avoidance and tolerance strategies that species have evolved against false springs.

## Plant Physiology and Diversity versus the Current False Spring Definition

Plants have evolved to minimize false spring damage through two strategies: avoidance and tolerance. Many temperate forest plants utilize various morphological strategies to be more frost tolerant: some have increased ‘packability’ of leaf primordia in winter buds, which may permit more rapid leafout (Edwards *et al.*, 2017) and minimize the exposure time of less resistant tissues. Other species have young leaves with more trichomes, which may act as a buffer against hard or radiative frosts (Agrawal *et al.*, 2004; Prozherina *et al.*, 2003). And many other individuals are able to respond to abiotic cues such as consistently dry winters. Species living in habitats with drier winters develop shoots and buds with decreased water content, which makes the buds more tolerant to drought and also to false spring events (Beck *et al.*, 2007; Kathke & Bruelheide, 2011; Hofmann & Bruelheide, 2015; Morin *et al.*, 2007; Muffler *et al.*, 2016; Norgaard Nielsen & Rasmussen, 2009; Poirier *et al.*, 2010). These morphological strategies are probably only a few of the many ways plants work to avoid certain types of frost damage, and more studies are needed to investigate the interplay between morphological traits and false spring tolerance.

Rather than being more tolerant of spring freezing temperatures, some temperate forest species have evolved to avoid frosts via their phenologies. Effective avoidance strategies require well-timed spring phenologies. Most temperate deciduous tree species optimize growth and minimize spring freeze damage by using three cues to initiate budburst: low winter temperatures (chilling), warm spring temperatures (forcing), and in-

creasing photoperiods (Chuine, 2010). The evolution of these three cues and their interactions have permitted temperate plant species to occupy more northern ecological niches (Kollas *et al.*, 2014) and decrease the risk of false spring damage (Charrier *et al.*, 2011). One avoidance strategy, for example, is the interaction between over-winter chilling and spring forcing temperatures. Warm temperatures earlier in the winter will not result in early budburst due to insufficient chilling (Basler & Körner, 2012). Likewise, photoperiod sensitivity is a common false spring avoidance strategy: species that respond strongly to photoperiod cues in addition to warm spring temperatures are unlikely to have large advances in budburst with warming, and thus may evade false spring events as warming continues (Basler & Korner, 2014).

## Defining Vegetative Risk

Phenology and frost tolerance are clearly intertwined — with important variation occurring across different phenological phases. However, plants will suffer greater long-term effects from the loss of photosynthetic tissue, which could impact multiple years of growth, reproduction, and canopy development (Vitasse *et al.*, 2014a; Xie *et al.*, 2015). However, there is high variability in defining a damaging temperature threshold across species, including between agricultural and ecological studies (Figure 2).




Figure 2: A comparison of damaging spring freezing temperature thresholds across ecological and agronomic studies. Each study is listed on the vertical axis along with the taxonomic group of focus. Next to the species name is the freezing definition used within that study (e.g., 100% is 100% whole plant lethality). Each point is the best estimate recorded for the temperature threshold with standard deviation if indicated in the study.

There is also important variation within certain phenological phases. Most notably, within the vegetative phases of spring leafout, plants that have initiated budburst but have not fully leafed out are more likely to sustain damage from a false spring than individuals past the leafout phase. This is because freezing tolerance is lowest after budburst begins until the leaf is fully unfolded (Lenz *et al.*, 2016). Therefore, the rate of budburst and the length of time between budburst and leafout is essential for predicting the level of damage from a false spring event. We will refer to the timing between these phenophases — budburst to leafout — as the duration of vegetative risk (Figure 3).






Figure 3: Differences in spring phenology and false spring risk across two species: *Ilex mucronata* (L.) and *Betula alleghaniensis* (Marsh.). We mapped a hypothetical false spring event based on historical weather data and long-term observational phenological data collected at Harvard Forest (O’Keefe, 2014). In this scenario, *Ilex mucronata*, which budbursts early and generally has a short period between budburst (light green squares) and leafout (dark green triangles), would be exposed to a false spring event during its duration of vegetative risk (i.e., from budburst to leafout), whereas *Betula alleghaniensis* would avoid it entirely (even though it has a longer duration of vegetative risk), due to later budburst.

## How Species Phenological Cues Shape Vegetative Risk

Predictions of false spring critically depend on understanding what controls the duration of vegetative risk across species. For temperate species, the three major cues (winter chilling temperatures, spring warming temperatures and photoperiod) that control budburst (Chuine, 2010) play a dominant role. Most phenological studies currently focus on one phenophase (i.e., budburst or leafout) but, to examine false spring risk, it is important to examine the effects of the three phenological cues and their interactions on the duration of vegetative risk—that is, researchers must collect data on both budburst and leafout timing.

Such cues may provide a starting point for predicting how climate change will alter the duration of vegetative risk. Robust predictions will require more information, especially the emissions scenario realized over coming

167 decades (IPCC, 2015), but some outcomes with warming are more expected than others. For example, higher  
168 temperatures are generally expected to increase forcing and decrease chilling in many locations, as well as  
169 to trigger budburst at times of the year when daylength is shorter. Using data from a recent study that  
170 manipulated all three cues and measured budburst and leafout (Flynn & Wolkovich, 2018) shows that any  
171 one of these effects alone can have a large impact on the duration of vegetative risk (Figure 4): more forcing  
172 shortens it substantially (-15 to -8 days), while shorter photoperiods and less chilling increase it to a lesser  
173 extent (+3 to 9 days). Together, however, the expected shifts generally shorten the duration of vegetative  
174 risk by 4-13 days, both due to the large effect of forcing and the combined effects of multiple cues. How  
175 shortened the risk period is, however, varies strongly by species and highlights how climate change may speed  
176 some species through this high risk period, but not others. Additionally, as our results are for a small set  
177 of species we expect other species may have more diverse responses, as has already been seen in shifts in  
178 phenology with warming (Cleland *et al.*, 2006; Fu *et al.*, 2015; Xin, 2016).




Figure 4: We examine the effects of phenological cues on the duration of vegetative risk across three species: *Acer pensylvanicum*, *Fagus grandifolia*, and *Populus grandidentata*. ‘More Forcing’ is a 5°C increase in spring warming temperatures, ‘Shorter Photoperiod’ is a 4 hour decrease in photoperiod and ‘Less Chilling’ is a 30 day decrease in over-winter chilling. Along with the estimated isolated effects, we the show the combined predicted shifts in phenological cues with potential climate change (i.e., more forcing with shorter photoperiod and more forcing with less chilling) and the subsequent shifts in duration of vegetative risk across species. To calculate the combined effects, we added the estimated isolated effects of each cue alone with the interaction effects for the relevant cues for each species.

These findings highlight the need for further studies on the interplay among chilling, forcing, and photoperiod cues and the duration of vegetative risk across species. This is especially true for species occupying ecological niches more susceptible to false spring events; even if warming causes a shortened duration of vegetative risk for such species, the related earlier budburst dates could still lead to greater risk of false spring exposure.

## Predictable Regional Differences in Climate, Species Responses and False Spring Risk

Robust predictions must consider the interplay of species cues with a specific location's climate. Climate and thus false spring risk vary across regions. Some regions experience harsher winters and greater temperature variability throughout the year (Figure 5 e.g., Maine, USA), and these more variable regions often have a much higher risk of false spring than others (Figure 5 e.g., Lyon, France). Understanding and integrating spatiotemporal effects and regional differences when investigating false spring risk and duration of vegetative risk would help improve predictions as climate change progresses. Such differences depend both on the local climate, the local species and the cues for each species at that location, as a single species may have varying cues across space. Therefore, based on cues alone, different regions may have different durations of vegetative risk for the same species (Caffarra & Donnelly, 2011; Partanen, 2004; Vihera-aarnio *et al.*, 2006). Studies also show that different species within the same location can exhibit different sensitivities to the three cues (Basler & Körner, 2012; Laube *et al.*, 2013), further amplifying the myriad of climatic and phenological shifts that determine false spring risk in a region.




Figure 5: False spring risk can vary dramatically across regions. Here we show the period when plants are most at risk to tissue loss – between budburst and leafout (upper, lines represent the range with the thicker line representing the interquartile range) and the variation in the number of freeze days ( $-2.2^{\circ}\text{C}$ ) (Schwartz, 1993) that occurred on average over the past 50 years for five different sites (lower, bars represent the range, points represent the mean). Data come from USA-NPN SI-x tool (1981-2016) and observational studies (1950-2016) for phenology (Schaber & Badeck, 2005; Soudani et al., 2012; USA-NPN, 2016; White et al., 2009) and NOAA Climate Data Online tool for climate (from 1950-2016).

How a single species' phenological cues varies across space is not yet well predicted. Some studies have investigated how phenological cues for budburst vary across space, including variation across populations, by using latitudinal gradients (Gauzere *et al.*, 2017; Sogaard, Gunnhild and Johnsen, Øystein and Nilsen, Jarle and Junttila, Olavi, 2008; Way & Montgomery, 2015; Zohner *et al.*, 2016). Fewer, however, have integrated distance from the coast (but see Aitken & Bemmels, 2015; Harrington & Gould, 2015; Myking & Skroppa, 2007) or regional effects. Some studies assert that the distance from the coast is a stronger indicator of budburst timing than latitude (Myking & Skroppa, 2007), with populations further inland initiating budburst

first, whereas those closer to the coast budburst later in the season. Therefore, to understand the interplay between duration of vegetative risk and climatic variation it is important to recognize how climate regime extremes (e.g., seasonal trends, annual minima and annual maxima) vary across regions and how they will shift in the future: as climatic regimes are altered by climate change false spring risk could vary in intensity across regions and time (i.e., regions currently at high risk of false spring damage could become low-risk regions in the future and vice versa).

## Conclusion

Temperate forest trees are most at risk to frost damage in the spring due to the stochasticity of spring freezes. With warm temperatures advancing in the spring but last spring freeze dates advancing at a slower rate, there could be more damaging false spring events in the future, especially in high-risk regions (Gu *et al.*, 2008; Inouye, 2008; Liu *et al.*, 2018). Current equations for evaluating false spring damage (e.g., Equation 1) largely simplify the myriad complexities involved in assessing false spring damage and risks. More studies aimed at understanding relationships among species avoidance and tolerance strategies, climatic regimes, and physiological cue interactions with the duration of vegetative risk would improve predictions. Additionally, research to establish temperature thresholds for damage across functional types and phenophases will help effectively predict false spring risk in the future. An integrated approach to assessing past and future spring freeze damage would provide novel insights into plant strategies, and offer more robust predictions as climate change progresses, which is essential for mitigating the adverse ecological and economic effects of false springs.

## Acknowledgments

We thank D. Buonaiuto, W. Daly, A. Ettinger, and I. Morales-Castilla for comments and insights that improved the manuscript.

## References

Agrawal AA, Conner JK, Stinchcombe JR (2004) Evolution of plant resistance and tolerance to frost damage. *Ecology Letters*, **7**, 1199–1208. doi:10.1111/j.1461-0248.2004.00680.x. URL <http://dx.doi.org/10.1111/>

j.1461-0248.2004.00680.x.

Aitken SN, Bemmels JB (2015) Time to get moving: assisted gene flow of forest trees. *Evolutionary Applications*, **9**, 271–290. doi:10.1111/eva.12293. URL <http://dx.doi.org/10.1111/eva.12293>.

Allstadt AJ, Vavrus SJ, Heglund PJ, Pidgeon AM, Wayne E, Radeloff VC (2015) Spring plant phenology and false springs in the conterminous U. S. during the 21st century. *Environmental Research Letters (submitted)*, **10**, 104008. doi:10.1088/1748-9326/10/10/104008. URL <http://dx.doi.org/10.1088/1748-9326/10/10/104008>.

Augspurger CK (2009) Spring 2007 warmth and frost: Phenology, damage and refoliation in a temperate deciduous forest. *Functional Ecology*, **23**, 1031–1039. doi:10.1111/j.1365-2435.2009.01587.x.

Augspurger CK (2013) Reconstructing patterns of temperature, phenology, and frost damage over 124 years: Spring damage risk is increasing. *Ecology*, **94**, 41–50. doi:10.1890/12-0200.1.

Ault TR, Henebry GM, de Beurs KM, Schwartz MD, Betancourt JL, Moore D (2013) The false spring of 2012, earliest in North American record. *Eos, Transactions American Geophysical Union*, **94**, 181–182. doi:10.1002/2013EO200001.

Ault TR, Schwartz MD, Zurita-Milla R, Weltzin JF, Betancourt JL (2015) Trends and natural variability of spring onset in the coterminous United States as evaluated by a new gridded dataset of spring indices. *Journal of Climate*, **28**, 8363–8378. doi:10.1175/JCLI-D-14-00736.1.

Barker D, Loveys B, Egerton J, Gorton H, Williams W, Ball M (2005) CO<sub>2</sub> Enrichment predisposes foliage of a eucalypt to freezing injury and reduces spring growth. *Plant, Cell and Environment*, **28**, 1506–1515.

Barlow K, Christy B, O’Leary G, Riffkin P, Nuttall J (2015) Simulating the impact of extreme heat and frost events on wheat crop production: A review. *Field Crops Research*, **171**, 109–119.

Basler D, Körner C (2012) Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agricultural and Forest Meteorology*, **165**, 73–81. doi:10.1016/j.agrformet.2012.06.001. URL <http://dx.doi.org/10.1016/j.agrformet.2012.06.001>.

Basler D, Korner C (2014) Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiology*, **34**, 377–388. doi:10.1093/treephys/tpu021. URL <http://dx.doi.org/10.1093/treephys/tpu021>.

255 Beck EH, Fettig S, Knake C, Hartig K, Bhattarai T (2007) Specific and unspecific responses of plants to cold  
256 and drought stress. *Journal of Biosciences*, **32**, 501–510.

257 Beck EH, Heim R, Hansen J (2004) Plant resistance to cold stress: Mechanisms and environmental signals  
258 triggering frost hardening and dehardening. *Journal of Biosciences*, **29**, 449–459.

259 Burke M, Gusta L, Quamme H, Weiser C, Li P (1976) Freezing and injury in plants. *Annual Review of Plant*  
260 *Physiology*, **27**, 507–528.

261 Caffarra A, Donnelly A (2011) The ecological significance of phenology in four different tree species: Effects  
262 of light and temperature on bud burst. *International Journal of Biometeorology*, **55**, 711–721. doi:10.1007/  
263 s00484-010-0386-1.

264 CaraDonna, Paul J and Bain, Justin A (2016) Frost sensitivity of leaves and flowers of subalpine plants is  
265 related to tissue type and phenology. *Journal of Ecology*, **104**, 55–64. doi:10.1111/1365-2745.12482.

266 Charrier G, Bonhomme M, Lacointe A, Améglio T (2011) Are budburst dates, dormancy and cold acclimation  
267 in walnut trees (*Juglans regia* L.) under mainly genotypic or environmental control? *International Jour-*  
268 *nal of Biometeorology*, **55**, 763–774. doi:10.1007/s00484-011-0470-1. URL [https://doi.org/10.1007/](https://doi.org/10.1007/s00484-011-0470-1)  
269 [s00484-011-0470-1](https://doi.org/10.1007/s00484-011-0470-1).

270 Chuine I (2010) Why does phenology drive species distribution? *Philosophical Transactions of the Royal*  
271 *Society B: Biological Sciences*, **365**, 3149–3160. doi:10.1098/rstb.2010.0142. URL [http://dx.doi.org/](http://dx.doi.org/10.1098/rstb.2010.0142)  
272 [10.1098/rstb.2010.0142](http://dx.doi.org/10.1098/rstb.2010.0142).

273 Cleland E, Chiariello N, Loarie S, Mooney H, Field C (2006) Diverse responses of phenology to global changes  
274 in a grassland ecosystem. *PNAS*, **103**, 13740–13744.

275 Edwards EJ, Chatelet DS, Spriggs EL, Johnson ES, Schlutius C, Donoghue MJ (2017) Correlation, causation,  
276 and the evolution of leaf teeth: A reply to Givnish and Kriebel. *Am J Bot*, **104**, 509–515. doi:10.3732/  
277 ajb.1700075.

278 Flynn DFB, Wolkovich EM (2018) Temperature and photoperiod drive spring phenology across all species in  
279 a temperate forest community. *New Phytologist*, **219**. doi:10.1111/nph.15232. URL [http://dx.doi.org/](http://dx.doi.org/10.1111/nph.15232)  
280 [10.1111/nph.15232](http://dx.doi.org/10.1111/nph.15232).

281 Fu YH, Zhao H, Piao S, *et al.* (2015) Declining global warming effects on the phenology of spring leaf unfolding.  
282 *Nature*, **526**, 104–107. doi:10.1038/nature15402. URL <http://dx.doi.org/10.1038/nature15402>.



- Gauzere J, Delzon S, Davi H, Bonhomme M, Garcia de Cortazar-Atauri I, Chuine I (2017) Integrating interactive effects of chilling and photoperiod in phenological process-based models. A case study with two European tree species: *Fagus sylvatica* and *Quercus petraea*. *Agricultural and Forest Meteorology*, **244-255**, 9–20.
- Gu L, Hanson PJ, Post WM, *et al.* (2008) The 2007 Eastern US spring freeze: Increased cold damage in a warming world. *BioScience*, **58**, 253. doi:10.1641/B580311.
- Harrington CA, Gould PJ (2015) Tradeoffs between chilling and forcing in satisfying dormancy requirements for Pacific Northwest tree species. *Frontiers in Plant Science*, **6**. doi:10.3389/fpls.2015.00120. URL <http://dx.doi.org/10.3389/fpls.2015.00120>.
- Hofmann M, Bruehlheide H (2015) Frost hardiness of tree species is independent of phenology and macroclimatic niche. *Journal of Biosciences*, **40**, 147–157. doi:10.1007/s12038-015-9505-9.
- Hufkens K, Friedl MA, Keenan TF, Sonnentag O, Bailey A, O’Keefe J, Richardson AD (2012) Ecological impacts of a widespread frost event following early spring leaf-out. *Global Change Biology*, **18**, 2365–2377. doi:10.1111/j.1365-2486.2012.02712.x.
- Inouye DW (2000) The ecological and evolutionary significance of frost in the context of climate change. *Ecology Letters*, **3**, 457–463. doi:10.1046/j.1461-0248.2000.00165.x.
- Inouye DW (2008) Effects of climate change on phenology, frost damage, and floral abundance of montane wildflowers. *Ecology*, **89**, 353–362.
- IPCC (2015) *Climate change 2014: mitigation of climate change*, vol. 3. Cambridge University Press.
- Kathke S, Bruehlheide H (2011) Differences in frost hardiness of two Norway spruce morphotypes growing at Mt. Brocken, Germany. *Flora - Morphology, Distribution, Functional Ecology of Plants*, **206**, 120–126. doi:10.1016/j.flora.2010.09.007. URL <http://dx.doi.org/10.1016/j.flora.2010.09.007>.
- Klosterman S, Hufkens K, Richardson AD (2018) Later springs green-up faster: the relation between onset and completion of green-up in deciduous forests of North America. *International Journal of Biometeorology*. doi:10.1007/s00484-018-1564-9. URL <https://doi.org/10.1007/s00484-018-1564-9>.
- Knudson W (2012) The economic impact of the spring’s weather on the fruit and vegetable sectors. URL <http://legislature.mi.gov/documents/2011-2012/CommitteeDocuments/House/Agriculture/Testimony/Committee1-5-30-2012.pdf>.

- 311 Kollas C, Körner C, Randin CF (2014) Spring frost and growing season length co-control the cold range  
312 limits of broad-leaved trees. *Journal of Biogeography*, **41**, 773–783. doi:10.1111/jbi.12238.
- 313 Laube J, Sparks TH, Estrella N, Höfler J, Ankerst DP, Menzel A (2013) Chilling outweighs photoperiod in  
314 preventing precocious spring development. *Global Change Biology*, **20**, 170–182. doi:10.1111/gcb.12360.  
315 URL <http://dx.doi.org/10.1111/gcb.12360>.
- 316 Lenz A, Hoch G, Körner C, Vitasse Y (2016) Convergence of leaf-out towards minimum risk of freezing  
317 damage in temperate trees. *Functional Ecology*, **30**, 1–11. doi:10.1111/1365-2435.12623.
- 318 Lenz A, Hoch G, Vitasse Y, Körner C (2013) European deciduous trees exhibit similar safety margins against  
319 damage by spring freeze events along elevational gradients. *New Phytologist*, **200**, 1166–1175. doi:10.1111/  
320 np.12452.
- 321 Liu Q, Piao S, Janssens IA, *et al.* (2018) Extension of the growing season increases vegetation exposure to  
322 frost. *Nature Communications*, **9**. doi:10.1038/s41467-017-02690-y. URL <http://dx.doi.org/10.1038/s41467-017-02690-y>.
- 324 Longstroth M (2012) Protect blueberries from spring freezes by using sprinklers. url. URL [https://www.](https://www.canr.msu.edu/news/protect_blueberries_from_spring_freezes_by_using_sprinklers)  
325 [canr.msu.edu/news/protect\\_blueberries\\_from\\_spring\\_freezes\\_by\\_using\\_sprinklers](https://www.canr.msu.edu/news/protect_blueberries_from_spring_freezes_by_using_sprinklers).
- 326 Longstroth M (2013) Assessing frost and freeze damage to flowers and buds of fruit trees.  
327 url. URL [https://www.canr.msu.edu/news/assessing\\_frost\\_and\\_freeze\\_damage\\_to\\_flowers\\_and\\_](https://www.canr.msu.edu/news/assessing_frost_and_freeze_damage_to_flowers_and_buds_of_fruit_trees)  
328 [buds\\_of\\_fruit\\_trees](https://www.canr.msu.edu/news/assessing_frost_and_freeze_damage_to_flowers_and_buds_of_fruit_trees).
- 329 Ludlum DM (1968) *Early American Winters: 1604-1820*. 3. Boston: American Meteorological Society.
- 330 Marino GP, Kaiser DP, Gu L, Ricciuto DM (2011) Reconstruction of false spring occurrences over the south-  
331 eastern United States, 1901–2007: an increasing risk of spring freeze damage? *Environmental Research*  
332 *Letters*, **6**, 24015. doi:10.1088/1748-9326/6/2/024015.
- 333 Martin M, Gavazov K, Körner C, Hattenschwiler S, Rixen C (2010) Reduced early growing season freezing  
334 resistance in alpine treeline plants under elevated atmospheric  $CO_2$ . *Global Change Biology*, **16**, 1057–1070.  
335 doi:10.1111/j.1365-2486.2009.01987.x. URL <http://dx.doi.org/10.1111/j.1365-2486.2009.01987.x>.
- 336 Mehdipoor H, Zurita-Milla EIVR (2017) Continental-scale monitoring and mapping of false spring: A cloud  
337 computing solution. URL <http://www.geocomputation.org/2017/papers/48.pdf>.

- Menzel A, Helm R, Zang C (2015) Patterns of late spring frost leaf damage and recovery in a European beech (*Fagus sylvatica* L.) stand in south-eastern Germany based on repeated digital photographs. *Frontiers in Plant Science*, **6**, 110. doi:10.3389/fpls.2015.00110. URL <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4338663/>.
- Mock CJ, Mojzisek J, McWaters M, Chenoweth M, Stahle DW (2007) The winter of 1827–1828 over eastern North America: a season of extraordinary climatic anomalies, societal impacts, and false spring. *Climatic Change*, **83**, 87–115. doi:10.1007/s10584-006-9126-2. URL <http://dx.doi.org/10.1007/s10584-006-9126-2>.
- Morin X, Ameglio T, Ahas R, *et al.* (2007) Variation in cold hardiness and carbohydrate concentration from dormancy induction to bud burst among provenances of three European oak species. *Tree Physiology*, **27**, 817–825.
- Muffler L, Beierkuhnlein C, Aas G, Jentsch A, Schweiger AH, Zohner C, Kreyling J (2016) Distribution ranges and spring phenology explain late frost sensitivity in 170 woody plants from the Northern Hemisphere. *Global Ecology and Biogeography*, **25**, 1061–1071. doi:10.1111/geb.12466. URL <http://dx.doi.org/10.1111/geb.12466>.
- Myking T, Skroppa T (2007) Variation in phenology and height increment of northern *Ulmus glabra* populations: Implications for conservation. *Scandinavian Journal of Forest Research*, **22**, 369–374.
- Norgaard Nielsen CC, Rasmussen HN (2009) Frost hardening and dehardening in *Abies procera* and other conifers under differing temperature regimes and warm-spell treatments. *Forestry*, **82**, 43–59. doi:10.1093/forestry/cpn048. URL <http://dx.doi.org/10.1093/forestry/cpn048>.
- O’Keefe J (2014) Phenology of woody species at Harvard Forest since 1990. URL <http://harvardforest.fas.harvard.edu:8080/exist/apps/datasets/showData.html?id=hf003>.
- Partanen J (2004) Dependence of photoperiodic response of growth cessation on the stage of development in *Picea abies* and *Betula pendula* seedlings. *Forest Ecology and Management*, **188**, 137–148. doi:10.1016/j.foreco.2003.07.017. URL <http://dx.doi.org/10.1016/j.foreco.2003.07.017>.
- Pearce R (2001) Plant freezing and damage. *Annals of Botany*, **87**, 417–424.
- Peterson AG, Abatzoglou JT (2014) Observed changes in false springs over the contiguous United States. *Geophysical Research Letters*, **41**, 2156–2162. doi:10.1002/2014GL059266.

- Poirier M, Lacointe A, Ameglio T (2010) A semi-physiological model of cold hardening and dehardening in walnut stem. *Tree Physiology*, **30**, 1555–1569. doi:10.1093/treephys/tpq087. URL <http://dx.doi.org/10.1093/treephys/tpq087>.
- Prozherina N, Freiwald V, Rousi M, Oksanen E (2003) Interactive effect of springtime frost and elevated ozone on early growth, foliar injuries and leaf structure of birch (*Betula pendula*). *New Phytologist*, **159**, 623–636. doi:10.1046/j.1469-8137.2003.00828.x. URL <http://dx.doi.org/10.1046/j.1469-8137.2003.00828.x>.
- Richardson A, O’Keefe J (2009) *Phenological differences between understory and overstory: a case study using the long-term Harvard Forest records*, pp. 87–117. A. Noormets (Ed.), *Phenology of Ecosystem Processes*, Springer, New York.
- Richardson AD (2015) PhenoCam images and canopy phenology at Harvard Forest since 2008. URL <http://harvardforest.fas.harvard.edu:8080/exist/apps/datasets/showData.html?id=hf158>.
- Richardson AD, Keenan TF, Migliavacca M, Ryu Y, Sonnentag O, Toomey M (2013) Climate change, phenology, and phenological control of vegetation feedbacks to the climate system. *Agricultural and Forest Meteorology*, **169**, 156 – 173. doi:<https://doi.org/10.1016/j.agrformet.2012.09.012>. URL <http://www.sciencedirect.com/science/article/pii/S0168192312002869>.
- Sakai A, Larcher W (1987) *Frost Survival of Plants*. Springer-Verlag.
- Sánchez B, Rasmussen A, Porter JR (2013) Temperatures and the growth and development of maize and rice: a review. *Global Change Biology*, **20**, 408–417. doi:10.1111/gcb.12389. URL <http://dx.doi.org/10.1111/gcb.12389>.
- Schaber J, Badeck FW (2005) Plant phenology in Germany over the 20th century. *Regional Environmental Change*, **5**, 37–46. doi:10.1007/s10113-004-0094-7. URL <http://dx.doi.org/10.1007/s10113-004-0094-7>.
- Schwartz MD (1993) Assessing the onset of spring: A climatological perspective. *Physical Geography*, **14**(6), 536–550.
- Søgaard, Gunnhild and Johnsen, Øystein and Nilsen, Jarle and Junttila, Olavi (2008) Climatic control of bud burst in young seedlings of nine provenances of Norway spruce. *Tree Physiology*, **28**, 311–320.
- Soudani K, Hmimina G, Delpierre N, *et al.* (2012) Ground-based Network of NDVI measurements for tracking temporal dynamics of canopy structure and vegetation phenology in different biomes. *Remote Sensing of*

*Environment*, **123**, 234–245. doi:10.1016/j.rse.2012.03.012. URL <http://dx.doi.org/10.1016/j.rse.2012.03.012>.

Taschler D, Beikircher B, Neuner G (2004) Frost resistance and ice nucleation in leaves of five woody timberline species measured in situ during shoot expansion. *Tree Physiology*, **24**, 331–337.

USA-NPN (2016) USA National Phenology Network Extended Spring Indices. URL <http://dx.doi.org/10.5066/F7XD0ZRK>.

Vihera-aarnio A, Hakkinen R, Junttila O (2006) Critical night length for bud set and its variation in two photoperiodic ecotypes of *Betula pendula*. *Tree Physiology*, **26**, 1013–1018.

Vitasse Y, Lenz A, Hoch G, Körner C (2014a) Earlier leaf-out rather than difference in freezing resistance puts juvenile trees at greater risk of damage than adult trees. *Journal of Ecology*, **102**, 981–988. doi:10.1111/1365-2745.12251.

Vitasse Y, Lenz A, Körner C (2014b) The interaction between freezing tolerance and phenology in temperate deciduous trees. *Frontiers in Plant Science*, **5**, 541. doi:10.3389/fpls.2014.00541. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4192447&tool=pmcentrez&rendertype=abstract>.

Vitra A, Lenz A, Vitasse Y (2017) Frost hardening and dehardening potential in temperate trees from winter to budburst. *New Phytologist*, **216**, 113–123. doi:10.1111/nph.14698. URL <http://dx.doi.org/10.1111/nph.14698>.

Way DA, Montgomery RA (2015) Photoperiod constraints on tree phenology, performance and migration in a warming world. *Plant, Cell & Environment*, **38**, 1725–1736. doi:10.1111/pce.12431. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/pce.12431>.

White MA, De Beurs KM, Didan K, *et al.* (2009) Intercomparison, interpretation, and assessment of spring phenology in North America estimated from remote sensing for 1982–2006. *Global Change Biology*, **15**, 2335–2359. doi:10.1111/j.1365-2486.2009.01910.x. URL <http://dx.doi.org/10.1111/j.1365-2486.2009.01910.x>.

Xie Y, Wang X, Silander JA (2015) Deciduous forest responses to temperature, precipitation, and drought imply complex climate change impacts. *Proceedings of the National Academy of Sciences*, **112**, 13585–13590. doi:10.1073/pnas.1509991112. URL <http://dx.doi.org/10.1073/pnas.1509991112>.

- 422 Xin Q (2016) A risk-benefit model to simulate vegetation spring onset in response to multi-decadal climate  
423 variability: Theoretical basis and applications from the field to the Northern Hemisphere. *Agriculture and*  
424 *Forest Meteorology*, **228-229**, 139–163.
- 425 Zohner CM, Benito BM, Svenning JC, Renner SS (2016) Day length unlikely to constrain climate-driven  
426 shifts in leaf-out times of northern woody plants. *Nature Climate Change*, **6**, 1120–1123. doi:10.1038/  
427 nclimate3138. URL <http://dx.doi.org/10.1038/NCLIMATE3138>.