

# Tree seasonality in a warming climate

Heikki Hänninen<sup>1</sup> and Karen Tanino<sup>2</sup>

<sup>1</sup> Plant Ecophysiology and Climate Change Group (PECC), Department of Biosciences, University of Helsinki, Finland

<sup>2</sup> Department of Plant Sciences, College of Agriculture and Bioresources, University of Saskatchewan, Saskatoon, SK, Canada

**Climate warming has increased researchers' interest in plant phenology and its modelling. Although the main focus is on projections of accelerated springtime phenological events, also a further extension of the growing season by delayed growth cessation is often projected. However, ecophysiological studies indicate that, for boreal and temperate trees, such generalisations are precluded owing to differential climatic conditions and inter- and intraspecific genetic differences. The annual cycle of these trees is an integrated system, where one phase affects subsequent phases, resulting in delayed impacts, which are only partially addressed in current ecophysiological models. Here, we outline an updated integrated conceptual model of the annual cycle by identifying ecophysiological phenomena that are particularly significant under climate warming.**

## Neglected impacts of the annual cycle

Climate warming has increased the attention paid to plant phenology and the duration of the growing season [1–4]. Changes in the phenological patterns caused by climatic warming can have major implications for both the global carbon cycle [5] and the ecology of plants [6] and their herbivores [7]. The main research focus has been on the acceleration of springtime phenological events of plants (e.g. bud burst in boreal and temperate trees), which can be caused by climate warming [8–11]. Such acceleration will have considerable implications for both the risk of frost damage [12–15] and the photosynthetic production [5,16,17] of the trees, with the photosynthetic production also feeding back to the climate system [2]. Somewhat less attention has been paid to the effects of climate warming on the growth cessation and senescence of plants in autumn; mainly, a prolongation of the growing season is often projected [2,10,18].

However, as emphasised recently, because of genetic interactions with chilling temperatures and photoperiod, climate warming will not accelerate the bud burst and growth onset in all boreal and temperate tree species [3]. A recent transplantation study carried out with three European *Quercus* species suggested that, even when bud burst and growth onset are accelerated by climatic warming, the acceleration response can be nonlinear, so that the acceleration levels off with increased levels of warming [19]. Furthermore, a recent review reveals that warming causes acceleration, rather than a delay, of growth cessation in many tree species, ecotypes and cultivars [20]. The projections of the effects of climate warming are further complicated by intraspecific genetic variation in the

environmental regulation of the annual cycle, which is related to the large geographical distribution areas of many boreal and temperate tree species [20,21]. Owing to these differences, not only the intensity, but also the direction of the response to warming can vary among ecotypes, so that warming accelerates the growth cessation in some and delays it in other ecotypes of a given species. To our knowledge, such differences among ecotypes have not been addressed in global change modelling.

It has been known for a long time that the annual cycle of growth and dormancy in boreal and temperate trees forms an integrated system, where one phase in the cycle affects the subsequent phases, so that environmental factors have several delayed impacts [22–28]. Many of these delayed impacts have been addressed in process-based ecophysiological simulation models [29,30] often used in scenario studies [6,13,14,31–33], but many remain unaddressed. Here, we outline an updated integrated conceptual model for the annual cycle of boreal and temperate trees by identifying phenomena of particular significance under climate warming.

## Bud burst and growth onset

Cell division and growth in the buds are temperature dependent, so that the rate of development towards bud burst and growth onset increases with rising air temperatures [26,27]. This is the physiological basis of the common projection of accelerated springtime phenological events in plants, often deduced in computer simulations by the accumulation of day degrees or other types of temperature sum (Figure 1) [12–14,31–33].

## Glossary

**Chilling:** exposure to low non-freezing temperatures required for rest break (see below) in both vegetative and reproductive buds.

**Dormancy:** general term indicating the opposite of growth; that is, a condition where no visible growth is observable in the bud.

**Fixed growth habit:** height growth pattern where growth consists entirely of the lengthening of the primordia formed in the bud during the previous growing season.


**Free growth habit:** height growth pattern where the lengthening of new primordia developed during the same growing season also contributes to height growth.

**Process-based model:** a simulation model where the development of a plant or an entire ecosystem is described by explicitly addressing physiological and biophysical phenomena.

**Rest:** special case of dormancy where growth is arrested by physiological conditions in the bud, so that the bud remains dormant regardless of the environmental conditions.

**Rest break:** removal of the growth-arresting physiological conditions in the bud, caused in most cases by the accumulation of chilling. After rest completion, the bud can remain dormant for several months if the air temperatures remain low.

Corresponding author: Hänninen, H. (heikki.hanninen@helsinki.fi).



**Figure 1.** An integrated conceptual model of the annual cycle of growth and dormancy in boreal and temperate trees, identifying the ecophysiological phenomena that determine the timing of growth onset and cessation. In addition to their short-term effects, air temperature and photoperiod have several delayed impacts on the timing of these phenological events. As depicted by the arrows for the periods of growth (red) and dormancy (blue), the delayed impacts are, in many cases, carried over for several months in the annual cycle. The alternation between growth and dormancy occurs gradually (dashed arrows). (i) Long-term accumulation of chilling during late autumn and early winter causes rest break (i.e. removal of the growth-arresting physiological conditions in the bud). In some cases, long photoperiods also affect rest break. (ii) After rest completion, growth onset (e.g. bud burst) is brought on by the cumulative effects of high temperatures, as described by temperature-sum models. (iii) Temperature sum accumulation regulates height growth and, in tree species with a fixed growth habit, also its cessation. (iv) In most species with a free growth habit, a short photoperiod is an essential environmental cue inducing growth cessation. However, several interactions with air temperature have been found, so that growth cessation might even be affected by earlier temperatures (i.e. those occurring before and during growth phase). In some species and ecotypes, growth cessation is induced by low night temperatures independently of the photoperiod. (v) Dormancy is induced by short photoperiods. High night temperatures during the induction increase the depth of dormancy by increasing the chilling requirement of rest completion and/or the temperature sum requirement of growth onset in some species. By contrast, low night temperatures advance dormancy in northern ecotypes under long photoperiods and in certain fruit cultivars that are insensitive to the photoperiod.

However, this projection might be complicated in several ways by dormancy (see [Glossary](#)), a typical feature of the buds of boreal and temperate trees [26,27,34–36]. Rest break, (i.e. the removal of growth-arresting conditions in the buds) requires prolonged exposure to chilling temperatures, typically several weeks (Figure 1). Trees with a low chilling requirement burst buds and commence growth in spring at an earlier date than do those with a high chilling requirement, the latter having a greater depth of rest [37,38]. This delayed effect of chilling temperatures on the timing of bud burst is projected to cause different acceleration responses to warming in boreal trees, depending on the chilling requirement: in trees with a low chilling requirement, warming accelerates bud burst more than in trees with a high chilling requirement [13,39].

In a few cases, the dormancy phenomenon and the related chilling requirement are projected to cause differences among trees in the timing of bud burst in terms of not only the intensity, but also the direction of the response to warming. The chilling required for bud burst will be delayed and, in milder climates, also reduced by climate warming. As shown by process-based ecophysiological modelling [12], this delaying effect of elevated air temperatures in autumn and winter can be carried over to the

spring (Figure 1). In mild and maritime climates, warming is thus projected to delay bud burst in some species, such as balsam poplar (*Populus trichocarpa*) and common hawthorn (*Crataegus monogyna*) [31]. Furthermore, process-based modelling has revealed that the lack of chilling caused by climate warming can cause abnormal bud burst in some temperate species [33]. This aspect has also been demonstrated experimentally under controlled conditions by using artificial rest breaks at different stages of dormancy [40].

Recently, a delay of growth onset as a result of warm winters was also found in native meadow and steppe vegetation in natural conditions in the Tibetan Plateau of western China, and the delay was attributed to insufficient chilling [41]. These pioneering satellite-based observations suggest that the delay of growth onset caused by climatic warming in winter is not restricted to woody plants. The satellite observations from the Tibetan Plateau [41] mark the importance of experimental studies addressing the dormancy phenomena in herbaceous plants.

There is increasing experimental evidence that elevated air temperatures during dormancy induction in late summer and early autumn increase the depth of dormancy, so that more chilling is required for rest break and/or more

accumulation of temperature sum for bud burst (Figure 1) [20,42–47]; some of the studies have made comparisons and found distinct ecotype- or cultivar-dependent responses. Thus, besides the impact of reduced chilling being carried over from the previous autumn and winter, the impacts of elevated air temperatures, with potentially delayed bud burst in spring, could be carried over from an even earlier time (i.e. from the previous summer; Figure 1). These phenomena and the differential ecotype- or cultivar-dependent responses are not addressed in current process-based ecophysiological models ([29,30]; however, see [48]).

Half a century ago, an experimental study revealed complicated interactions and delayed effects in the annual cycle of native sugar maple (*Acer saccharum*) [22]. The author concluded that the differential depth of dormancy was associated with the previous growing season and, in particular, the photoperiod prevailing after bud burst. When plants were grown in a 20-h photoperiod, a second flush of growth was produced, as opposed to only one flush in a 9-h photoperiod. The plants grown in the 20-h photoperiod with two flushes of growth over spring and summer subsequently evinced a greater depth of dormancy in the following autumn and winter than did those grown in the 9-h photoperiod. Although 20-h and 9-h photoperiods are extreme treatments, these results suggest a potentially complicated case of delayed interactive effects of temperature and photoperiod, which can affect the seasonality of the trees under climate warming. Accordingly, if spring-time bud burst is advanced or delayed for some reason, photoperiod can interact to induce differential spring and summer growth patterns, which will then influence the subsequent rest during the autumn and winter and finally affect the timing of bud burst the next spring. These effects are not addressed in current process-based ecophysiological models [29,30] and, for the sake of simplicity, they are similarly not addressed in the conceptual model of Figure 1.

It has also been suggested that additional rest break requirements based on prevailing light conditions, such as the requirement of a long photoperiod, prevent or slow down the accelerating effects of springtime warming on bud burst even when the chilling requirement is met (Figure 1) [3,48–51]. With the exception of a few species such as beech (*Fagus sylvatica*) [52], however, the role of photoperiod in regulating the timing of bud burst remains unclear [4,30].

### Growth cessation

Since the discovery of photoperiodism during the early 20th century [53], autumn growth cessation in boreal and temperate trees has generally been explained in terms of the attainment of a critical short photoperiod [24,54–57]. This notion suggests that climate warming should cause no change in the timing of growth cessation. Experimental evidence accumulated over several decades, however, points at interactive effects of photoperiod and air temperature on growth cessation in both coniferous and deciduous broad-leaved trees. In most cases, elevated night temperatures have induced earlier growth cessation (Figure 1) [20,42,44–47,58–64]. In tree species with a fixed growth habit, such as Scots pine (*Pinus sylvestris*), height growth

cessation is regulated solely by temperature sum accumulation [65]. Furthermore, in many boreal tree species, an accelerated growth onset is followed by an accelerated growth cessation [65,66], suggesting that the accelerating impact of greater temperature sum accumulation in spring is carried over to the autumn (Figure 1). Collectively, these findings suggest that, contrary to the projection of a prolonged growing season in autumn [2,10,18], climate warming can accelerate growth cessation in many boreal and temperate trees. This scenario might be further substantiated by the observation that daily minimum temperatures have been increasing more than have daily maximums during the past few decades [67]. If this trend continues, then night temperatures will generally increase more than day temperatures; it is the high night temperatures in particular that have been shown to be an essential interactive factor accelerating tree growth cessation (Figure 1, see above).

In some trees, however, low night temperatures induce growth cessation (Figure 1). In most of these cases, growth cessation occurs in combination with a long photoperiod. This group of trees appears to consist of northern ecotypes and species whose dormancy induction is insensitive to the photoperiod. It is exemplified by northern ecotypes of Norway spruce (*Picea abies*), pubescent birch (*Betula pubescens*), red osier dogwood (*Cornus sericea*), laurel willow (*Salix pentandra*), and several species of the Rosaceae family [20,60,62,68–73]. In these species and ecotypes, climate warming is projected to cause an extension of the growing season in the autumn through delayed growth cessation. Furthermore, according to a recent modelling study, low temperatures accelerate the autumn colouration in three European broad-leaved tree species after the colouration has been induced by a short photoperiod [74]; thus, climatic warming is also projected to delay the autumn phenology of the trees in this case.

### Concluding remarks

Climate warming is likely to accelerate bud burst and growth onset in most, but not all, boreal and temperate trees. Growth cessation might be either accelerated or delayed by warming, depending on the species and even on the ecotype. The differences among tree species, ecotypes and cultivars in their phenological responses to warming can have crucial impacts on the structure and functioning of boreal and temperate forest ecosystems, thus contributing to an overall multitude of ecological responses to climate change. The differences will also have important implications for practical forestry and horticulture, such as the need to reassess the suitability of a given tree provenance or cultivar used earlier in forest regeneration or fruit production.

The take-home message of the integrated conceptual model is that one part of the annual cycle impacts subsequent parts and that individual seasonal responses should not be examined in isolation without taking into account the history of the plant. Thus, because of several delayed impacts of environmental factors, any projections of the effects of climate warming on the phenology of boreal and temperate trees need to consider the entire annual cycle. Current process-based ecophysiological simulation models

of the annual cycle [29,30] provide a useful tool for generating quantitative projections for the trees [14,31–33] and for scaling up these projections to higher organisational levels, ranging from stands and ecosystems to entire biogeographical areas [75–78]. For reliable projections of phenological changes, however, further species-, ecotype- and even cultivar-specific experimental data are needed in order to develop these models further and to better understand the underlying ecophysiology summarised in Figure 1.

### Acknowledgements

This study was supported by the Academy of Finland (Project 122194), and the Agroforestry Division, Agriculture and Agri-Food Canada. We wish to thank Risto Häkkinen and Robin Lundell for comments on an earlier version of the manuscript, Pekka Hirvonen for linguistic revision of the text, and Olavi Junttila for insightful discussions.

### References

- Steltzer, H. and Post, E. (2009) Seasons and life cycles. *Science* 324, 886–887
- Peñuelas, J. *et al.* (2009) Phenology feedbacks on climate change. *Science* 324, 887–888
- Körner, C. and Basier, D. (2010) Phenology under global warming. *Science* 327, 1461–1462
- Chuine, I. *et al.* (2010) Warming, photoperiods, and tree phenology. *Science* 329, 277–278
- Leinonen, I. and Kramer, K. (2002) Applications of phenological models to predict the future carbon sequestration potential of boreal forests. *Clim. Change* 55, 99–113
- Morin, X. *et al.* (2008) Tree species range shifts at a continental scale: new predictive insights from a process-based model. *J. Ecol.* 96, 784–794
- Visser, M.E. and Holleman, L.J.M. (2001) Warmer springs disrupt the synchrony of oak and winter moth phenology. *Proc. R. Soc. Lond. B: Biol. Sci.* 268, 289–294
- Kramer, K. *et al.* (2000) The importance of phenology for the evaluation of impact of climate change on growth of boreal, temperate and Mediterranean forests ecosystems: an overview. *Int. J. Biometeorol.* 44, 67–75
- Beaubien, E.G. and Freeland, H.J. (2000) Spring phenology events in Alberta, Canada: links to ocean temperature. *Int. J. Biometeorol.* 44, 53–59
- Menzel, A. *et al.* (2006) European phenological response to climate change matches the warming pattern. *Glob. Change Biol.* 12, 1969–1976
- Linkosalo, T. *et al.* (2009) The time series of flowering and leaf bud burst of boreal trees (1846–2005) support the direct temperature observations of climatic warming. *Agric. Forest Meteorol.* 149, 453–461
- Cannell, M.G.R. and Smith, R.I. (1986) Climatic warming, spring budburst and frost damage on trees. *J. Appl. Ecol.* 23, 177–191
- Hänninen, H. (1991) Does climatic warming increase the risk of frost damage in northern trees? *Plant Cell Environ.* 14, 449–454
- Hänninen, H. (2006) Risk of frost damage to boreal forest trees under climatic warming: identification of critical ecophysiological traits. *Tree Physiol.* 26, 889–898
- Gu, L. *et al.* (2008) The 2007 eastern US spring freeze: increased cold damage in a warming world? *BioScience* 58, 253–262
- Ensminger, I. *et al.* (2009) Conifer cold hardiness, climate change and the likely effects of warmer temperatures on photosynthesis. In *Plant Cold Hardiness: From the Laboratory to the Field* (Gusta, L.V. *et al.*, eds), pp. 249–261, CABI
- Richardson, A.D. *et al.* (2010) Influence of spring and autumn phenological transitions on forest ecosystem productivity. *Proc. R. Soc. Lond. B: Biol. Sci.* 365, 3227–3246
- Defila, C. and Clot, B. (2001) Phenological trends in Switzerland. *Int. J. Biometeorol.* 45, 203–207
- Morin, X. *et al.* (2010) Changes in leaf phenology of three European oak species in response to experimental climate change. *New Phytol.* 186, 900–910
- Tanino, K. *et al.* (2010) Temperature-driven plasticity in growth cessation and dormancy development in deciduous woody plants: a working hypothesis suggesting how molecular and cellular function is affected by temperature during dormancy induction. *Plant Mol. Biol.* 73, 49–65
- Rohde, A. *et al.* (2011) Bud set in poplar – genetic dissection of a complex trait in natural and hybrid populations. *New Phytol.* 189, 106–121
- Olmsted, C.E. (1954) Experiments on photoperiodism, dormancy and leaf age and abscission in sugar maple. *Bot. Gaz.* 112, 365–393
- Vegis, A. (1964) Dormancy in higher plants. *Annu. Rev. Plant Physiol.* 15, 185–224
- Weiser, C.J. (1970) Cold resistance and injury in woody plants. *Science* 169, 1269–1278
- Perry, T.O. (1971) Dormancy of trees in winter. *Science* 171, 29–36
- Sarvas, R. (1972) Investigations on the annual cycle of development of forest trees, Active period. *Communicationes Instituti Forestalis Fenniae* 76, 1–110
- Sarvas, R. (1974) Investigations on the annual cycle of development of forest trees, II. Autumn dormancy and winter dormancy. *Communicationes Instituti Forestalis Fenniae* 84, 1–101
- Fuchigami, L.H. *et al.* (1977) The relationship of vegetative maturity to rest development and spring bud-break. *J. Am. Soc. Hort. Sci.* 102, 450–452
- Chuine, I. *et al.* (2003) Plant development models. In *Phenology: An Integrative Environmental Science* (Schwartz, M.D., ed.), pp. 305–333, Kluwer Academic
- Hänninen, H. and Kramer, K. (2007) A framework for modelling the annual cycle of trees in boreal and temperate regions. *Silva Fenn.* 41, 167–205
- Murray, M.B. *et al.* (1989) Date of budburst of fifteen tree species in Britain following climatic warming. *J. Appl. Ecol.* 26, 693–700
- Kramer, K. (1994) A modelling analysis of the effects of climatic warming on the probability of spring frost damage to tree species in The Netherlands and Germany. *Plant Cell Environ.* 17, 367–377
- Morin, X. *et al.* (2009) Leaf phenology in 22 North American tree species during the 21st century. *Glob. Change Biol.* 15, 961–975
- Fuchigami, L.H. *et al.* (1982) A degree growth stage (GS) model and cold acclimation in temperate woody plants. In *Plant Cold Hardiness and Freezing Stress* (Vol. 2) (Li, P.H. and Sakai, A., eds), pp. 93–116, Academic Press
- Rohde, A. and Bhalerao, R.P. (2007) Plant dormancy in the perennial context. *Trends Plant Sci.* 12, 217–223
- Ruttink, T. *et al.* (2007) A molecular timetable for apical bud formation and dormancy induction in poplar. *Plant Cell* 19, 2370–2390
- Powell, L.E. (1986) The chilling requirement in apple and its role in regulating time of flowering in spring in cold-winter climates. *Acta Horticulturae* 179, 129–140
- Hänninen, H. (1990) Modelling bud dormancy release in trees from cool and temperate regions. *Acta Forestalia Fennica* 213, 1–47
- Linkosalo, T. (2000) Analyses of the spring phenology of boreal trees and its response to climate change. *University of Helsinki Department of Forest Ecology Publications* 22, 1–55
- Nee, C.-C. and Fuchigami, L.H. (1992) Overcoming rest at different growth stages with hydrogen cyanamide. *Scientia Horticulturae* 50, 107–113
- Yu, H. (2010) Winter and spring warming result in delayed spring phenology on the Tibetan plateau. *Proc. Natl. Acad. Sci. U.S.A.* 107, 22151–22156
- Dormling, I. (1989) The role of photoperiod and temperature in the induction and release of dormancy in *Pinus sylvestris* L. seedlings. *Ann. Forest Sci.* 46, 228–232
- Westergaard, L. and Eriksen, E.N. (1997) Autumn temperature affects the induction of dormancy in first-year seedlings of *Acer platanoides* L. *Scand. J. Forest Res.* 12, 11–16
- Heide, O.M. (2003) High autumn temperature delays spring bud burst in boreal trees, counterbalancing the effect of climatic warming. *Tree Physiol.* 23, 931–936
- Sogaard, G. *et al.* (2008) Climatic control of bud burst in young seedlings of nine provenances of Norway spruce. *Tree Physiol.* 28, 311–320
- Granhus, A. *et al.* (2009) Bud burst timing in *Picea abies* seedlings as affected by temperature during dormancy induction and mild spells during chilling. *Tree Physiol.* 29, 497–503



- 47 Kalcsits, L. *et al.* (2009) Warm temperature accelerates short photoperiod induced growth cessation and dormancy induction in hybrid poplar (*Populus x spp.*). *Trees* 23, 973–979
- 48 Caffarra, A. *et al.* (2011) Modelling the timing of *Betula pubescens* budburst, II. Integrating complex effects of photoperiod into process-based models. *Clim. Res.* 46, 159–170
- 49 Linkosalo, T. *et al.* (2000) Predicting spring phenology and frost damage risk of *Betula* spp. under climatic warming: a comparison of two models. *Tree Physiol.* 20, 1175–1182
- 50 Caffarra, A. *et al.* (2011) Modelling the timing of *Betula pubescens* budburst, I. Temperature and photoperiod: a conceptual model. *Clim. Res.* 46, 147–157
- 51 Caffarra, A. and Donnelly, A. (2010) The ecological significance of phenology in four different tree species: effects of light and temperature on bud burst. *Int. J. Biometeorol.* DOI: 10.1007/s00484-010-0386-1
- 52 Heide, O.M. (1993) Dormancy release in beech buds (*Fagus sylvatica*) requires both chilling and long days. *Physiol. Plant* 89, 187–191
- 53 Garner, W.W. and Allard, H.A. (1923) Further studies in photoperiodism, the response of the plant to the relative length of day and night. *J. Agric. Res.* 23, 871–920
- 54 Kramer, P.J. (1936) Effect of variation in length of day on growth and dormancy of trees. *Plant Physiol.* 11, 127–137
- 55 Vaartaja, O. (1954) Photoperiodic ecotypes of trees. *Can. J. Bot.* 32, 392–399
- 56 Wareing, P.F. (1956) Photoperiodism in woody plants. *Annu. Rev. Plant Physiol.* 7, 191–214
- 57 Nitsch, J.P. (1957) Growth responses of woody plants to photoperiodic stimuli. *Proc. Am. Soc. Hort. Sci.* 70, 512–525
- 58 Van der Veen, R. (1951) Influence of daylength on the dormancy of some species of the genus *Populus*. *Physiol. Plant.* 4, 35–40
- 59 Kramer, P.J. (1957) Some effects of various combinations of day and night temperatures and photoperiod on the height growth of loblolly pine seedlings. *Forest Sci.* 3, 45–55
- 60 Håbørg, A. (1972) Effects of photoperiod and temperature on growth and development of three latitudinal and three altitudinal populations of *Betula pubescens* Ehrh. *Scientific Reports of the Agricultural University of Norway* 51(2), 1–27
- 61 Heide, O.M. (1974) Growth and dormancy in Norway spruce ecotypes (*Picea abies*), I. Interaction of photoperiod and temperature. *Physiol. Plant* 30, 1–12
- 62 Junttila, O. (1980) Effect of photoperiod and temperature on apical growth cessation in two ecotypes of *Salix* and *Betula*. *Physiol. Plant* 48, 347–352
- 63 Downs, R.J. and Bevington, J.M. (1981) Effect of temperature and photoperiod on growth and dormancy of *Betula papyrifera*. *Am. J. Bot.* 68, 795–800
- 64 Mölmann, J.A. *et al.* (2005) Low night temperature and inhibition of gibberellin biosynthesis override phytochrome action and induce bud set and cold acclimation, but not dormancy in PHYA overexpressors and wild-type of hybrid aspen. *Plant Cell Environ.* 28, 1579–1588
- 65 Koski, V. and Sievänen, R. (1985) Timing of growth cessation in relation to the variations in the growing season. In *Crop Physiology of Forest Trees* (Tigerstedt, P.M.A. *et al.*, eds), pp. 167–193, Helsinki University Press
- 66 Viherä-Aarnio, A. *et al.* (2005) Effects of seed origin and sowing time on timing of height growth cessation of *Betula pendula* seedlings. *Tree Physiol.* 25, 101–108
- 67 Vose, R.S. *et al.* (2005) Maximum and minimum temperature trends for the globe: an update through 2004. *Geophys. Res. Lett.* 32, L23822 DOI: 10.1029/2005GL024379
- 68 Dormling, I. *et al.* (1968) The experimental control of the life cycle in *Picea abies* (L.) Karst. *Silvae Gen.* 17, 44–64
- 69 Heide, O.M. and Pestrud, A.K. (2005) Low temperature, but not photoperiod, controls growth cessation and dormancy induction and release in apple and pear. *Tree Physiol.* 25, 109–114
- 70 Junttila, O. (2007) Regulation of annual shoot growth cycle in northern tree species. In *Physiology of Northern Plants Under Changing Environment* (Taulavuori, E. and Taulavuori, K., eds), pp. 177–210, Res. Signpost
- 71 Svendsen, E. *et al.* (2007) A molecular marker associated with low-temperature induction of dormancy in red osier dogwood (*Cornus sericea*). *Tree Physiol.* 27, 385–397
- 72 Heide, O.M. (2008) Interaction of photoperiod and temperature in the control of growth and dormancy of *Prunus* species. *Scientia Horticulturae* 115, 309–314
- 73 Junttila, O. (1982) The cessation of apical growth in latitudinal ecotypes and ecotype crosses of *Salix pentandra* L. *J. Exp. Bot.* 33, 1021–1029
- 74 Delpierre, N. *et al.* (2009) Modelling interannual and spatial variability of leaf senescence for three deciduous tree species in France. *Agric. Forest Meteorol.* 149, 938–948
- 75 Chuine, I. and Beaubien, E.G. (2001) Phenology is a major determinant of tree species range. *Ecol. Lett.* 4, 500–510
- 76 Kramer, K. *et al.* (2002) Evaluation of 6 process-based forest growth models based on eddy-covariance measurements of CO<sub>2</sub> and H<sub>2</sub>O fluxes at 6 forest sites in Europe. *Glob. Change Biol.* 8, 213–230
- 77 Hänninen, H. *et al.* (2005) Seasonal development of Scots pine under climatic warming: effects on photosynthetic production. *Can. J. Forest Res.* 35, 2092–2099
- 78 Kramer, K. and Hänninen, H. (2009) The annual cycle of development of trees and process-based modelling of growth to scale up from the tree to the stand. In *Phenology of Ecosystem Processes* (Noormets, A., ed.), pp. 201–227, Springer