

# Rethinking False Spring Risk

C. J. Chamberlain <sup>1,2</sup>, E. M. Wolkovich <sup>1,2</sup>

June 5, 2017

## Introduction

### 1. Introduce False Spring Concept

- (a) Plants growing in temperate environments are at risk of being exposed to late spring freezes, which can be detrimental to growth.
- (b) Individuals that leaf out before the last frost are at risk of leaf loss, damaging wood tissue, and slowed or stalled canopy development (Gu *et al.*, 2008; Hufkens *et al.*, 2012).
- (c) Therefore, temperate deciduous tree species must have plastic phenological responses in the spring in order to optimize photosynthesis and minimize frost or drought risk (Polgar & Primack, 2011).
- (d) These late spring freezing events are known as false springs. False spring events can result in highly adverse ecological and economic consequences (Knudson, 2012; Ault *et al.*, 2013).

### 2. Introduce Climate Change and Importance of False Spring Studies

- (a) Climate change is expected to increase damage from false spring events around the world due to earlier spring onset and greater fluctuations in temperature (Cannell & Smith, 1986; Inouye, 2008; Martin *et al.*, 2010).
- (b) Temperate forest species around the world are initiating leaf out about 4.6 days earlier per degree Celsius (Wolkovich *et al.*, 2012; Polgar *et al.*, 2014).
- (c) It is anticipated that there will be a decrease in false spring frequency overall but the magnitude of temperature variation is likely to increase, therefore amplifying the expected intensity of false spring events (Kodra *et al.*, 2011; Allstadt *et al.*, 2015).
- (d) Multiple studies have documented false spring events in recent years (Gu *et al.*, 2008; Augspurger, 2009; Knudson, 2012; Augspurger, 2013) and some have linked this to climate change (Ault *et al.*, 2013; Allstadt *et al.*, 2015; Muffler *et al.*, 2016; Xin, 2016).

- (e) Due to these reasons, it is crucial for researchers to properly evaluate the effects of false spring events on temperate forests and agricultural crops in order to make more accurate predictions on future trends.

### 3. Introduce Current False Spring Index Equation

- (a) Different species respond differently to late spring freezing events.
- (b) The level of damage sustained by plants from a false spring also varies across phenophases.
- (c) Various studies have assessed the risk of damage or the intensity of particular false spring events but at this time false spring studies fail to incorporate all potential factors that could affect the level of frost damage risk.
- (d) A False Spring Index (FSI) signifies the likelihood of a damage to occur from a late spring freeze.
- (e) Currently, FSI evaluates day of budburst, number of growing degree days, and day of last spring freeze through a simple equation as seen below (Marino *et al.*, 2011).

$$FSI = JulianDate(LastSpringFreeze) - JulianDate(Budburst) \quad (1)$$

- (f) False spring studies largely simplify the various ecological elements that could predict the level of plant damage from late spring freezing events.
- (g) In contrast to these simplifications, we argue that a wealth of factors greatly impacts plants' frost spring risk such that simple indices will most likely lead to inaccurate predictions and ultimately do little to advance the field.

### 4. State the Purpose of the Paper

- (a) In this paper we aim to highlight the complexity of factors driving a plant's false spring risk.
- (b) We outline in particular how life stage of the individual (Caffarra & Donnelly, 2011), location within a forest or canopy (Augspurger, 2013), winter chilling hours (Flynn & Wolkovich 2017?), proximity to water (Gu *et al.*, 2008), level of precipitation prior to the freezing event (Anderegg *et al.*, 2013), freeze duration/intensity, and range limits of the species (Martin *et al.*, 2010) unhinge simple metrics of false spring.
- (c) The ultimate intent is to demonstrate how an integrated view of false spring that incorporates these factors would rapidly advance progress in this field.

## Defining False Spring

### 1. Definition and Threat

- (a) Episodic spring frosts present the most risk to frost damage for temperate forest plants (Sakai & Larcher, 1987).
- (b) Abnormally warm conditions in the late winter or early spring can cause budburst to initiate early in trees and shrubs.
- (c) Freezing temperatures following a warm spell could result in plant damage or even death (Ludlum, 1968; Mock *et al.*, 2007).
- (d) False springs are defined by two phases: rapid vegetative growth prior to a freeze and a post freeze setback (Gu *et al.*, 2008).
- (e) Freeze and thaw fluctuations can cause defoliation, xylem embolism and decreased xylem conductivity which can result in crown dieback (Gu *et al.*, 2008).
- (f) Species that are better able to phenologically track the shifts in spring advancement due to climate change are more likely to sustain damaging events such as false springs (Scheifinger *et al.*, 2003).

## 2. Define Chilling requirements to specify timing of damaging false spring events

- (a) Deciduousness and the evolution of two dormancy phases (i.e. endodormancy and ecodormancy) in temperate forest trees has permitted species to occupy more northern ecological niches (Samish, 1954).
- (b) Endodormancy is the period of winter when temperate trees are inhibited from growing, regardless of the outdoor environment.
- (c) Ecodormancy is the period of time when growth can occur but the external environment is not conducive to growth (e.g. too cold) (Basler & Körner, 2012).
- (d) Therefore, warm temperatures earlier in the year (i.e. in February) do not seem to affect species, most likely because trees have not yet left the endodormancy phase.
- (e) Frost damage usually occurs when there is a warmer than average March, a freezing April, and enough growing degree days between budburst and the last freeze date (Augspurger, 2013).
- (f) A damaging false spring is currently defined as having more than 7 days between budburst and the last freeze date, using Equation 1 (Peterson & Abatzoglou, 2014).
- (g) The 7 day parameter exposes less resistant foliate phenophases to a false spring, thus putting the plant at a higher risk of damage.
- (h) Once budburst has initiated, buds cannot respond to cold temperatures and freeze resistance is greatly reduced (Taschler *et al.*, 2004; Lenz *et al.*, 2013; Vitasse *et al.*, 2014).
- (i) There are two types of freezes: a “hard freeze” at  $-2.2^{\circ}\text{C}$  and a “soft freeze” at  $-1.7^{\circ}\text{C}$  (Vavrus *et al.*, 2006; Kodra *et al.*, 2011; Augspurger, 2013).

- (j) However, the definition is still largely under debate.

### 3. Damage and drought

- (a) Freezing damage can occur directly via intracellular ice formation or indirectly via freezing dehydration (Pearce, 2001; Beck *et al.*, 2004; Hofmann & Bruelheide, 2015).
- (b) Intracellular ice formation often results in defoliation and increased xylem cavitation or embolism in the stem.
- (c) Freezing tolerance in plants is usually against extracellular freezing or freezing dehydration (Burke *et al.*, 1976).
- (d) Drought and desiccation within the xylem mimic the adverse effects of false spring events (Cavender-Bares *et al.*, 2015).
- (e) Dry winters typically result in new, frost-tolerant shoots due to the decreased water content and osmotic potential from the reduced number of accumulated solutes (Morin *et al.*, 2007; Hofmann & Bruelheide, 2015).
- (f) Therefore, it is hypothesized that increased bud dehydration results in increased frost hardiness (Beck *et al.*, 2007; Norgaard Nielsen & Rasmussen, 2009; Poirier *et al.*, 2010; Kathke & Bruelheide, 2011; Hofmann & Bruelheide, 2015).
- (g) However, more studies are needed to investigate the interplay between false spring events and precipitation and how that relationship impacts the level of damage a plant sustains.

## References

- Allstadt, A.J., Vavrus, S.J., Heglund, P.J., Pidgeon, A.M., Wayne, E. & Radeloff, V.C. (2015) Spring plant phenology and false springs in the conterminous U. S. during the 21st century. *Environmental Research Letters (submitted)* **10**, 104008.
- Anderegg, W.R.L., Plavcová, L., Anderegg, L.D.L., Hacke, U.G., Berry, J.A. & Field, C.B. (2013) Drought's legacy: Multiyear hydraulic deterioration underlies widespread aspen forest die-off and portends increased future risk. *Global Change Biology* **19**, 1188–1196.
- Augspurger, C.K. (2009) Spring 2007 warmth and frost: Phenology, damage and refoliation in a temperate deciduous forest. *Functional Ecology* **23**, 1031–1039.
- Augspurger, C.K. (2013) Reconstructing patterns of temperature, phenology, and frost damage over 124 years: Spring damage risk is increasing. *Ecology* **94**, 41–50.

- Ault, T.R., Henebry, G.M., de Beurs, K.M., Schwartz, M.D., Betancourt, J.L. & Moore, D. (2013) The False Spring of 2012, Earliest in North American Record. *Eos, Transactions American Geophysical Union* **94**, 181–182.
- Basler, D. & Körner, C. (2012) Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agricultural and Forest Meteorology* **165**, 73–81.
- Beck, E.H., Fettig, S., Knake, C., Hartig, K. & Bhattarai, T. (2007) Specific and unspecific responses of plants to cold and drought stress. *Journal of Biosciences* **32**, 501–510.
- Beck, E.H., Heim, R. & Hansen, J. (2004) Plant resistance to cold stress: Mechanisms and environmental signals triggering frost hardening and dehardening. *Journal of Biosciences* **29**, 449–459.
- Burke, M., Gusta, L., Quamme, H., Weiser, C. & Li, P. (1976) Freezing and injury in plants. *Annual Review of Plant Physiology* **27**, 507–528.
- Caffarra, A. & Donnelly, A. (2011) The ecological significance of phenology in four different tree species: Effects of light and temperature on bud burst. *International Journal of Biometeorology* **55**, 711–721.
- Cannell, M. & Smith, R. (1986) Climatic Warming , Spring Budburst and Forest Damage on Trees Author ( s ): M . G . R . Cannell and R . I . Smith Published by : British Ecological Society Stable URL : <http://www.jstor.org/stable/2403090> JSTOR is a not-for-profit service that helps schol. *Journal of Applied Ecology* **23**, 177–191.
- Cavender-Bares, J., González-Rodríguez, A., Eaton, D.A.R., Hipp, A.A.L., Beulke, A. & Manos, P.S. (2015) Phylogeny and biogeography of the american live oaks (quercussubsectionvirentes): a genomic and population genetics approach. *Molecular Ecology* **24**, 3668–3687.
- Gu, L., Hanson, P.J., Post, W.M., Kaiser, D.P., Yang, B., Nemani, R., Pallardy, S.G. & Meyers, T. (2008) The 2007 Eastern US Spring Freeze: Increased Cold Damage in a Warming World. *BioScience* **58**, 253.
- Hofmann, M. & Bruelheide, H. (2015) Frost hardiness of tree species is independent of phenology and macro-climatic niche. *Journal of Biosciences* **40**, 147–157.
- Hufkens, K., Friedl, M.A., Keenan, T.F., Sonnentag, O., Bailey, A., O’Keefe, J. & Richardson, A.D. (2012) Ecological impacts of a widespread frost event following early spring leaf-out. *Global Change Biology* **18**, 2365–2377.
- Inouye, D.W. (2008) Effects of climate chnage on phenology, frost damage, and floral abundance of montane wildflowers. *Ecology* **89**, 353–362.

- Kathke, S. & Bruelheide, H. (2011) Differences in frost hardiness of two norway spruce morphotypes growing at mt. brocken, germany. *Flora - Morphology, Distribution, Functional Ecology of Plants* **206**, 120–126.
- Knudson, W. (2012) The economic impact of the spring's weather on the fruit and vegetable sectors. *The Strategic Marketing Institute Working Paper* .
- Kodra, E., Steinhäuser, K. & Ganguly, A.R. (2011) Persisting cold extremes under 21st-century warming scenarios. *Geophysical Research Letters* **38**, 1–5.
- Lenz, A., Hoch, G., Vitasse, Y. & Körner, C. (2013) European deciduous trees exhibit similar safety margins against damage by spring freeze events along elevational gradients. *New Phytologist* **200**, 1166–1175.
- Ludlum, D.M. (1968) *Early American Winters: 1604-1820*. 3, American Meteorological Society.
- Marino, G.P., Kaiser, D.P., Gu, L. & Ricciuto, D.M. (2011) Reconstruction of false spring occurrences over the southeastern United States, 1901–2007: an increasing risk of spring freeze damage? *Environmental Research Letters* **6**, 24015.
- Martin, M., Gavazov, K., Körner, C., Hattenschwiler, S. & Rixen, C. (2010) Reduced early growing season freezing resistance in alpine treeline plants under elevated atmospheric co<sub>2</sub>. *Global Change Biology* **16**, 1057–1070.
- Mock, C.J., Mojzisek, J., McWaters, M., Chenoweth, M. & Stahle, D.W. (2007) The winter of 1827–1828 over eastern north america: a season of extraordinary climatic anomalies, societal impacts, and false spring. *Climatic Change* **83**, 87–115.
- Morin, X., Ameglio, T., Ahas, R., Kurz-Besson, C., Lanta, V., Lebourgeois, F., Miglietta, F. & Chuine, I. (2007) Variation in cold hardiness and carbohydrate concentration from dormancy induction to bud burst among provenances of three european oak species. *Tree Physiology* **27**, 817–825.
- Muffler, L., Beierkuhnlein, C., Aas, G., Jentsch, A., Schweiger, A.H., Zohner, C. & Kreyling, J. (2016) Distribution ranges and spring phenology explain late frost sensitivity in 170 woody plants from the northern hemisphere. *Global Ecology and Biogeography* **25**, 1061–1071.
- Norgaard Nielsen, C.C. & Rasmussen, H.N. (2009) Frost hardening and dehardening in abies procera and other conifers under differing temperature regimes and warm-spell treatments. *Forestry* **82**, 43–59.
- Pearce, R. (2001) Plant freezing and damage. *Annals of Botany* **87**, 417–424.
- Peterson, A.G. & Abatzoglou, J.T. (2014) Observed changes in false springs over the contiguous United States. *Geophysical Research Letters* **41**, 2156–2162.

- Poirier, M., Lacointe, A. & Ameglio, T. (2010) A semi-physiological model of cold hardening and dehardening in walnut stem. *Tree Physiology* **30**, 1555–1569.
- Polgar, C., Gallinat, A. & Primack, R.B. (2014) Drivers of leaf-out phenology and their implications for species invasions: Insights from Thoreau’s Concord. *New Phytologist* **202**, 106–115.
- Polgar, C.A. & Primack, R.B. (2011) Leaf-out phenology of temperate woody plants: From trees to ecosystems. *New Phytologist* **191**, 926–941.
- Sakai, A. & Larcher, W. (1987) *Frost Survival of Plants*. Springer-Verlag.
- Samish, R. (1954) Dormancy in woody plants. *Annual Review of Plant Physiology and Plant Molecular Biology* **5**, 183–204.
- Scheifinger, H., Menzel, A., Koch, E. & Peter, C. (2003) Trends of spring time frost events and phenological dates in central europe. *Theoretical and Applied Climatology* **74**, 41–51.
- Taschler, D., Beikircher, B. & Neuner, G. (2004) Frost resistance and ice nucleation in leaves of five woody timberline species measured in situ during shoot expansion. *Tree Physiology* **24**, 331–337.
- Vavrus, S., Walsh, J.E., Chapman, W.L. & Portis, D. (2006) The behavior of extreme cold air outbreaks under greenhouse warming. *International Journal of Climatology* **26**, 1133–1147.
- Vitasse, Y., Lenz, A., Hoch, G. & Körner, C. (2014) Earlier leaf-out rather than difference in freezing resistance puts juvenile trees at greater risk of damage than adult trees. *Journal of Ecology* **102**, 981–988.
- Wolkovich, E.M., Cook, B.I., Allen, J.M., Crimmins, T.M., Betancourt, J.L., Travers, S.E., Pau, S., Regetz, J., Davies, T.J., Kraft, N.J.B., Ault, T.R., Bolmgren, K., Mazer, S.J., McCabe, G.J., McGill, B.J., Parmesan, C., Salamin, N., Schwartz, M.D. & Cleland, E.E. (2012) Warming experiments underpredict plant phenological responses to climate change. *Nature* **485**, 18–21.
- Xin, Q. (2016) A risk-benefit model to simulate vegetation spring onset in response to multi-decadal climate variability: Theoretical basis and applications from the field to the Northern Hemisphere. *Agriculture and Forest Meteorology* pp. 139–163.