

# Rethinking False Spring Risk

Chamberlain, Wolkovich

December 14, 2016

## Introduction

Plants that grow in temperate environments are at risk of being exposed to late spring freezes, which can be detrimental to plant growth. According to Gu *et al.* (2008), there are two phases involved in late spring freezing: rapid vegetative growth prior to the freeze and the post freeze setback. This combined process is known as a false spring. Freeze and thaw fluctuations can cause xylem embolism and decreased xylem conductivity which can result in crown dieback (Gu *et al.*, 2008). More frequently, plants that have been exposed to a false spring will experience leaf loss and slower canopy development (Hufkens *et al.*, 2012). With anthropogenic climate change, the severity of damage incurred from a false spring phenomena is predicted to be heightened due to earlier spring onset and greater fluctuations in temperatures. Temperate forest species around the world are initiating leaf out about 4.6 days earlier per degree Celcius (Polgar *et al.*, 2014; Wolkovich *et al.*, 2012). It is anticipated that there will be a decrease in false spring frequency overall but the severity of temperature variation is likely to increase, therefore amplifying the expected intensity of false spring events (Allstadt *et al.*, 2015). In 2012, a false spring event resulted in half a billion dollars worth of fruit tree damage in Michigan alone (Ault *et al.*, 2013; Knudson, 2012). Due to these reasons, it is crucial for researchers to properly evaluate the effects of false spring events on not only agricultural crops but in temperate forests as well.

Different species exhibit varying responses to late spring freezing events. The level of damage sustained by plants from a false spring also varies across phenophases. Generally, reproductive phases are more sensitive to false spring events than vegetative phases and developing leaves are more susceptible to damage than opening buds or expanding shoots (Peterson & Abatzoglou, 2014). However, trees that suffer severe vegetative growth damage from a false spring event will suffer greater long-term effects than trees that lose one year of reproductive growth due to the loss in photosynthetic tissue in one growing season. False spring events put seedling and sapling trees at greater risk to damage than adult trees (Vitasse *et al.*, 2014). Younger trees

are more likely to incur lastly damage to the leaf buds and vegetative growth, whereas adult trees are at risk of xylem embolism. In order for xylem embolism to occur, extreme cavitation must first occur. This would require more intensive freezing events than it would take to damage seedling and sapling leaf buds. Although, in a study performed by Augspurger (2009), it was noted that some individuals of the same life stage but at different phenophases suffered varying degrees of damage, indicating that phenophase is a greater indicator for level of risk than life stage. Warm temperatures earlier in the year (i.e. in February) do not seem to affect species, most likely because it is too soon for budburst to take place and sufficient chilling has not yet occurred. Frost damage usually occurs when there is a warmer than average March, a freezing April, and enough growing days between the high temperatures and the last freeze date (Augspurger, 2013). In a study performed by Peterson and Abatzoglou (2014), it had been determined that 7 days between budburst and last freeze date is a significant parameter. During this time, it was determined that leaf buds will have enough growing degree days to begin budburst but is not too much time where they are past the leaf out phenophase. There is much debate over the definition of freezing temperatures and has resulted in two types of freezes: a "hard" freeze at  $-2.2^{\circ}\text{C}$  and a "soft" freeze at  $-1.7^{\circ}\text{C}$  (Augspurger, 2013; Kodra *et al.*, 2011; Vavrus *et al.*, 2006).

At this time false spring studies fail to incorporate all potential factors that could affect the level of frost damage risk. A False Spring Index (FSI) signifies the likelihood of a damage to occur from a late spring freeze. Currently, FSI evaluates day of budburst, number of growing degree days, and day of last spring freeze through a simple equation as seen below (Marino *et al.*, 2011).

$$FSI = JulianDate(LastSpringFreeze) - JulianDate(Budburst)$$

If FSI is a positive number and greater than 7, then crown dieback is more likely to occur. In this study, we aim to integrate a more thorough model for predicting false spring risk, which would ideally incorporate life stage of the individual (Caffarra & Donnelly, 2011), location within a forest or canopy, winter chilling hours (Flynn, Wolkovich 2017?), proximity to water (Gu *et al.*, 2008), level of precipitation prior to the freezing event (Anderegg *et al.*, 2013), and freeze duration. Another highly crucial factor to consider is the rate of budburst and the length of time between budburst and leaf out at the species level. We will use the BBCH Scale Phase 09 to define budburst and Phase 15 to define leaf out (Meier, 2001).

## Climatic Implications: How climate change is affecting spring freezing events

Table 1: Number of False Springs along a Latitudinal Gradient in Western Europe

Station	Elevation	Latitude	Longitude	False Springs
Kempton, Germany	705m	47.724	10.336	8
Augsburg, Germany	461m	48.426	10.943	8
Bamberg, Germany	210m	49.875	10.921	7
Jena, Germany	155m	50.927	11.584	4
Hannover, Germany	55.0m	52.466	9.679	6
Bremen, Germany	4.00m	53.046	8.799	5
Hamburg, Germany	11.0m	53.635	9.99	4
Schleswig, Germany	43.0m	54.529	9.549	0
Flyvestation, Denmark	3.00m	57.093	9.849	0
Oslo, Norway	94.0m	59.943	10.721	0

Table 2: Number of False Springs along a Latitudinal Gradient in North America

Station	Elevation	Latitude	Longitude	False Springs
Anthony, Kansas	415m	37.155	-98.028	13
Hastings, Nebraska	587m	40.583	-98.350	7
West Point, Nebraska	399m	41.845	-96.714	5
Yankton, South Dakota	360m	42.883	-97.350	5
Brookings, South Dakota	497m	44.325	-96.769	0
Aberdeen, South Dakota	395m	45.443	-98.413	2
Grand Forks, North Dakota	253m	47.933	-97.083	0
Pembina, North Dakota	241m	48.971	-97.242	1

## Determining Spring Index

Before a suitable model for determining false spring risk can be established, an appropriate determination of spring onset is crucial. There are many methods that can be used to determine first day of spring and there also many definitions. In order to test the best method for calculating spring onset (or budburst), we gathered data using three different methodologies. The first method for collecting budburst was from observational data recorded for 33 tree species by Dr. John O’Keefe at Harvard Forest from 1990 to 2014 (O’Keefe, 2014). Dr. O’Keefe defines budburst as 50% leaf emergence. We subsetting this dataset down to include only the tree species that were most consistently observed, which ended up being eight species. The second data set was provided from PhenoCam data, which are field cameras, placed in Harvard Forest, take real-time images of plant growth and are programmed to record initial green up. The final set was collected

through the USA National Phenology Network (USA-NPN), using their Data Visualization tool to gather Extended Spring Index values (SI-x) by accessing the "Spring Indices, Historic Annual" gridded layer and looking specifically at "First Leaf - Spring Onset" (USA-NPN, 2016b). The SI-x value is the time of leaf out was monitored from historical dates of budburst using honeysuckle and lilac clones around the U.S. and combining that information with daily recordings from local weather stations (USA-NPN, 2016a; Ault *et al.*, 2015b,a; Schwartz *et al.*, 2013; Schwartz, 1997). Through assessing past years' weather and budburst, scientists are able to determine general weather trends that subsequently lead to leaf out. Based on these trends, SI-x values can be calculated from daily weather data (USA-NPN, 2016a).

The date of last spring freeze was gathered from the Fisher Meteorological Station which was downloaded from the Harvard Forest web page (data available online<sup>1</sup>). The Tmin values were used and the Last Spring Freeze was determined from the latest Julian date that the temperature reached -1.7°C or below.

PhenoCam data is not available for Harvard Forest until 2008 and observation data is only recorded through 2014, so this evaluation assesses FSI values from 2008 through 2014. The FSI values were calculated for each methodology using the formula based on the study performed by Marino *et al.* (2011). Table 2 shows that the Observed and PhenoCam FSI values are all negative from 2008 through 2014. The FSI values from the USA-NPN are, on average, much higher in comparison to the other two methods.

A graphical representation of the FSI values compared across the three methodologies can be seen in Figure 1. In 2008 and 2012, FSI is higher than the significant parameter given of 7 for the NPN data, indicating a possibly damaging false spring event. However, the PhenoCam data did not indicate a false spring in either of the two years and the FSI value for 2012 found through the observed data was not significant. The NPN data always had the highest FSI values.

---

<sup>1</sup><http://harvardforest.fas.harvard.edu/meteorological-hydrological-stations>




Figure 1: A scatterplot indicating FSI values from 2008 to 2014 for each methodology used in this study. PhenoCam FSI values are red, Observed FSI values are blue, and USA-NPN FSI values are green.

A Pearson Correlation was used to determine the strength of association between the three methods used in the study. As indicated in Table 1, the FSI values from the observed data and the SI-x NPN data are strongly correlated ( $r=0.93947095$ ), whereas the FSI values calculated using the PhenoCam data is not as strongly correlated to either the observed FSI values ( $r=0.4679934$ ) or the NPN FSI values ( $r=0.4242059$ ). Although, according to the pearson correlation, all results were similar.

Table 3: Pearson Correlation Coefficients shown comparing the strength of association between the FSI values calculated across all three methodologies.

	year	nbn	okeefe	phenocam
year	1.00	-0.03	-0.20	-0.37
nbn	-0.03	1.00	0.94	0.42
okeefe	-0.20	0.94	1.00	0.47
phenocam	-0.37	0.42	0.47	1.00

Our projections indicate that observational FSI values are highly comparable to the USA-NPN FSI values, rendering both justifiable methods for determining potential risk involved in late spring freezes. Even though budburst is defined differently between Dr. O’Keefe, USA-NPN, and the PhenoCam, the dates of budburst are similar. The spring onset dates gathered from the PhenoCam dataset are different from the other two methods, which is likely due to fact that the PhenoCam data is assessing budburst for the forest canopy. Through the use of USA-NPN data, researchers could gather dates of budburst across multiple locations at once in order to determine False Spring risk, making it a more effective method than observational data. Although, all three methods are viable. It largely depends on the functional group of interest for the study, definition of budburst, and resources available.

# Determining the Duration of Vegetative Risk (?)

## Tree Spotters Data




Figure 2: A timeline plot indicating the duration of vegetative risk for each species studied at the Arnold Arboretum in 2016.

## Dan's Data

It is possible, that with anthropogenic climate change progressing, leaf out timing may be delayed. As winter seasons begin to warm and chilling requirements are not met, more warming in the spring must first occur for budburst to begin (Polgar *et al.*, 2014; Fu *et al.*, 2012; Morin *et al.*, 2009; McCreary *et al.*, 1990)




Figure 3: A timeline plot indicating the duration of vegetative risk for each species from experimental chilling study.



Dan's Data




Figure 4: A timeline plot indicating the duration of vegetative risk for each species from collected from Harvard Forest.

## References

- Allstadt, A.J., Vavrus, S.J., Heglund, P.J., Pidgeon, A.M., Wayne, E. & Radeloff, V.C. (2015) Spring plant phenology and false springs in the conterminous U. S. during the 21st century. *Environmental Research Letters (submitted)* **10**, 104008.
- Anderegg, W.R.L., Plavcová, L., Anderegg, L.D.L., Hacke, U.G., Berry, J.A. & Field, C.B. (2013) Drought's legacy: Multiyear hydraulic deterioration underlies widespread aspen forest die-off and portends increased future risk. *Global Change Biology* **19**, 1188–1196.
- Augsburger, C.K. (2013) Reconstructing patterns of temperature, phenology, and frost damage over 124 years: Spring damage risk is increasing. *Ecology* **94**, 41–50.
- Ault, T.R., Henebry, G.M., de Beurs, K.M., Schwartz, M.D., Betancourt, J.L. & Moore, D. (2013) The False Spring of 2012, Earliest in North American Record. *Eos, Transactions American Geophysical Union* **94**, 181–182.
- Ault, T.R., Schwartz, M.D., Zurita-Milla, R., Weltzin, J.F. & Betancourt, J.L. (2015a) Trends and natural variability of spring onset in the coterminous united states as evaluated by a new gridded dataset of spring indices. *Journal of Climate* **28**, 8363–8378.
- Ault, T.R., Zurita-Milla, R. & Schwartz, M.D. (2015b) A Matlab{©} toolbox for calculating spring indices from daily meteorological data. *Computers {&} Geosciences* **83**, 46–53.
- Caffarra, A. & Donnelly, A. (2011) The ecological significance of phenology in four different tree species: Effects of light and temperature on bud burst. *International Journal of Biometeorology* **55**, 711–721.
- Fu, Y.H., Campioli, M., Van Oijen, M., Deckmyn, G. & Janssens, I.A. (2012) *Bayesian comparison of six different temperature-based budburst models for four temperate tree species*, vol. 230.
- Gu, L., Hanson, P.J., Post, W.M., Kaiser, D.P., Yang, B., Nemani, R., Pallardy, S.G. & Meyers, T. (2008) The 2007 Eastern US Spring Freeze: Increased Cold Damage in a Warming World. *BioScience* **58**, 253.
- Hufkens, K., Friedl, M.A., Keenan, T.F., Sonnentag, O., Bailey, A., O'Keefe, J. & Richardson, A.D. (2012) Ecological impacts of a widespread frost event following early spring leaf-out. *Global Change Biology* **18**, 2365–2377.
- Knudson, W. (2012) The economic impact of the spring's weather on the fruit and vegetable sectors. *The Strategic Marketing Institute Working Paper* .
- Kodra, E., Steinhäuser, K. & Ganguly, A.R. (2011) Persisting cold extremes under 21st-century warming scenarios. *Geophysical Research Letters* **38**, 1–5.

- Marino, G.P., Kaiser, D.P., Gu, L. & Ricciuto, D.M. (2011) Reconstruction of false spring occurrences over the southeastern United States, 1901–2007: an increasing risk of spring freeze damage? *Environmental Research Letters* **6**, 24015.
- McCreary, D.D., Lavender, D.P. & Hermann, R.K. (1990) Predicted global warming and Douglas-fir chilling requirements. *Annales des Sciences Forestieres* **47**, 325–330.
- Meier, U. (2001) Growth stages of mono-and dicotyledonous plants BBCH Monograph Edited by Uwe Meier Federal Biological Research Centre for Agriculture and Forestry. *Agriculture* **12**, 141—147 ST — Geochemical study of the organic mat.
- Morin, X., Lechowicz, M.J., Augspurger, C., O’keefe, J., Viner, D. & Chuine, I. (2009) Leaf phenology in 22 North American tree species during the 21st century. *Global Change Biology* **15**, 961–975.
- O’Keefe, J. (2014) Phenology of Woody Species at Harvard Forest since 1990. Tech. rep.
- Peterson, A.G. & Abatzoglou, J.T. (2014) Observed changes in false springs over the contiguous United States. *Geophysical Research Letters* **41**, 2156–2162.
- Polgar, C., Gallinat, A. & Primack, R.B. (2014) Drivers of leaf-out phenology and their implications for species invasions: Insights from Thoreau’s Concord. *New Phytologist* **202**, 106–115.
- Schwartz, M.D. (1997) Spring Index Models: An approach to connecting Satellite and surface phenology. *Phenology of Seasonal climates*, pp. 23–38.
- Schwartz, M.D., Ault, T.R. & Betancourt, J.L. (2013) Spring onset variations and trends in the continental United States: Past and regional assessment using temperature-based indices. *International Journal of Climatology* **33**, 2917–2922.
- USA-NPN (2016a) USA National Phenology Network.
- USA-NPN (2016b) USA National Phenology Network Data Visualizer Tool.
- Vavrus, S., Walsh, J.E., Chapman, W.L. & Portis, D. (2006) The behavior of extreme cold air outbreaks under greenhouse warming. *International Journal of Climatology* **26**, 1133–1147.
- Vitasse, Y., Lenz, A., Hoch, G. & Körner, C. (2014) Earlier leaf-out rather than difference in freezing resistance puts juvenile trees at greater risk of damage than adult trees. *Journal of Ecology* **102**, 981–988.
- Wolkovich, E.M., Cook, B.I., Allen, J.M., Crimmins, T.M., Betancourt, J.L., Travers, S.E., Pau, S., Regetz, J., Davies, T.J., Kraft, N.J.B., Ault, T.R., Bolmgren, K., Mazer, S.J., McCabe, G.J., McGill, B.J., Parmesan, C., Salamin, N., Schwartz, M.D. & Cleland, E.E. (2012) Warming experiments underpredict plant phenological responses to climate change. *Nature* **485**, 18–21.