

Topic 10: Eigenvalues and Eigenvectors

02-680: Essentials of Mathematics and Statistics

September 30, 2024

For a square matrix $A \in \mathbb{R}^{n \times n}$ for which $Ax = \lambda x$, we define $\lambda \in \mathbb{R}$ as an **eigenvalue** and $x \in \mathbb{R}^n \setminus \mathbf{0}$ as an **eigenvector**. Together we call these **eigenelements**.

The main idea is that the **transformation** of x by A results in a vector that points in the same direction of x , but scaled by λ .

Lets look at an example: let $A = \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix}$, $x = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$, and $\lambda = 5$.

$$\begin{aligned} Ax &\stackrel{?}{=} \lambda x \\ \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} &\stackrel{?}{=} 5 \begin{bmatrix} 2 \\ 1 \end{bmatrix} \\ \begin{bmatrix} (3 \cdot 2) + (4 \cdot 1) \\ (2 \cdot 2) + (1 \cdot 1) \end{bmatrix} &\stackrel{?}{=} \begin{bmatrix} (5 \cdot 2) \\ (5 \cdot 1) \end{bmatrix} \\ \begin{bmatrix} 10 \\ 5 \end{bmatrix} &= \begin{bmatrix} 10 \\ 5 \end{bmatrix} \end{aligned}$$

but given only A , how can we find the eigenelements?

1 Finding Eigenelements

Lets break down the original equation:

$$\begin{aligned} Ax &= \lambda x \\ Ax &= \lambda I_n x \\ Ax - \lambda I_n x &= \lambda I_n x - \lambda I_n x \\ (A - \lambda I_n)x &= 0 \end{aligned}$$

Since we know x cannot be $\mathbf{0}$ then $(A - \lambda I_n)$ but be 0 to satisfy the equation. This is actually a linear system on one variable! It turns out this only has non-trivial solutions when $(A - \lambda I_n)$ is singular (that is, it can't be invertible).

And remember we know that a matrix is singular when the determinant is 0:

$$|A - \lambda I_n| = 0.$$

We call this a *characteristic equation*.

Lets look at the example from above, we can rewrite it this way:

$$\begin{aligned} \begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} &= 0 \\ \begin{bmatrix} 3-\lambda & 4 \\ 2 & 1-\lambda \end{bmatrix} &= 0 \\ (3-\lambda)(1-\lambda) - (4)(2) &= 0 \\ (3-4\lambda+\lambda^2) - (8) &= 0 \\ \lambda^2 - 4\lambda - 5 &= 0 \\ (\lambda+1)(\lambda-5) &= 0 \end{aligned}$$

So we know that this matrix is singular when $\lambda = 5$ and -1 . These are the eigenvalues of A , to find the eigenvectors we plug this back into the original equation and solve for x .

$$\begin{aligned} \left(\begin{bmatrix} 3 & 4 \\ 2 & 1 \end{bmatrix} - 5 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} &= 0 \\ \begin{bmatrix} -2 & 4 \\ 2 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} &= 0 \end{aligned}$$

We can use whatever method we want to solve the system but we will find that for

$$\lambda = 5, x = \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

Similarly for $\lambda = -1$ we get the system:

$$\begin{bmatrix} 4 & 4 \\ 2 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0 \rightarrow x = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

Note in both of these cases there is a trivial solution of $x = \mathbf{0}$ but this is disallowed by definition.

Aside

A consequence of above is that the determinant is the sum of the eigenvalues:

$$|A| = \sum_{i=1}^n \lambda_i$$

A larger example. Find the eigenelements of $B = \begin{bmatrix} 1 & 1 & -1 \\ 2 & 3 & -4 \\ 4 & 1 & -4 \end{bmatrix}$.

We first write down the characteristic equation:

$$\begin{aligned} & \begin{vmatrix} 1-\lambda & 1 & -1 \\ 2 & 3-\lambda & -4 \\ 4 & 1 & -4-\lambda \end{vmatrix} = 0 \\ (1) & \begin{vmatrix} 2 & -4 \\ 4 & -4-\lambda \end{vmatrix} - (3-\lambda) \begin{vmatrix} 1-\lambda & -1 \\ 4 & -4-\lambda \end{vmatrix} + (1) \begin{vmatrix} 1-\lambda & -1 \\ 2 & -4 \end{vmatrix} = 0 \\ & \text{(note we chose column 2 because it had more 1's)} \\ & \dots \text{ details left to the reader } \dots \\ & \lambda^3 - 7\lambda + 6 = 0 \\ & (\lambda - 1)(\lambda - 2)(\lambda + 3) = 0 \end{aligned}$$

Therefore our eigenvalues are $\lambda = 1, 2$, and -3 .

If we solve the system:

$$\left(\begin{bmatrix} 1 & 1 & -1 \\ 2 & 3 & -4 \\ 4 & 1 & -4 \end{bmatrix} - \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

we find that $x_1 = x_2 = x_3$. So we can say $x = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$.

Finding the other eigenvectors is left as an exercise.

2 Triangular Matrices

A matrix $T \in \mathbb{R}^{n \times n}$ is triangular if all values on one side of the diagonal are 0:

$$\begin{bmatrix} T_{11} & T_{12} & \dots & T_{1(n-1)} & T_{1n} \\ 0 & T_{22} & \dots & T_{2(n-1)} & T_{2n} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & T_{(n-1)(n-1)} & T_{(n-1)n} \\ 0 & 0 & \dots & 0 & T_{nn} \end{bmatrix} \text{ or } \begin{bmatrix} T_{11} & 0 & \dots & 0 & 0 \\ T_{11} & T_{22} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ T_{(n-1)1} & T_{(n-1)2} & \dots & T_{(n-1)(n-1)} & 0 \\ T_{n1} & T_{n2} & \dots & T_{n(n-1)} & T_{nn} \end{bmatrix}.$$

It turns out triangular matrices have some special properties:

- $|T| = \sum_{i=1}^n T_{ii}$, and
- $\langle \lambda_1, \lambda_2, \dots, \lambda_n \rangle = \langle T_{11}, T_{22}, \dots, T_{nn} \rangle$.

Example. Find the eigenvalues of $D = \begin{bmatrix} 0 & 1 & 2 & 1 \\ 0 & 1 & 0 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}$.

$$\begin{aligned} \begin{vmatrix} -\lambda & 1 & 2 & 1 \\ 0 & 1-\lambda & 0 & 2 \\ 0 & 0 & 1-\lambda & 1 \\ 0 & 0 & 0 & 2-\lambda \end{vmatrix} &= 0 \\ -\lambda \begin{vmatrix} 1-\lambda & 0 & 2 \\ 0 & 1-\lambda & 1 \\ 0 & 0 & 2-\lambda \end{vmatrix} &= 0 \\ (-\lambda)(1-\lambda) \begin{vmatrix} 1-\lambda & 1 \\ 0 & 2-\lambda \end{vmatrix} &= 0 \\ (-\lambda)(1-\lambda)^2(2-\lambda) &= 0 \end{aligned}$$

We know λ s are 0, 1, 2 (since they are the distinct values).

We can solve for each λ :

$\lambda = 0$:

$$(D - 0I_4)x = 0 \rightarrow \begin{pmatrix} x_2 + 2x_3 + x_4 = 0 \\ x_2 + + 2x_4 = 0 \\ x_3 = 0 \\ 2x_4 = 0 \end{pmatrix} \rightarrow x = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Notice that x_1 is an arbitrary value.

$\lambda = 2$:

$$(D - 2I_4)x = 0 \rightarrow \begin{pmatrix} -2x_1 + x_2 + 2x_3 + x_4 = 0 \\ -x_2 + + 2x_4 = 0 \\ -x_3 = 0 \\ 0 = 0 \end{pmatrix} \rightarrow x = \begin{bmatrix} \frac{3}{2}x_4 \\ 2x_4 \\ 0 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3 \\ 4 \\ 0 \\ 2 \end{bmatrix}$$

Notice that x_4 is an arbitrary value, but chosen to make the vector integers.

$\lambda = 1$:

$$\begin{aligned} (D - I_4)x = 0 \rightarrow \begin{pmatrix} -x_1 + x_2 + 2x_3 + x_4 = 0 \\ 2x_4 = 0 \\ 0 = 0 \\ x_4 = 0 \end{pmatrix} \rightarrow \\ \begin{pmatrix} -x_1 + x_2 + 2x_3 = 0 \\ x_4 = 0 \end{pmatrix} \rightarrow x = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix} \text{ and } \begin{bmatrix} 2 \\ 0 \\ 1 \\ 0 \end{bmatrix} \end{aligned}$$

(Note these are *bases* of the space defined by the system above.)

3 Eigendecomposition and Diagonalization

Useful References

Isaak and Monougian, “Basic Concepts of Linear Algebra”. §7

Wilder, “10-606-f23:Lecture 6” GitHub repository, https://github.com/bwilder0/10606-f23/blob/main/files/notes_vectorspace.pdf

Kolter, “Linear Algebra Review and Reference”, <https://www.cs.cmu.edu/~zkolter/course/15-884/linalg-review.pdf> §3.12