## Lecture 1: Causal Identification

POL-GA 1251 Quantitative Political Analysis II Prof. Cyrus Samii NYU Politics

January 28, 2019

## Plan for the week

#### Mon:

- Discuss overview of class and syllabus.
- Explain what "causal identification" means.
- ▶ Introduce the "potential outcomes" framework
- Relate it to linear regression model.

#### Wed:

- Randomized experiments.
- Explain estimation concepts (estimand, estimators, bias, consistency, efficiency).
- Explain statistical inference concepts (sampling distribution, randomization distribution, CLT, confidence intervals, p-value).
- First homework distributed.

## Where this class fits in

## Model of quantitative research process:

- ▶ Theory motivates causal hypothesis or target of inference:
  - H: manipulating X results in (...) effect on Y.
- Hypothesis, statistical theory, and substantive theory motivate a research design:
  - Operationalize X and Y.
  - Define ways to get optimal variation in X and Y given constraints.
- ▶ Research design and statistical theory motivate analysis plan:
  - Optimal estimation strategy, given constraints.
  - Optimal testing strategy, given constraints.

## Where this class fits in

## Model of quantitative research process:

- ► Theory motivates causal hypothesis or target of inference:
  - H: manipulating X results in (...) effect on Y.
- Hypothesis, statistical theory, and substantive theory motivate a research design:
  - Operationalize X and Y.
  - Define ways to get optimal variation in X and Y given constraints.
- ► Research design and statistical theory motivate analysis plan:
  - Optimal estimation strategy, given constraints.
  - Optimal testing strategy, given constraints.

Identification refers generally to sufficiency for drawing a conclusion given *the type of data* that are available.

Identification refers generally to sufficiency for drawing a conclusion given *the type of data* that are available.

Distinguish from sampling uncertainty:

Identification refers generally to sufficiency for drawing a conclusion given *the type of data* that are available.

Distinguish from sampling uncertainty:

Manski (1995, 4): "Studies of identification seek to characterize the conclusions that could be drawn if one could...obtain an unlimited number of observations. Identification problems cannot be solved by gathering more of the same kind of data."

Identification refers generally to sufficiency for drawing a conclusion given *the type of data* that are available.

Distinguish from sampling uncertainty:

Manski (1995, 4): "Studies of identification seek to characterize the conclusions that could be drawn if one could...obtain an unlimited number of observations. Identification problems cannot be solved by gathering more of the same kind of data."

When we do not have identification, "the danger exists that the desired quantity will not be discernible unambiguously from the data – even when infinitely many samples are taken" (Pearl 2009, 77).

Identification refers generally to sufficiency for drawing a conclusion given *the type of data* that are available.

Distinguish from sampling uncertainty:

Manski (1995, 4): "Studies of identification seek to characterize the conclusions that could be drawn if one could...obtain an unlimited number of observations. Identification problems cannot be solved by gathering more of the same kind of data."

When we do not have identification, "the danger exists that the desired quantity will not be discernible unambiguously from the data – even when infinitely many samples are taken" (Pearl 2009, 77).

In this class we focus on causal identification. This is a specific application of the general idea of "identification," distinct from some other applications:

Alternative application of "identification" (I):

Suppose someone says...

...they prefer Harris over Sanders, and

...they prefer Warren over Harris.

Does this information (data) identify the person's preference ordering over these three candidates?

Alternative application of "identification" (II):

Suppose none of the coefficients below are equal to zero but the error terms (last ones) are iid mean zero draws. Which system of simultaneous equations identifies its coefficients?

$$\begin{aligned} x_t &= \alpha_1^a + \alpha_2^a y_t + \nu_t^a \\ y_t &= \beta_1^a + \beta_2^a x_t + \epsilon_t^a \\ \\ x_t &= \alpha_1^b + \alpha_2^b y_t + \alpha_3^b w_t + \alpha_4^b v_t + \nu_t^b \\ y_t &= \beta_1^b + \beta_2^b x_t + \beta_3^b w_t + \beta_4^b v_t + \epsilon_t^b \\ \\ x_t &= \alpha_1^c + \alpha_2^c y_t + \alpha_3^c w_t + \nu_t^c \\ y_t &= \beta_1^c + \beta_2^c x_t + \beta_4^c v_t + \epsilon_t^c \end{aligned}$$

Causal identification refers to sufficiency for drawing a conclusion about a *causal effect* given the type of data at hand.

Causal identification refers to sufficiency for drawing a conclusion about a *causal effect* given the type of data at hand.

This class focuses on strategies for non-parametric causal identification and, in turn, estimation of causal effects.

Causal identification refers to sufficiency for drawing a conclusion about a *causal effect* given the type of data at hand.

This class focuses on strategies for non-parametric causal identification and, in turn, estimation of causal effects.

Angrist and Krueger (1999):

The combination of a clearly labeled source of identifying variation in a causal variable and the use of a particular econometric[/statistical] technique to exploit this information is what we call an identification strategy.

#### The Road Not Taken

Two roads diverged in a yellow wood, And sorry I could not travel both And be one traveler, long I stood And looked down one as far as I could To where it bent in the undergrowth;

Then took the other, as just as fair, And having perhaps the better claim, Because it was grassy and wanted wear; Though as for that the passing there Had worn them really about the same,


And both that morning equally lay
In leaves no step had trodden black.
Oh, I kept the first for another day!
Yet knowing how way leads on to way,
I doubted if I should ever come back.

I shall be telling this with a sigh Somewhere ages and ages hence: Two roads diverged in a wood, and I— I took the one less traveled by, And that has made all the difference. Modern frameworks for causal analysis:

- Potential outcomes (Neyman, 1923; Rubin, 1974, 1978).
- ► Causal graphs (Pearl, 2009).

Both rely on "counterfactual" logic.

# Running Example: Effect of College on Earnings\*



A causal effect can be defined as a contrast between "potential outcomes."

	Pretreatment values			Which treatment		Po	osttr	eatment	: val	ues Y <sup>T</sup>		Missing data indicator  M  M  M  M  M  M  M  M  M  M  M  M  M									
	х,		x <sub>c</sub>		Υ <sub>1</sub>		Y <sub>d</sub>		$\mathbf{Y}_{1}^{\mathrm{T}}$		$Y_d^T$	н×		M <sup>X</sup> C	м <sub>1</sub>		$M_d^1$		$M_1^T$		$M_d^T$
Experimental units in population P										,											

Fig. 1. All values in a study of T treatments.




Fig. 1. All values in a study of T treatments.

▶ Population  $\mathcal{P}$  indexed by i = 1, ..., N.




Fig. 1. All values in a study of T treatments.

- ▶ Population  $\mathcal{P}$  indexed by i = 1, ..., N.
- ► Covariates, *x<sub>i</sub>*.




Fig. 1. All values in a study of T treatments.

- ▶ Population  $\mathcal{P}$  indexed by i = 1, ..., N.
- ► Covariates, *x<sub>i</sub>*.
- ► Treatment variable, W<sub>i</sub> with support W.




Fig. 1. All values in a study of T treatments.

- ▶ Population  $\mathcal{P}$  indexed by i = 1, ..., N.
- ► Covariates, *x<sub>i</sub>*.
- Treatment variable,
   W<sub>i</sub> with support W.
- ▶ Potential outcomes,  $y_{wi}$  given treatment  $w \in \mathcal{W}$ .




Fig. 1. All values in a study of T treatments.

- ▶ Population  $\mathcal{P}$  indexed by i = 1, ..., N.
- Covariates, x<sub>i</sub>.
  - Treatment variable,
     W<sub>i</sub> with support W.
- ▶ Potential outcomes,  $y_{wi}$  given treatment  $w \in \mathcal{W}$ .
- Missing data indicators,  $M_{i,j}^{(k)}$ .

Things to note:

#### Things to note:

► For now, we will assume outcomes for i depend only on treatment values for i and no one else — "stable unit treatment value assumption" (SUTVA), a.k.a. "no interference."

#### Things to note:

- ► For now, we will assume outcomes for i depend only on treatment values for i and no one else — "stable unit treatment value assumption" (SUTVA), a.k.a. "no interference."
- ▶ Unit level causal effects for members of  $\mathcal{P}$  are fixed a priori, and they compare  $y_{wi}$  to  $y_{\tilde{w}i}$  for  $w \neq \tilde{w}$  and  $i \in \mathcal{P}$ .

### Things to note:

- ► For now, we will assume outcomes for i depend only on treatment values for i and no one else — "stable unit treatment value assumption" (SUTVA), a.k.a. "no interference."
- ▶ Unit level causal effects for members of  $\mathcal{P}$  are fixed a priori, and they compare  $y_{wi}$  to  $y_{\tilde{w}i}$  for  $w \neq \tilde{w}$  and  $i \in \mathcal{P}$ .
- ▶ Population causal effects for compare aggregates of unit level causal effects for members of  $\mathcal{P}$ .
- ▶ Effects are defined in an "agnostic" or "non-parametric" way.
- Potential outcomes and covariates are fixed, treatments and response indicators stochastic.
- Effects are defined by letting only treatments vary, holding units fixed.
- Thus, causal effects are clearly defined for units that can conceivably receive different treatment values.
- ▶ A test for the above is "manipulation" (Holland, 1986).

# Potential outcomes, causal effects, and manipulability

Holland (1986): "For causal inference, it is critical that each unit be potentially exposable to any one of the causes."

Angrist and Krueger (1999): "The problem of ambiguous counterfactuals is typically resolved by focusing on hypothetic manipulations in the world as is."

# Potential outcomes, causal effects, and manipulability

Holland (1986): "For causal inference, it is critical that each unit be potentially exposable to any one of the causes."

Angrist and Krueger (1999): "The problem of ambiguous counterfactuals is typically resolved by focusing on hypothetic manipulations in the world as is."

Issues arise when trying to interpret things like race or gender. See VanderWeele and Robinson (2014) for a formal treatment of ways to interpret "race effects."

# Potential outcomes and fundamental problem of causal inference

Recall, a unit level causal effect compares  $y_{wi}$  to  $y_{\tilde{w}i}$  for  $w \neq \tilde{w}$ .

"Fundamental problem of causal inference" (Holland, 1986): For each i potential outcomes for all w exist, but we only observe the potential outcome for the treatment value that i receives.

- "Scientific solution": Use theory to determine when units are interchangeable.
- "Statistical solution": Study features of conditional distributions, such as averages.

Basic statistical setting (adjusting notation to conform to MHE):

Basic statistical setting (adjusting notation to conform to MHE):

▶ Consider a random draw, i, from P, countable but large.

Basic statistical setting (adjusting notation to conform to MHE):

- ▶ Consider a random draw, i, from P, countable but large.
- Each draw is characterized by
  - ightharpoonup a covariate vector,  $X_i$ ,
  - ▶ potential outcomes that under SUTVA are characterized as  $Y_{di}$  for all  $d \in \mathcal{D}$ , as well as
  - ▶ treatment assignments,  $D_i \in \mathcal{D}$ .

▶ Suppose  $\mathcal{D} = \{0,1\}$ . Then under SUTVA, potential outcomes for an arbitrary draw from  $\mathcal{P}$  are  $Y_{1i}$  and  $Y_{0i}$ .

- ▶ Suppose  $\mathcal{D} = \{0,1\}$ . Then under SUTVA, potential outcomes for an arbitrary draw from  $\mathcal{P}$  are  $Y_{1i}$  and  $Y_{0i}$ .
- ightharpoonup A unit level treatment effect for an arbitrary draw from  ${\cal P}$  is,

$$\rho_i = Y_{1i} - Y_{0i},$$

for which,

$$E[\rho_i] = E[Y_{1i} - Y_{0i}] = \rho,$$
 (1)

the average treatment effect (ATE).

- ▶ Suppose  $\mathcal{D} = \{0,1\}$ . Then under SUTVA, potential outcomes for an arbitrary draw from  $\mathcal{P}$  are  $Y_{1i}$  and  $Y_{0i}$ .
- lacktriangle A unit level treatment effect for an arbitrary draw from  ${\cal P}$  is,

$$\rho_i = Y_{1i} - Y_{0i},$$

for which,

$$E[\rho_i] = E[Y_{1i} - Y_{0i}] = \rho, \tag{1}$$

the average treatment effect (ATE).

▶ For an arbitrary draw from P, we observe,

$$Y_i = D_i Y_{1i} + (1 - D_i) Y_{i0}.$$

- ▶ Suppose  $\mathcal{D} = \{0,1\}$ . Then under SUTVA, potential outcomes for an arbitrary draw from  $\mathcal{P}$  are  $Y_{1i}$  and  $Y_{0i}$ .
- lacktriangle A unit level treatment effect for an arbitrary draw from  ${\cal P}$  is,

$$\rho_i = Y_{1i} - Y_{0i},$$

for which,

$$E[\rho_i] = E[Y_{1i} - Y_{0i}] = \rho, \tag{1}$$

the average treatment effect (ATE).

▶ For an arbitrary draw from  $\mathcal{P}$ , we observe,

$$Y_i = D_i Y_{1i} + (1 - D_i) Y_{i0}.$$

For our running example, we have  $D_i = 1$  if college,  $D_i = 0$  if not. Outcome of interest is income.  $\rho$  is the average income benefit of college.

Consider simple difference mean college grad incomes vs mean no college incomes:

$$E[Y_i|D_i=1] - E[Y_i|D_i=0] = E[Y_{1i}|D_i=1] - E[Y_{0i}|D_i=0]$$

Now, suppose in the population  $D_i = 1[f(X_i, U_i) > \eta_i]$ .

Consider simple difference mean college grad incomes vs mean no college incomes:

$$E[Y_i|D_i=1] - E[Y_i|D_i=0] = E[Y_{1i}|D_i=1] - E[Y_{0i}|D_i=0]$$

- Now, suppose in the population  $D_i = 1[f(X_i, U_i) > \eta_i]$ .
- ▶ Then what does  $E[Y_{1i}|D_i=1]$  equal?

Consider simple difference mean college grad incomes vs mean no college incomes:

$$E[Y_i|D_i=1] - E[Y_i|D_i=0] = E[Y_{1i}|D_i=1] - E[Y_{0i}|D_i=0]$$

- Now, suppose in the population  $D_i = 1[f(X_i, U_i) > \eta_i]$ .
- ▶ Then what does  $E[Y_{1i}|D_i=1]$  equal?
- ► For the full population,

$$E[Y_{1i}] = E[Y_{1i}|f(X_i, U_i) > \eta_i] \Pr[f(X_i, U_i) > \eta_i] + E[Y_{1i}|f(X_i, U_i) \le \eta_i] \Pr[f(X_i, U_i) \le \eta_i],$$

Consider simple difference mean college grad incomes vs mean no college incomes:

$$E[Y_i|D_i=1] - E[Y_i|D_i=0] = E[Y_{1i}|D_i=1] - E[Y_{0i}|D_i=0]$$

- Now, suppose in the population  $D_i = 1[f(X_i, U_i) > \eta_i]$ .
- ▶ Then what does  $E[Y_{1i}|D_i=1]$  equal?
- ► For the full population,

$$E[Y_{1i}] = E[Y_{1i}|f(X_i, U_i) > \eta_i] \Pr[f(X_i, U_i) > \eta_i] + E[Y_{1i}|f(X_i, U_i) \le \eta_i] \Pr[f(X_i, U_i) \le \eta_i],$$

We can also write

$$E[Y_{1i}|f(X_i, U_i) > \eta_i] = E[Y_{1i}|f(X_i, U_i) > \eta_i] \Pr[f(X_i, U_i) > \eta_i] + E[Y_{1i}|f(X_i, U_i) > \eta_i] \Pr[f(X_i, U_i) \leq \eta_i]$$

Consider simple difference mean college grad incomes vs mean no college incomes:

$$E[Y_i|D_i=1] - E[Y_i|D_i=0] = E[Y_{1i}|D_i=1] - E[Y_{0i}|D_i=0]$$

- Now, suppose in the population  $D_i = 1[f(X_i, U_i) > \eta_i]$ .
- ▶ Then what does  $E[Y_{1i}|D_i=1]$  equal?
- For the full population,

$$E[Y_{1i}] = E[Y_{1i}|f(X_i, U_i) > \eta_i] \Pr[f(X_i, U_i) > \eta_i] + E[Y_{1i}|f(X_i, U_i) \le \eta_i] \Pr[f(X_i, U_i) \le \eta_i],$$

We can also write

$$E[Y_{1i}|f(X_i, U_i) > \eta_i] = E[Y_{1i}|f(X_i, U_i) > \eta_i] \Pr[f(X_i, U_i) > \eta_i] + E[Y_{1i}|f(X_i, U_i) > \eta_i] \Pr[f(X_i, U_i) \leq \eta_i]$$

- ▶ Shows source of bias from selection into treatment.
- Similar could be shown for no-college outcomes.

Various decompositions are possible to illustrate sources of bias.

- Various decompositions are possible to illustrate sources of bias.
- ► E.g, consider a decomposition wrt ATT:

$$\begin{split} E[Y_i|D_i = 1] - E[Y_i|D_i = 0] &= E[Y_{1i}|D_i = 1] - E[Y_{0i}|D_i = 0] \\ &= \underbrace{E[Y_{1i}|D_i = 1] - E[Y_{0i}|D_i = 1]}_{\text{Average treatment effect on the treated (ATT)}} \\ &+ \underbrace{E[Y_{0i}|D_i = 1] - E[Y_{0i}|D_i = 0]}_{\text{Selection bias}}. \end{split}$$

- Various decompositions are possible to illustrate sources of bias.
- ► E.g, consider a decomposition wrt ATT:

$$E[Y_i|D_i = 1] - E[Y_i|D_i = 0] = E[Y_{1i}|D_i = 1] - E[Y_{0i}|D_i = 0]$$

$$= \underbrace{E[Y_{1i}|D_i = 1] - E[Y_{0i}|D_i = 1]}_{\text{Average treatment effect on the treated (ATT)}}$$

$$+ \underbrace{E[Y_{0i}|D_i = 1] - E[Y_{0i}|D_i = 0]}_{\text{Selection bias}}.$$

$$(2)$$

Could do similar wrt to ATC or effect heterogeneity (cf. CCI).

Now consider some identifying assumptions:

Now consider some identifying assumptions:

Identifying assumption 1 (random assignment):

$$D_i \perp (Y_{1i}, Y_{0i}) \text{ and } 0 < Pr[D_i = 1] < 1$$
 (3)

Now consider some identifying assumptions:

Identifying assumption 1 (random assignment):

$$D_i \perp (Y_{1i}, Y_{0i}) \text{ and } 0 < Pr[D_i = 1] < 1$$
 (3)

Under (3),

$$\underbrace{E[Y_{0i}|D_i=1] - E[Y_{0i}|D_i=0]}_{\text{Selection bias}} = 0.$$

(Why? E.g., if  $A \perp B$ , what does this imply about E[A|B]?)

Now consider some identifying assumptions:

Identifying assumption 1 (random assignment):

$$D_i \perp (Y_{1i}, Y_{0i}) \text{ and } 0 < Pr[D_i = 1] < 1$$
 (3)

Under (3),

$$\underbrace{E[Y_{0i}|D_i=1] - E[Y_{0i}|D_i=0]}_{\text{Selection bias}} = 0.$$

(Why? E.g., if  $A \perp B$ , what does this imply about E[A|B]?) As such,

$$\underbrace{E[Y_{1i}|D_i = 1] - E[Y_{0i}|D_i = 1]}_{ATT} = E[Y_{1i} - Y_{0i}],$$

so the simple difference, (2), equals  $\rho$ .

Identifying assumption 2 (conditionally independent/unconfounded/stronglignorable assignment):

$$D_i \perp (Y_{1i}, Y_{0i})|X_i$$
 and  $0 < Pr[D_i = 1|X_i = x] < 1$  for all  $x \in \mathcal{X}$  (4)

Identifying assumption 2 (conditionally independent/unconfounded/stronglignorable assignment):

$$D_i \perp (Y_{1i}, Y_{0i})|X_i \text{ and } 0 < Pr[D_i = 1|X_i = x] < 1 \text{ for all } x \in \mathcal{X}$$
(4)

The conditional average treatment effect (CATE) is given by,

$$\rho(x) = E[Y_{1i} - Y_{0i} | X_i = x].$$

Identifying assumption 2 (conditionally independent/unconfounded/stronglignorable assignment):

$$D_i \perp (Y_{1i}, Y_{0i})|X_i \text{ and } 0 < Pr[D_i = 1|X_i = x] < 1 \text{ for all } x \in \mathcal{X}$$
 (4)

The conditional average treatment effect (CATE) is given by,

$$\rho(x) = E[Y_{1i} - Y_{0i}|X_i = x].$$

By the same logic as before, assumption (4) implies,

$$E[Y_i|D_i = 1, X_i = x] - E[Y_i|D_i = 0, X_i = x] = \rho(x).$$

(Show this.)

Identifying assumption 2 (conditionally independent/unconfounded/stronglignorable assignment):

$$D_i \perp (Y_{1i}, Y_{0i})|X_i \text{ and } 0 < Pr[D_i = 1|X_i = x] < 1 \text{ for all } x \in \mathcal{X}$$
 (4)

The conditional average treatment effect (CATE) is given by,

$$\rho(x) = E[Y_{1i} - Y_{0i}|X_i = x].$$

By the same logic as before, assumption (4) implies,

$$E[Y_i|D_i = 1, X_i = x] - E[Y_i|D_i = 0, X_i = x] = \rho(x).$$

(Show this.)

Marginalization over X, the support of  $X_i$ , yields,

$$\int_{\mathcal{X}} \rho(x) dF(x) = \rho.$$

► Alternative framework is causal graph framework (Pearl 2009).

- ► Alternative framework is causal graph framework (Pearl 2009).
- Working with our example, we have in the population



- Alternative framework is causal graph framework (Pearl 2009).
- Working with our example, we have in the population



Our target of inference is based on an hypothetical intervention:

$$\mathsf{E}\left[Y|do(D_i=1)\right] - \mathsf{E}\left[Y|do(D_i=0)\right],$$

equivalent to the post-intervention graph



- Alternative framework is causal graph framework (Pearl 2009).
- Working with our example, we have in the population



Our target of inference is based on an hypothetical intervention:

$$\mathsf{E}\left[Y|do(D_i=1)\right] - \mathsf{E}\left[Y|do(D_i=0)\right],$$

equivalent to the post-intervention graph



▶ Difference between  $E[Y|do(D_i = 1)] - E[Y|do(D_i = 0)]$  and  $E[Y|D_i = 1] - E[Y|D_i = 0]$  is "backdoor paths" from D to Y.

▶ Random assignment ⇒ intervention graph is population graph:



► Random assignment ⇒ intervention graph *is* population graph:



► CIA implies no backdoor path through *U*:



▶ Random assignment ⇒ intervention graph is population graph:



CIA implies no backdoor path through U:



► Conditioning on *X* removes the other backdoor path:



▶ Marginalize over x to recover the intervention graph.

- ▶ These are examples of "closing" backdoor paths.
- ▶ Other operations, e.g., "opening" backdoor paths by conditioning on "colliders":





## Looking forward to the rest of the class

- With respect to our example and cases that resemble it:
  - ▶ These identifying assumptions rule out confounding by  $U_i$ .
  - ▶ If true, sufficient for identifying average treatment effect.
  - Plausibility is suspect, so need either other sources of identification or ways to check sensitivity to violations.

## Looking forward to the rest of the class

- With respect to our example and cases that resemble it:
  - ▶ These identifying assumptions rule out confounding by  $U_i$ .
  - ▶ If true, sufficient for identifying average treatment effect.
  - ▶ Plausibility is suspect, so need either other sources of identification or ways to check sensitivity to violations.
- More generally, this semester we will look at...
  - ...these and other identifying assumptions and then statistical methods that make use of them.
  - ...other target parameters (e.g., distributional effects), relax SUTVA, etc.
  - ...both potential outcomes and causal graphs representations
     each yields insights.

## Looking forward to the rest of the class

- With respect to our example and cases that resemble it:
  - ▶ These identifying assumptions rule out confounding by  $U_i$ .
  - ▶ If true, sufficient for identifying average treatment effect.
  - ▶ Plausibility is suspect, so need either other sources of identification or ways to check sensitivity to violations.
- More generally, this semester we will look at...
  - ...these and other identifying assumptions and then statistical methods that make use of them.
  - ...other target parameters (e.g., distributional effects), relax SUTVA, etc.
  - ...both potential outcomes and causal graphs representations
     each yields insights.
- Time for questions.