

Math 300: Advanced Boundary Value Problems

Week 8

1.1 Sturm-Liouville Theory

1. Theorem 4.5.

Given the regular Sturm-Liouville problem,

$$\begin{aligned}(p(x)\phi')' + [q(x) + \lambda\sigma(x)]\phi &= 0, \quad a < x < b, \\ \alpha_1\phi(a) + \beta_1\phi'(a) &= 0, \\ \alpha_2\phi(b) + \beta_2\phi'(b) &= 0,\end{aligned}$$

with eigenvalues λ_n and corresponding eigenfunctions ϕ_n .

(a) The regular Sturm-Liouville problem has an infinite spectrum

$$S = \{\lambda_1, \lambda_2, \dots, \lambda_n, \dots\}$$

and $\lim_{n \rightarrow \infty} \lambda_n = +\infty$.

(b) If $\alpha_1\beta_1 \leq 0$ and $\alpha_2\beta_2 \geq 0$, the spectrum is bounded below and the eigenvalues may be ordered as

$$\lambda_1 < \lambda_2 < \dots < \lambda_n < \dots$$

Moreover, if $q(x) \leq 0$ for $a \leq x \leq b$, then $\lambda_n \geq 0$ for all $n \geq 1$.

(c) If the eigenvalues are ordered as $\lambda_1 < \lambda_2 < \dots < \lambda_n < \dots$, the eigenfunction corresponding to λ_n has exactly $(n - 1)$ zeros in the interval $a < x < b$.

2. Theorem 4.6. (*Dirichlet's Theorem*)

If f is piecewise smooth on $[a, b]$, the **generalized Fourier series**,

$$f(x) \sim \sum_{n=1}^{\infty} c_n \phi_n(x), \quad \text{where,} \quad c_n = \frac{\langle f, \phi_n \rangle}{\|\phi_n\|^2} = \frac{1}{\|\phi_n\|^2} \int_a^b f(x) \phi_n(x) \sigma(x) dx,$$

for $n \geq 1$, converges pointwise to $[f(x^+) + f(x^-)]/2$ for each $x \in (a, b)$.

3. *Example 4.6.* Consider the regular Sturm-Liouville problem

$$\phi'' + \lambda\phi = 0, \quad 0 < x < 1,$$

$$\phi(0) = 0,$$

$$2\phi(1) - \phi'(1) = 0.$$

(continue Example 4.6)

4. *Example 4.7.* Consider the regular Sturm-Liouville problem

$$\begin{aligned}\phi'' + \lambda^2 \phi &= 0, & 0 < x < \pi, \\ \phi'(0) &= 0, \\ \phi(\pi) &= 0.\end{aligned}$$

- (a) Find the eigenvalues λ_n^2 and the corresponding eigenfunctions ϕ_n for this problem.
- (b) Show directly, by integration, that eigenfunctions corresponding to distinct eigenvalues are orthogonal.
- (c) Given the function $f(x) = \pi^2 - x^2/2, 0 < x < \pi$, find the eigenfunction expansion of f .
- (d) Show that

$$\frac{\pi^3}{32} = 1 - \frac{1}{3^3} + \frac{1}{5^3} - \frac{1}{7^3} + \frac{1}{9^3} - + \cdots.$$

(continue Example 4.7)

5. **Theorem 4.7.**

If (ϕ_n, λ_n) is an eigenpair for the regular Sturm-Liouville problem

$$\begin{aligned}(p(x)\phi')' + [q(x) + \lambda\sigma(x)]\phi &= 0, & a < x < b, \\ \alpha_1\phi(a) + \beta_1\phi'(a) &= 0, \\ \alpha_2\phi(b) + \beta_2\phi'(b) &= 0,\end{aligned}$$

then λ_n can be calculated from the **Rayleigh quotient**:

$$\lambda_n = \frac{-p(x)\phi_n(x)\phi'_n(x)\Big|_a^b + \int_a^b (p(x)\phi'_n(x)^2 - q(x)\phi_n(x)^2) dx}{\int_a^b \phi_n(x)^2 \sigma(x) dx}.$$

6. **Corollary 4.1.**

If

$$-p(x)\phi_n(x)\phi'_n(x)\Big|_a^b = -[p(b)\phi_n(b)\phi'_n(b) - p(a)\phi_n(a)\phi'_n(a)] \geq 0,$$

and $q(x) \leq 0$ for $a < x < b$, then $\lambda_n > 0$.

7. The **Rayleigh quotient** for **any** PWS function $u = u(x)$ on $[a, b]$ is given by

$$\mathcal{R}(u) = \frac{-p(x)u(x)u'(x)\Big|_a^b + \int_a^b (p(x)u'(x)^2 - q(x)u(x)^2) dx}{\int_a^b u(x)^2 \sigma(x) dx}.$$

8. **Theorem 4.8.**

Given the regular Sturm-Liouville problem

$$\begin{aligned}(p(x)\phi')' + [q(x) + \lambda\sigma(x)]\phi &= 0, & a < x < b, \\ \alpha_1\phi(a) + \beta_1\phi'(a) &= 0, \\ \alpha_2\phi(b) + \beta_2\phi'(b) &= 0,\end{aligned}$$

with spectrum

$$\lambda_1 < \lambda_2 < \cdots < \lambda_n < \cdots,$$

Then, the **leading eigenvalue** is

$$\lambda_1 = \min_u \mathcal{R}(u)$$

for all continuous functions u satisfying the boundary conditions

$$\alpha_1 u(a) + \beta_1 u'(a) = 0, \quad \alpha_2 u(b) + \beta_2 u'(b) = 0.$$

9. *Example 4.9.* Find good upper and lower bounds for the leading eigenvalue of the regular Sturm-Liouville problem

$$\begin{aligned}\phi'' - x\phi + \lambda\phi &= 0, & 0 < x < 1, \\ \phi'(0) &= 0, \\ 2\phi(1) + \phi'(1) &= 0.\end{aligned}$$

(continue Example 4.9)

10. *Example 4.10.* Find the generalized Fourier series solution to the homogeneous Neumann problem for the wave equation. Use the Rayleigh quotient to show that $\lambda_1 > 0$.

$$\alpha(x) \frac{\partial^2 u}{\partial t^2} = \frac{\partial}{\partial x} \left(\tau(x) \frac{\partial u}{\partial x} \right) - \beta(x)u, \quad 0 < x < l, \quad t > 0,$$

$$\frac{\partial u}{\partial x}(0, t) = 0, \quad t > 0,$$

$$\frac{\partial}{\partial x}(l, t) = 0, \quad t > 0,$$

$$u(x, 0) = f(x), \quad 0 < x < l,$$

$$\frac{\partial u}{\partial t}(x, 0) = g(x),$$

where $\alpha(x) > 0$, $\tau(x) > 0$, and $\beta(x) > 0$ for $0 < x < l$.

(continue Example 4.10)