







Simulation of Dynamical Scattering Effect in Electron Diffraction Patterns

Tarik Drevon, David Waterman, Eugene Krissinel




Discrepancy between theory and experiments

Macromolecules solved from experimental
Electron Diffraction (ED) data

Sample	Year	Resolution
Lysozyme (14.4 kDa) 	2013	2.9 Å
Sup35 prion protein core (905 Da) 	2016	1.0 Å
Au ₁₄₆ (p-MBA) ₅₇ nanoparticle (37.5 kDa) 	2017	0.85 Å


B. L. Nannenga and T. Gonen, “The cryo-EM method microcrystal electron diffraction (MicroED),” *Nat. Methods*, vol. 16, no. May, pp. 369–379, 2019

Multiple scattering should prevent use
of standard X-ray kinematic technique




G. Subramanian, S. Basu, H. Liu, J. Zuo, and J. C. H. Spence, “Solving protein nanocrystals by cryo-EM diffraction : Multiple scattering artifacts,” *Ultramicroscopy*, vol. 148, pp. 87–93, 2015

Electron diffraction and dynamical scattering



Atomic interaction cross section

$$P_{scattered} = \sigma \phi_{incident}$$



X-rays (Thomson scattering)


$$\sigma_{th} = \frac{8\pi}{3} r_e^2 = 66 fm^2$$

Electrons (Coulomb scattering)

$$\sigma_{th} \approx 1.87 \times 10^6 Z^{4/3} (c/v)^2 = 5 \times 10^7 fm^2$$


Mean free path

$$l_e = 1/\sigma_e \rho$$



Numerical simulation tools of ED patterns

Born approximation (used for X-rays)



Schrodinger's fast electron wave equation

$$\frac{\partial^2}{\partial z^2} \gg 2ik_0 \frac{\partial}{\partial z}$$

$$\left\{ \frac{\hbar^2}{2m_0} \nabla^2 + V(\mathbf{r}) \right\} \Psi(\mathbf{r}) = E\Psi(\mathbf{r})$$

$$\frac{\partial \Psi(x, y, z)}{\partial z} = \left\{ \frac{i\lambda}{4\pi} \nabla_{xy}^2 + i\sigma V(x, y, z) \right\} \Psi(x, y, z)$$

Method	Exact/ consistent	Speed	Memory (per atoms)	Periodic structure	Grid based	Package
Multislice (MS) (physical optics based approach)	no	fast	10 points	yes	yes	TEMSIM (pyMS) PRISM,...
Bloch wave (diagonalization in reciprocal space)	yes	slow	10 points	yes	no	EDM,PyQt,...
Near bragg (real space path differences)	no	??	1 point	no	no	NearBragg (James Holton)


Multisllice algorithm (MS)

$$\frac{\partial \Psi(x, y, z)}{\partial z} = \left\{ \frac{i\lambda}{4\pi} \nabla_{xy}^2 + i\sigma V(x, y, z) \right\} \Psi(x, y, z) \quad \longrightarrow \quad \Psi(z + \Delta z) = p(x, y, \Delta z) \boxed{*} \left(t(x, y, z) \Psi(z) \right)$$

FFT

Fresnel propagator

$$p(x, y, \Delta z) = \frac{1}{i\lambda \Delta z} e^{ik_0 \frac{x^2 + y^2}{2\Delta z}}$$



Transmission function



$$t(x, y, z) = e^{i\sigma \nu \Delta z}$$

$$\approx 1 + i\sigma \nu \Delta z$$

Weak potential


Projected potential

$$\nu_{\Delta z} = \int_z^{z+\Delta z} V(x, y, z') dz'$$




Multislice simple example


Transmission function
and projected potential




Real space wave
function as function of
propagation







Major beams
intensities in
reciprocal space




Multislice Silicon Diamond Structure



[110] vs [100]




Simulation of biotin





Structure $P2_12_12_1$



$$\begin{aligned}a &= 5.24\text{\AA}, \\b &= 10.35\text{\AA}, \\c &= 21.04\text{\AA}\end{aligned}$$





$[001]=0\text{deg}$



$[401]=45\text{deg}$



$[801]=64\text{deg}$



What's next?

- *Use CCP4 with the simulated data*
- *Include effects of defects and solvent scattering*
- *Develop nearBragg approach*
- *Make python interface available from the python index pypi*

More info <http://brno.rc-harwell.ac.uk:8000/> RCaH VPN required