

# P2Y12 Inhibitor Monotherapy Versus Conventional Dual Antiplatelet Therapy or Aspirin Monotherapy in Acute Coronary Syndrome: A Pooled Analysis of the SMART-DATE and SMART-CHOICE Trials



Pil Sang Song, MD<sup>a,1</sup>, Yong Hwan Park, MD<sup>b,1</sup>, Ju-Hyeon Oh, MD, PhD<sup>b</sup>, Young Bin Song, MD, PhD<sup>c</sup>, Seung-Hyuk Choi, MD, PhD<sup>c</sup>, Hyeon-Cheol Gwon, MD, PhD<sup>c</sup>, Deok-Kyu Cho, MD, PhD<sup>d</sup>, Seung-Woon Rha, MD, PhD<sup>e</sup>, Jang-Whan Bae, MD, PhD<sup>f</sup>, Jin-Ok Jeong, MD, PhD<sup>a,\*</sup>, and Joo-Yong Hahn, MD, PhD<sup>c,\*\*</sup>, for the SMART-DATE and the SMART-CHOICE investigators

**Controversy remains regarding the optimal antiplatelet regimen in patients with acute coronary syndrome (ACS). This study sought to investigate the efficacy and safety of P2Y12 inhibitor monotherapy compared with conventional dual antiplatelet therapy (DAPT) and aspirin monotherapy in patients with ACS undergoing percutaneous coronary intervention. Data on 4,453 patients were pooled from SMART-DATE and SMART-CHOICE randomized trials. Antiplatelet therapy regimens were categorized as P2Y12 inhibitor monotherapy (P2Y12 inhibitor monotherapy after 3-month DAPT), conventional DAPT (12-month or longer DAPT), and aspirin monotherapy (aspirin monotherapy after 6-month DAPT). The primary endpoint was major adverse cardiac and cerebrovascular events (MACCE, a composite of all-cause death, myocardial infarction, and stroke). Inverse-probability of treatment-weighted (IPTW) analysis was performed. At 1 year, patients in the P2Y12 inhibitor monotherapy had a comparable risk of MACCE compared with those in the conventional DAPT (IPTW-adjusted hazard ratio [HR], 0.655; 95% confidence interval [CI] 0.393 to 1.094;  $p = 0.106$ ), and tended to have a lower risk of MACCE than those in the aspirin monotherapy (IPTW-adjusted HR, 0.606; 95% CI, 0.347 to 1.058;  $p = 0.078$ ). The adjusted hazard for the Bleeding Academic Research Consortium (BARC) type 2 to 5 bleeding was significantly lower in P2Y12 inhibitor monotherapy than in conventional DAPT (IPTW-adjusted HR, 0.341; 95% CI, 0.190 to 0.614;  $p < 0.001$ ) and in aspirin monotherapy (IPTW-adjusted HR, 0.359; 95% CI, 0.182 to 0.708;  $p = 0.003$ ). In conclusion, among patients with ACS undergoing PCI, P2Y12 inhibitor monotherapy after 3-month DAPT reduced risk of bleeding compared with conventional DAPT and aspirin monotherapy after 6-month DAPT without increasing MACCE. © 2021 Elsevier Inc. All rights reserved. (Am J Cardiol 2021;150:47–54)**

<sup>a</sup>Division of Cardiology, Department of Internal Medicine, Chungnam National University Hospital, Chungnam National University College of Medicine, Daejeon, South Korea; <sup>b</sup>Division of Cardiology, Department of Internal Medicine, Samsung Changwon Hospital, Sungkyunkwan University School of Medicine, Changwon, South Korea; <sup>c</sup>Division of Cardiology, Department of Medicine, Heart Vascular Stroke Institute, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, South Korea; <sup>d</sup>Division of Cardiology, Department of Internal Medicine, Yonsei University College of Medicine and Cardiovascular Center, Yonjin Severance Hospital, Yonjin, South Korea; <sup>e</sup>Cardiovascular Center, Korea University Guro Hospital, Seoul, South Korea; and <sup>f</sup>Department of Internal Medicine, College of Medicine, Chungbuk National University Hospital, Cheongju, Republic of Korea. Manuscript received December 30, 2020; revised manuscript received and accepted March 22, 2021.

**Acknowledgment:** This study was supported by unrestricted grants from the Korean Society of Interventional Cardiology (grant number 2013-3), Abbott Vascular, Biotronik, and Boston Scientific.

**Clinical Trial Registration:** ClinicalTrials.gov Identifier - NCT01701453 and NCT02079194

<sup>1</sup>PSS and YHP contributed equally to this work. See page 53 for disclosure information.

\*Corresponding author: Tel: +82-42-280-8796; Fax: +82-42-280-8796.

\*\*Corresponding author: Tel: +82-2-3410-6653; Fax: +82-2-3410-6278

E-mail addresses: [jojeong@cnu.ac.kr](mailto:jojeong@cnu.ac.kr) (J.-O. Jeong), [jyhahn@skku.edu](mailto:jyhahn@skku.edu) (J.-Y. Hahn).

Dual antiplatelet therapy (DAPT) with aspirin and a P2Y12 inhibitor is the cornerstone of treatment after percutaneous coronary intervention (PCI). The current international guidelines recommend DAPT for at least 12 months after the implantation of drug-eluting stents in patients with acute coronary syndrome (ACS).<sup>1,2</sup> However, conventional DAPT for 12 months or longer after ACS increases the bleeding complications that counterbalance the benefit from preventing ischemic events.<sup>3,4</sup> Recently, an abbreviated DAPT strategy followed by a P2Y12 inhibitor monotherapy has emerged as a potential alternative to conventional DAPT regimens, aiming to reduce an excess of bleeding risk associated with the addition of aspirin while maintaining an anti-ischemic efficacy.<sup>5</sup> However, the ischemic and bleeding risks of an abbreviated DAPT strategy followed by a P2Y12 inhibitor monotherapy compared with conventional or 6-month DAPT in patients with ACS remain uncertain. To address these issues, we performed an analysis of the pooled data of the Smart Angioplasty Research Team-Safety of 6-Month Duration of Dual Antiplatelet Therapy After Percutaneous Coronary Intervention in Patients with Acute Coronary Syndrome (SMART-

DATE)<sup>6</sup> and Smart Angioplasty Research Team: Comparison Between P2Y12 Antagonist Monotherapy vs Dual Antiplatelet Therapy in Patients Undergoing Implantation of Coronary Drug-Eluting Stents (SMART-CHOICE)<sup>7</sup> trials to investigate the efficacy and safety of P2Y12 inhibitor monotherapy after 3 months of DAPT strategy after DES implantation in patients with ACS.

## Methods

The SMART-DATE was a multicenter, randomized, open-label trial to investigate whether a 6-month duration of DAPT would be non-inferior to the conventional 12-month or longer duration of DAPT in patients with ACS after coronary stenting. The SMART-CHOICE trial was a multicenter, randomized, open-label trial designed to test the non-inferiority of P2Y12 inhibitor monotherapy compared with aspirin plus a P2Y12 inhibitor after mandatory 3-month DAPT in patients undergoing PCI with current-generation drug-eluting stents. Among 5,705 patients included in the pooled data, 4,453 patients who underwent PCI for ACS were selected, comprising the population used in the present analysis. The antiplatelet therapy regimens were categorized as P2Y12 inhibitor monotherapy (P2Y12 inhibitor monotherapy after 3-month DAPT); conventional DAPT (12-month or longer DAPT); and aspirin monotherapy (aspirin monotherapy after 6-month DAPT); and we compared clinical endpoints among three antiplatelet therapy regimens. *e-Figure 1* in the *Supplementary Data* displays the network of antiplatelet regimens used in the present analysis. An independent institutional review board at each study center approved two clinical trials, and all patients provided written informed consent.

In the SMART-DATE trial, all patients received 300 mg of aspirin and a 300 or 600 mg clopidogrel loading dose orally at least 12 hours before PCI unless they had previously received these antiplatelet medications. After December 2014, prasugrel and ticagrelor became available in Korea. Since then, prasugrel or ticagrelor could be used instead of clopidogrel during the course of the SMART-DATE trial. After the procedure, aspirin (100 mg orally once daily) was used indefinitely and clopidogrel (75 mg orally once daily) was maintained according to the randomization scheme (6 months vs 12 months or longer). In the SMART-CHOICE trial, patients received DAPT with aspirin 100 mg once daily plus clopidogrel 75 mg once daily or prasugrel 10 mg once daily or ticagrelor 90 mg twice daily for 3 months in both groups after the procedure. The administration of aspirin was stopped at 3 months after the index procedure in the P2Y12 inhibitor monotherapy group but was continued indefinitely in the DAPT group. A P2Y12 inhibitor was prescribed continuously in both groups. It was recommended that all patients receive optimal pharmacologic therapy, including statins,  $\beta$ -blockers, or renin-angiotensin system blockade, if indicated, following clinical guidelines.<sup>1,2</sup> At follow-up, data from patients, including clinical status, all interventions, endpoint events, and adverse events, were recorded. In particular, information on the use of aspirin and a P2Y12 inhibitor was assessed at each follow-up. Patients who discontinued antiplatelet therapy as a result of clinically significant active bleeding or for

other procedures were monitored carefully for cardiac events, and, once they were stabilized, their allocated antiplatelet therapy was restarted as soon as possible. An independent clinical event adjudication committee, whose members were masked to the study group assignments, assessed all clinical endpoints.

In this pooled analysis, the primary endpoint is major adverse cardiac and cerebrovascular events (MACCE), defined as a composite of all-cause death, myocardial infarction (MI), and stroke at 12 months after the index procedure; secondary endpoints were individual components of the MACCE and the Bleeding Academic Research Consortium (BARC) type 2 to 5 bleeding at 12 months after the index procedure. In both trials, all clinical outcomes are commonly defined according to the Academic Research Consortium.<sup>8</sup> The BARC bleeding was defined as reported previously.<sup>9</sup>

The primary analysis was done according to an intention-to-treat principle. Because the follow-up duration of the SMART-CHOICE trial was 1 year, patients enrolled in the SMART-DATE trial were censored at 12 months in the current study. The cumulative rate of clinical events was evaluated by Kaplan-Meier analyses and the significance level was assessed with the Log-rank test. To compare the clinical endpoints, the hazard ratios (HRs) of three antiplatelet regimens were calculated by a Cox proportional hazards model. An extended description of the statistical analysis is presented in the *Methods of Supplementary Data*. Categorical variables are presented as numbers of events and percentages and were compared using the chi-square test or Fisher's exact test. Continuous variables are presented as mean  $\pm$  standard deviation or median (interquartile range, in the case of a non-normal distribution), and were compared using the *t* test or Mann Whitney test. All probability values were 2-sided and *p* values < 0.05 were considered statistically significant. For all analyses, we used SPSS software version 20.0 (SPSS Inc, Chicago, IL) and R software (version 3.3; "MatchIt" and "survival" packages; R Foundation for Statistical Computing, Vienna, Austria) for Windows for the statistical analyses.

## Results

Out of a total population of 4,453 ACS patients, 870 (19.5%) patients were treated P2Y12 inhibitor monotherapy, 2,226 (50.0%) with conventional DAPT, and 1,357 (30.5%) with aspirin monotherapy. Baseline characteristics according to the different treatment regimens are shown in [Table 1](#) and [Table 2](#). Compared to patients in the conventional DAPT group and the aspirin monotherapy group, patients in the P2Y12 inhibitor monotherapy group were older, had a higher prevalence of diabetes, hypertension, dyslipidemia, chronic renal failure, and previous coronary revascularization. Beta-blocker was less frequently prescribed at discharge in the P2Y12 inhibitor monotherapy group than in the conventional DAPT group and the aspirin monotherapy group. After adjustment with the use of inverse probability of treatment weighted (IPTW), clinical characteristics were well balanced except age, left ventricular ejection fraction, ST-segment elevation MI as a

Table 1  
Demographic and baseline characteristics of patients

Variable	Unadjusted data				Adjusted date with the use of IPTW			
	P2Y12 inhibitor monotherapy (n = 870)	Conventional DAPT (n = 2,226)	Aspirin monotherapy (n = 1,357)	p value	P2Y12 inhibitor monotherapy	Conventional DAPT	Aspirin monotherapy	p value
Age (year)	64.4 ± 11.3	63.0 ± 11.6	62.0 ± 11.5	<0.001	64.2 ± 10.0	63.1 ± 11.5	62.9 ± 11.4	0.028
Gender	629 (72.3%)	1677 (75.3%)	1016 (74.9%)	0.210	73.1%	74.5%	73.7%	0.816
Body Mass Index (Kg/m <sup>2</sup> )	24.4 ± 3.3	24.5 ± 3.1	24.3 ± 3.2	0.299	24.3 ± 2.9	24.4 ± 3.2	24.5 ± 3.2	0.331
STEMI as Presentation	164 (18.9%)	664 (29.8%)	509 (37.5%)	<0.001	22.3%	29.4%	30.3%	0.019
Diabetes	318 (36.6%)	698 (31.4%)	365 (26.9%)	<0.001	31.1%	31.4%	30.7%	0.926
Hypertension	530 (60.9%)	1175 (52.8%)	669 (49.3%)	<0.001	56.2%	53.7%	51.7%	0.227
Dyslipidemia	420 (48.3%)	755 (33.9%)	322 (23.7%)	<0.001	37.7%	34.1%	32.7%	0.126
Current smoker	306 (35.2%)	783 (35.2%)	506 (37.3%)	0.400	33.1%	35.7%	35.3%	0.549
History of CVA	52 (6.0%)	105 (4.7%)	52 (3.8%)	0.065	3.9%	4.9%	4.2%	0.472
History of CRF	25 (2.9%)	32 (1.4%)	13 (1.0%)	0.001	1.6%	1.7%	1.4%	0.770
History of previous MI	34 (3.9%)	60 (2.7%)	30 (2.2%)	0.056	2.4%	2.7%	2.2%	0.616
Previous revascularization	79 (9.1%)	147 (6.6%)	65 (4.8%)	<0.001	6.4%	6.5%	5.7%	0.628
LVEF (%)	58.2 ± 11.2	56.7 ± 10.5	55.7 ± 10.6	<0.001	57.9 ± 9.9	56.7 ± 10.7	56.5 ± 10.6	0.028

Plus–minus values are means ± SD.

CRF = chronic renal failure; CVA = cerebrovascular accident; DAPT = dual antiplatelet therapy; IPTW = inverse probability treatment weighting; LVEF = left ventricular ejection fraction; MI = myocardial infarction; STEMI = ST-segment elevation myocardial infarction.

presentation, trans-radial approach for PCI, and potent P2Y12 inhibitor prescription at discharge.

At 12 months, the primary endpoint of the MACCE occurred in 25 (2.9%) patients in the P2Y12 inhibitor monotherapy group, 72 (3.2%) patients in the conventional DAPT group, and 49 (3.6%) in the aspirin monotherapy group (Figure 1). After adjusting for potential confounders

using IPTW, there was no significant difference between the P2Y12 inhibitor monotherapy and conventional DAPT with respect to the MACCE (Table 3). The rate of MACCEs tended to be lower in the P2Y12 inhibitor monotherapy group than in the aspirin monotherapy group. Regarding MI, there was no significant difference between the P2Y12 inhibitor monotherapy and conventional DAPT. When

Table 2  
Angiographic and procedural characteristics of patients

Variable	Unadjusted data				Adjusted date with the use of IPTW			
	P2Y12 inhibitor monotherapy (n = 870)	Conventional DAPT (n = 2,226)	Aspirin monotherapy (n = 1,357)	p value	P2Y12 inhibitor monotherapy	Conventional DAPT	Aspirin monotherapy	p value
Multi-vessel disease	435 (50.0%)	1069 (48.0%)	591 (43.6%)	0.005	47.0%	47.4%	45.9%	0.788
Trans-radial access	569 (65.4%)	1203 (54.0%)	637 (46.9%)	<0.001	63.1%	54.0%	53.3%	0.001
Use of IVUS	217 (24.9%)	582 (26.1%)	311 (22.9%)	0.096	23.6%	24.9%	24.3%	0.856
PCI for calcified lesion	110 (12.6%)	291 (13.1%)	165 (12.2%)	0.727	11.6%	12.5%	12.8%	0.730
PCI for thrombotic lesion	100 (11.5%)	434 (19.5%)	325 (23.9%)	<0.001	13.5%	19.3%	19.7%	0.062
Bifurcation PCI	110 (12.6%)	227 (10.2%)	124 (9.1%)	0.028	9.3%	10.3%	10.5%	0.659
Treated vessel								
LM	11 (1.3%)	37 (1.7%)	29 (2.1%)	0.287	0.9%	1.6%	1.7%	0.235
LAD	507 (58.3%)	1370 (61.5%)	767 (56.5%)	0.009	62.5%	59.7%	59.5%	0.424
LCx	249 (28.6%)	561 (25.2%)	331 (24.4%)	0.068	24.6%	25.7%	24.5%	0.810
RCA	311 (35.7%)	817 (36.7%)	504 (37.1%)	0.799	33.0%	36.6%	38.1%	0.111
Multi-vessel PCI	199 (22.9%)	506 (22.7%)	263 (19.4%)	0.041	20.4%	21.7%	22.4%	0.657
Degradable polymer DES	282 (32.4%)	697 (31.3%)	391 (28.8%)	0.146	30.8%	31.4%	30.1%	0.821
Total Stent Length (mm)	37.6 ± 21.6	35.1 ± 21.4	32.3 ± 18.1	<0.001	34.8 ± 18.1	34.5 ± 20.4	34.5 ± 20.4	0.923
No. of stent implanted	1.46 ± 0.72	1.46 ± 0.77	1.39 ± 0.75	0.004	1.37 ± 0.63	1.44 ± 0.74	1.46 ± 0.81	0.051
Discharge medications								
Statin	840 (96.6%)	2064 (92.7%)	1212 (89.3%)	<0.001	94.3%	92.6%	92.1%	0.535
Beta-blocker	531 (61.0%)	1509 (67.8%)	961 (70.8%)	<0.001	66.1%	67.9%	68.1%	0.642
RAAS blocker	538 (61.8%)	1430 (64.2%)	929 (68.5%)	0.003	66.4%	64.5%	65.7%	0.730
Potent P2Y12 Inhibitor	305 (35.1%)	541 (24.3%)	275 (20.3%)	<0.001	29.8%	25.1%	23.7%	0.028

Plus–minus values are means ± SD

DAPT = dual antiplatelet therapy; DES = drug-eluting stent; IPTW = inverse probability treatment weighting; IVUS = intravascular ultrasound; LAD = left anterior descending coronary artery; LCx = left circumflex coronary artery; LM = left main coronary artery; PCI = percutaneous coronary intervention; RAAS = renin-angiotensin-aldosterone system; RCA = right coronary artery.




Figure 1. Time-to-Event Curves for Major Adverse Cardiovascular and Cerebrovascular Events (MACCE) (A), All-Cause Death (B), Myocardial Infarction (MI) (C), and Stroke (D). DAPT: dual antiplatelet therapy, FU: follow-up.

compared with aspirin monotherapy, however, P2Y12 inhibitor monotherapy was associated with a significantly lower risk of MI (Table 3). A total of 118 BARC type 2–5 bleeding events (2.6%) occurred at 12 months after the index PCI. Pairwise comparisons between groups are presented in Figure 2. The adjusted HR for bleeding endpoint was significantly lower for P2Y12 inhibitor monotherapy regimen compared with the conventional DAPT and aspirin monotherapy regimens (Table 3). In addition, patients treated with P2Y12 inhibitor monotherapy had a lower IPTW-adjusted all-cause death rate than those treated with conventional DAPT. The rate of all-cause death was also lower in the P2Y12 inhibitor monotherapy group than in the aspirin monotherapy group. Results from the per-protocol analysis were similar to those from the intention-to-treat analysis (e-Table 2 in the Supplementary Data). Details of results in patients with per-protocol treatment are presented in the Results in the Supplementary Data.

The treatment effects of the P2Y12 inhibitor monotherapy compared with conventional DAPT or aspirin monotherapy were consistent across various subgroups for the MI, including in subgroups analyzed according to the

presence versus absence of ST-segment elevation and type of P2Y12 inhibitors (e-Figure 2 in the Supplementary Data). In subgroup analyses for the risk of BARC type 2–5 bleeding, the results were also consistent across various subgroups (e-Figure 3 in the Supplementary Data).

## Discussion

Our analysis demonstrates that there were no significant differences between P2Y12 inhibitor monotherapy after 3-month DAPT and conventional DAPT for the primary endpoint of MACCE at 12 months after the index procedure. Compared with conventional DAPT, P2Y12 inhibitor monotherapy was associated with a significant reduction in BARC type 2 to 5 bleeding. Compared to aspirin monotherapy, P2Y12 inhibitor monotherapy significantly reduced the risk for MI. The results were consistent in per-protocol analyses and exploratory subgroup analysis.

Practice guidelines recommend DAPT for 12 months or longer in patients with ACS.<sup>1,2</sup> Recently, with the improved safety and efficacy of DES, a couple of studies challenged these guidelines and tested the safety of aspirin

Variable	Overall (n = 4,453)	P2Y12 inhibitor monotherapy (n = 870)	vsConventional DAPT (n = 2,226)	vsAspirin monotherapy (n = 1,357)	p-Value
<i>MACCE</i>					
Number of events	146 (3.3%)	25 (2.9%)	72 (3.2%)	49 (3.6%)	
Unadjusted HR (95% CI)		<i>Reference</i>	0.901 (0.571–1.420) p = 0.652	0.804 (0.496–1.300) p = 0.374	0.654
Adjusted HR (95% CI)		<i>Reference</i>	0.806 (0.510–1.274) p = 0.356	0.705 (0.434–1.147) p = 0.160	0.371
IPTW-Adjusted		<i>Reference</i>	0.655 (0.393–1.094) p = 0.106	0.606 (0.347–1.058) p = 0.078	0.186
<i>All-Cause Death</i>					
Number of events	85 (1.9%)	12 (1.4%)	46 (2.1%)	27 (2.0%)	
Unadjusted HR (95% CI)		<i>Reference</i>	0.676 (0.358–1.276) p = 0.227	0.702 (0.356–1.385) p = 0.307	0.474
Adjusted HR (95% CI)		<i>Reference</i>	0.566 (0.296–1.083) p = 0.086	0.541 (0.272–1.076) p = 0.080	0.178
IPTW-Adjusted		<i>Reference</i>	0.386 (0.197–0.756) p = 0.005	0.428 (0.202–0.908) p = 0.027	0.020
<i>Myocardial Infarction</i>					
Number of events	63 (1.4%)	8 (0.9%)	27 (1.2%)	28 (2.1%)	
Unadjusted HR (95% CI)		<i>Reference</i>	0.766 (0.348–1.686) p = 0.507	0.448 (0.204–1.018) p = 0.045	0.050
Adjusted HR (95% CI)		<i>Reference</i>	0.707 (0.320–1.558) p = 0.389	0.398 (0.180–0.876) p = 0.022	0.026
IPTW-Adjusted		<i>Reference</i>	0.655 (0.267–1.610) p = 0.357	0.353 (0.141–0.883) p = 0.026	0.033
<i>Stroke</i>					
Number of events	27 (0.6%)	6 (0.7%)	12 (0.5%)	9 (0.7%)	
Unadjusted HR (95% CI)		<i>Reference</i>	1.292 (0.484–3.436) p = 0.609	1.045 (0.372–2.933) p = 0.934	0.836
Adjusted HR (95% CI)		<i>Reference</i>	1.183 (0.443–3.155) p = 0.737	0.927 (0.329–2.611) p = 0.885	0.850
IPTW-Adjusted		<i>Reference</i>	1.211 (0.432–3.390) p = 0.717	0.664 (0.209–2.105) p = 0.487	0.476
<i>BARC type 2–5 Bleeding</i>					
Number of events	118 (2.6%)	15 (1.7%)	70 (3.1%)	33 (2.4%)	
Unadjusted HR (95% CI)		<i>Reference</i>	0.553 (0.317–0.965) p = 0.037	0.7158 (0.389–1.318) p = 0.283	0.086
Adjusted HR (95% CI)		<i>Reference</i>	0.499 (0.285–0.874) p = 0.015	0.605 (0.327–1.120) p = 0.109	0.049
IPTW-Adjusted		<i>Reference</i>	0.341 (0.190–0.614) p < 0.001	0.359 (0.182–0.708) p = 0.003	0.001

BARC = Bleeding Academic Research Consortium, CI = confidence interval, DAPT = dual antiplatelet therapy, HR = hazard ratio, IPTW = inverse probability treatment weighting, ITT = intention-to-treat, MACCE = major adverse cardiac and cerebrovascular events (a composite of all-cause death, myocardial infarction, and stroke).






Figure 2. Time-to-Event Curves for BARC type 2-5 Bleeding. BARC: Bleeding Academic Research Consortium, DAPT: dual antiplatelet therapy.

monotherapy after shortened DAPT compared with conventional DAPT in patients with ACS.<sup>10,11</sup> However, in the SMART-DATE trial, the risk of recurrent MI significantly increased with 6-month DAPT than with 12-month or longer DAPT in patients with ACS undergoing PCI in contemporary practice.<sup>6</sup> Therefore, the conventional DAPT regimen remains the standard of care in patients with ACS without excessive risk of bleeding. However, conventional DAPT increases bleeding and counteracts the benefit of reducing ischemic events. Therefore, the abbreviated duration of DAPT followed by P2Y12 inhibitor monotherapy has recently emerged as a promising alternative strategy.<sup>12</sup> Several randomized trials have demonstrated that P2Y12 inhibitor monotherapy is non-inferior to conventional DAPT regimen in the broad spectrum of patients.<sup>7,13,14</sup> However, data in patients with ACS are limited and, moreover, there have been no comparisons between 6-month DAPT followed by aspirin monotherapy and 3-month DAPT followed by P2Y12 inhibitor monotherapy. Therefore, in this pooled analysis of 2 randomized trials, we investigated the efficacy and safety of P2Y12 inhibitor

monotherapy compared with conventional DAPT or aspirin monotherapy in patients with ACS after coronary stenting.

In the present study, 3-month DAPT followed by P2Y12 inhibitor monotherapy was comparable to conventional DAPT with regards to the prevention of recurrent ischemic events, such as MI and stroke. Compared with 6-month DAPT followed by aspirin monotherapy, 3-month DAPT followed by P2Y12 inhibitor monotherapy was associated with a lower risk of MI. Although there are no randomized comparisons between P2Y12 inhibitor monotherapy and aspirin monotherapy after the current DES implantation, our results are in line with previous studies. In recent systematic review and meta-analysis, P2Y<sub>12</sub> inhibitor monotherapy is associated with a risk reduction for MI compared with aspirin monotherapy.<sup>15</sup> There is also supportive pharmacodynamic evidence. P2Y12 receptor activation is important to platelet thromboxane A<sub>2</sub> production, and the inhibition of P2Y12 can result in a substantial degree of thromboxane A<sub>2</sub> inhibition.<sup>16,17</sup>

Another important finding is that, compared with conventional DAPT, a significant reduction in BARC type 2 to

5 bleeding was prominent with P2Y12 inhibitor monotherapy. Even when compared to 6-month DAPT followed by aspirin monotherapy, the risk of bleeding was lower in the 3-month DAPT followed by P2Y12 inhibitor monotherapy. Three-month vs 6-month DAPT and the less adverse effect of clopidogrel on the gastrointestinal system when compared with aspirin might explain our results.<sup>18,19</sup>

Thereby, the rate of all-cause death was significantly lower in the P2Y12 inhibitor monotherapy group than in the conventional DAPT and aspirin monotherapy groups. In addition, the 3-month DAPT followed by P2Y12 inhibitor monotherapy tended to be better to aspirin monotherapy following 6-month DAPT with respect to the prevention of MACCE. Taken together, an abbreviated duration of DAPT followed by P2Y12 inhibitor monotherapy might be a promising strategy in patients with ACS after coronary stenting with a similar risk of ischemic events and reduction in bleeding complications compared with conventional DAPT, and with a lower risk of ischemic events compared with aspirin monotherapy.

This study has several limitations. First, a substantial proportion of patients in the P2Y12 inhibitor monotherapy group or in the aspirin monotherapy group received DAPT after 3 months or after 6 months, respectively. Therefore, the possibility of biases from low adherence in the P2Y12 inhibitor monotherapy group or in the aspirin monotherapy group could not be excluded. However, we also did a per-protocol analysis, which showed results consistent with the intention-to-treat analysis, suggesting that any potential biases caused by differential adherence and treatment crossover are likely to be small. Second, this was a post hoc exploratory analysis of 2 randomized trials; our report can identify correlations but does not prove causality. Third, the choice of the antiplatelet therapy regimen was randomly assigned in each trial, but significant differences were found in several variables between three antiplatelet regimen groups in the present analysis. Although we rigorously adjusted for these differences using multivariable Cox models and IPTW, unmeasured confounders may have affected our results. Fourth, event rates were low, so the present analysis may not be able to detect between-group differences in clinical endpoints. In terms of interaction tests, subgroup analyses reduced the sample size even more, therefore most are underpowered. Consequently, all the presented findings must be interpreted strictly as hypothesis-generating, which requires further prospective randomized validation.

In conclusion, 3-month DAPT followed by P2Y12 inhibitor monotherapy was comparable to conventional DAPT with regard to the prevention of ischemic events at 1 year after index PCI for ACS. Compared with conventional DAPT, P2Y12 inhibitor monotherapy was associated with a lower risk of BARC type 2 to 5 bleeding. In addition, P2Y12 inhibitor monotherapy after 3-month DAPT seems to more effectively prevent ischemic events than aspirin monotherapy after 6-month DAPT. These findings indicate that 3-month DAPT followed by P2Y12 inhibitor monotherapy could be a reasonable strategy in patients with ACS after implantation of current-generation DES. However, these findings should be interpreted as exploratory and

hypothesis-generating and should be tested in dedicated randomized trials.

## Author Contributions

Pil Sang Song: Conceptualization, Validation, Formal analysis, Investigation, Writing - Original Draft, Writing - Review & Editing, Visualization; Yong Hwan Park: Conceptualization, Validation, Formal analysis, Investigation, Writing - Original Draft, Writing - Review & Editing, Visualization; Ju-Hyeon Oh: Methodology, Software, Resources, Data Curation, Supervision, Project administration, Funding acquisition; Young Bin Song: Conceptualization, Investigation, Writing - Original Draft, Writing - Review & Editing, Supervision; Seung-Hyuk Choi: Conceptualization, Investigation, Writing - Original Draft, Writing - Review & Editing, Supervision; Hyeon-Cheol Gwon: Methodology, Software, Resources, Data Curation, Writing - Review & Editing, Supervision, Project administration, Funding acquisition; Deok-Kyu Cho: Methodology, Software, Resources, Data Curation, Supervision, Project administration, Funding acquisition; Seung-Woon Rha: Methodology, Software, Resources, Data Curation, Supervision, Project administration, Project administration, Funding acquisition; Jang-Whan Bae: Methodology, Software, Resources, Data Curation, Supervision, Project administration, Project administration, Funding acquisition; Jin-Ok Jeong: Conceptualization, Methodology, Software, Resources, Data Curation, Writing - Original Draft, Writing - Review & Editing, Supervision, Project administration, Project administration, Funding acquisition; Joo-Yong Hahn: Conceptualization, Methodology, Software, Resources, Data Curation, Writing - Original Draft, Writing - Review & Editing, Supervision, Project administration, Project administration, Funding acquisition.

## Disclosures

JY Hahn has received grants from Abbott Vascular, Biotronik, Boston Scientific, Daiichi Sankyo, and Medtronic; and speaker's fees from AstraZeneca, Daiichi Sankyo, and Sanofi-Aventis. HC Gwon has received research grants from Abbott Vascular, Boston Scientific, and Medtronic; and speaker's fees from Abbott Vascular, Boston Scientific, and Medtronic. All other authors declare that they have no conflicts of interest. The authors declare the financial interests/personal relationships which may be considered as potential competing interests.

## Supplementary materials

Supplementary material associated with this article can be found in the online version at <https://doi.org/10.1016/j.amjcard.2021.03.053>.

1. Levine GN, Bates ER, Bittl JA, Brindis RG, Fihn SD, Fleisher LA, Granger CB, Lange RA, Mack MJ, Mauri L, Mehran R, Mukherjee D, Newby LK, O'Gara PT, Sabatine MS, Smith PK, Smith SC Jr.. 2016 ACC/AHA guideline focused update on duration of dual antiplatelet therapy in patients with coronary artery disease. *Circulation* 2016;134:e123–e155.

2. Neumann FJ, Sousa-Uva M, Ahlsson A, Alfonso F, Banning AP, Benedetto U, Byrne RA, Collet JP, Falk V, Head SJ, Jüni P, Kastrati A, Koller A, Kristensen SD, Niebauer J, Richter DJ, Seferovic PM, Sibbing D, Stefanini GG, Windecker S, Yadav R, Zembala MO, ESC Scientific Document Group. 2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J* 2019;40:87–165.
3. Mauri L, Kereiakes DJ, Yeh RW, Driscoll-Shempp P, Cutlip DE, Steg PG, Normand SL, Braunwald E, Wiviott SD, Cohen DJ, Holmes DR Jr, Krucoff MW, Hermiller J, Dauerman HL, Simon DI, Kandzari DE, Garratt KN, Lee DP, Pow TK, Ver Lee P, Rinaldi MJ, Massaro JM. DAPT study investigators. Twelve or 30 months of dual antiplatelet therapy after drug-eluting stents. *N Engl J Med* 2014;371:2155–2166.
4. Gajananana D, Rogers T, Weintraub WS, Kolm P, Iantorno M, Khalid N, Chen Y, Shlofmitz E, Khan JM, Musallam A, Ben-Dor I, Satler LF, Zhang C, Torguson R, Waksman R. Ischemic versus bleeding outcomes after percutaneous coronary interventions in patients with high bleeding risk. *Am J Cardiol* 2020;125:1631–1637.
5. Malik AH, Yandrapalli S, Shetty SS, Aronow WS, Cooper HA, Panza JA. Meta-analysis of dual antiplatelet therapy versus monotherapy with P2Y12 inhibitors in patients after percutaneous coronary intervention. *Am J Cardiol* 2020;127:25–29.
6. Hahn JY, Song YB, Oh JH, Cho DK, Lee JB, Doh JH, Kim SH, Jeong JO, Bae JH, Kim BO, Cho JH, Suh IW, Kim DI, Park HK, Park JS, Choi WG, Lee WS, Kim J, Choi KH, Park TK, Lee JM, Yang JH, Choi JH, Choi SH, Gwon HC. SMART-DATE investigators. 6-month versus 12-month or longer dual antiplatelet therapy after percutaneous coronary intervention in patients with acute coronary syndrome (SMART-DATE): a randomised, open-label, non-inferiority trial. *Lancet* 2018;391:1274–1284.
7. Hahn JY, Song YB, Oh JH, Chun WJ, Park YH, Jang WJ, Im ES, Jeong JO, Cho BR, Oh SK, Yun KH, Cho DK, Lee JY, Koh YY, Bae JW, Choi JW, Lee WS, Yoon HJ, Lee SU, Cho JH, Choi WG, Rha SW, Lee JM, Park TK, Yang JH, Choi JH, Choi SH, Lee SH, Gwon HC, SMART-CHOICE Investigators. Effect of P2Y12 inhibitor monotherapy vs dual antiplatelet therapy on cardiovascular events in patients undergoing percutaneous coronary intervention: The SMART-CHOICE randomized clinical trial. *JAMA* 2019;321:2428–2437.
8. Cutlip DE, Windecker S, Mehran R, Boam A, Cohen DJ, van Es GA, Steg PG, Morel MA, Mauri L, Vranckx P, McFadden E, Lansky A, Hamon M, Krucoff MW, Serruys PW. Academic research consortium. Clinical end points in coronary stent trials: a case for standardized definitions. *Circulation* 2007;115:2344–2351.
9. Mehran R, Rao SV, Bhatt DL, Gibson CM, Caixeta A, Eikelboom J, Kaul S, Wiviott SD, Menon V, Nikolsky E, Serebruany V, Valgimigli M, Vranckx P, Taggart D, Sabik JF, Cutlip DE, Krucoff MW, Ohman EM, Steg PG, White H. Standardized bleeding definitions for cardiovascular clinical trials: a consensus report from the bleeding academic research consortium. *Circulation* 2011;123:2736–2747.
10. Kedhi E, Fabris E, van der Ent M, Buszman P, von Birgelen C, Roolvink V, Zurakowski A, Schotborgh CE, Hoorntje JCA, Eek CH, Cook S, Togni M, Meuwissen M, van Royen N, van Vliet R, Wedel H, Delawi R, Zijlstra F. Six months versus 12 months dual antiplatelet therapy after drug-eluting stent implantation in ST-elevation myocardial infarction (DAPT-STEMI): randomised, multicentre, non-inferiority trial. *BMJ* 2018;363:k3793.
11. Yerasi C, Case BC, Forrestal BJ, Torguson R, Weintraub WS, Garcia-Garcia HM, Waksman R. Optimizing monotherapy selection, aspirin versus P2Y12 inhibitors, following percutaneous coronary intervention. *Am J Cardiol* 2020;135:154–165.
12. Santos-Gallego CG, Badimon J. Overview of aspirin and platelet biology. *Am J Cardiol* 2021;144(Suppl 1):S2–S9.
13. Vranckx P, Valgimigli M, Jüni P, Hamm C, Steg PG, Heg D, van Es GA, McFadden EP, Onuma Y, van Meijeren C, Chichareon P, Benit E, Möllmann H, Janssens L, Ferrario M, Moschovitis A, Zurakowski A, Dominici M, Van Geuns RJ, Huber K, Slagboom T, Serruys PW, Windecker S, Investigators GLOBAL LEADERS. Ticagrelor plus aspirin for 1 month, followed by ticagrelor monotherapy for 23 months vs aspirin plus clopidogrel or ticagrelor for 12 months, followed by aspirin monotherapy for 12 months after implantation of a drug-eluting stent: a multicentre, open-label, randomised superiority trial. *Lancet* 2018;392:940–949.
14. Mehran R, Baber U, Sharma SK, Cohen DJ, Angiolillo DJ, Briguori C, Cha JY, Collier T, Dangas G, Dudek D, Dzavik V, Escaned J, Gil R, Gurbel P, Hamm CW, Henry T, Huber K, Kastrati A, Kaul U, Kornowski R, Krucoff M, Kunadian V, Marx SO, Mehta SR, Moliterno D, Ohman EM, Oldroyd K, Sardella G, Sartori S, Shlofmitz R, Steg PG, Weisz G, Witenbichler B, Han YL, Pocock S, Gibson CM. Ticagrelor with or without aspirin in high-risk patients after PCI. *N Engl J Med* 2019;381:2032–2042.
15. Chiarito M, Sanz-Sánchez J, Cannata F, Cao D, Sturla M, Panico C, Godino C, Regazzoli D, Reimers B, De Caterina R, Condorelli G, Ferrante G, Stefanini GG. Monotherapy with a P2Y12 inhibitor or aspirin for secondary prevention in patients with established atherosclerosis: a systematic review and meta-analysis. *Lancet* 2020;395:1487–1495.
16. Armstrong PC, Leadbeater PD, Chan MV, Kirkby NS, Jakubowski JA, Mitchell JA, Warner TD. In the presence of strong P2Y12 receptor blockade, aspirin provides little additional inhibition of platelet aggregation. *J Thromb Haemost* 2011;9:552–561.
17. Traby L, Kollars M, Kaider A, Eichinger S, Wolzt M, Kyrle PA. Effects of P2Y12 receptor inhibition with or without aspirin on hemostatic system activation: a randomized trial in healthy subjects. *J Thromb Haemost* 2016;14:273–281.
18. Sudlow CL, Mason G, Maurice JB, Wedderburn CJ, Hankey GJ. Thienopyridine derivatives versus aspirin for preventing stroke and other serious vascular events in high vascular risk patients. *Cochrane Database Syst Rev* 2009;7:CD001246.
19. Tillman H, Johnston SC, Farrant M, Barsan W, Elm JJ, Kim AS, Lindblad AS, Palesch YY, Easton JD. Risk for major hemorrhages in patients receiving clopidogrel and aspirin compared with aspirin alone after transient ischemic attack or minor ischemic stroke: a secondary analysis of the POINT randomized clinical trial. *JAMA Neurol* 2019;76:774–782.