

## Report from 2015 Brainhack Americas (LA)

# Calculating the Laterality Index Using FSL for Stroke Neuroimaging Data

Project URL: [http://github.com/npnl/LI\\_FSL](http://github.com/npnl/LI_FSL)

Kaori L. Ito<sup>??</sup> and Sook-Lei Liew

## 1 Introduction

The laterality index (LI) is one way to assess hemispheric dominance in a variety of tasks, such as language, cognitive functions, and changes in laterality in clinical populations, such as after stroke. In stroke neuroimaging, however, an optimal method of calculating the LI remains controversial, largely due to lesion variability in post-stroke brains.

Two main methods of calculating LI have evolved in neuroimaging literature [1]. The first, more traditional approach counts the number of active voxels in a given region of interest (ROI) for each hemisphere. This method has been criticized for its inability to account for differences in signal intensity. Hence, a second approach calculates laterality based on the percent signal change within a given region; however, this method also has problems, such as difficulty handling negative values.

A laterality toolbox that addresses some of these issues has been implemented in the statistical neuroimaging analysis package SPM, which provides users with options of using either method, along with more advanced statistical tests for robust LI calculations [2]. No such toolbox is yet available for FSL. Therefore, we developed a series of scripts to calculate LI in FSL using both voxel count and percent signal change methods. However, in the interest of space, here we present only results from the more robust method of the two (voxel count method).

## 2 Approach

We used fMRI data from two groups of stroke participants who either had right or left hemisphere lesions. Participants observed videos of right or left hand actions, and resulting statistical maps were calculated for each individual. The LI was then calculated per participant, based on the number of active voxels within a given anatomically-defined ROI (the inferior frontal gyrus, pars opercularis). Using the “cluster” tool in FSL, we set a threshold on the second-level whole-brain map. We set a range of z-values ( $z=1.0$ ,  $z=1.5$ ,  $z=2.3$ ) to test the effects of different thresholds. We then utilized “fslstats” to determine the total number of active voxels in both left and right hemisphere ROIs. Finally, we calculated LI based on the equation:

$$LI = (L - R) / (L + R)$$

where  $L$  represents the number of active voxels in the left-hemisphere ROI and  $R$  is the number of active voxels in the right-hemisphere ROI. This yields a value for LI such that  $-1 < LI < +1$ , where a positive value indicates left-hemisphere dominance and a negative value indicates right-hemisphere dominance.

## 3 Results/Discussion

We examined the variability in LI at different z-value thresholds to look at laterality differences in individuals with cortical versus subcortical stroke as well as the affected hemisphere (R vs. L). The LI values of four representative individuals (see Fig. 1) with the following types of stroke were as follows (see Table 1): subcortical left-hemisphere stroke (mean LI = -0.23; right lateralized), subcortical right-hemisphere stroke (mean LI = 0.79; left lateralized), cortical left-hemisphere stroke (mean LI = 0.96, left lateralized), and cortical right-hemisphere stroke (mean LI = 0.94, left lateralized). These LI results corresponded with our whole

<sup>??</sup> Correspondence: [kaoriito@usc.edu](mailto:kaoriito@usc.edu)


Neural Plasticity and Neurorehabilitation Laboratory, Chan Division of Occupational Science and Occupational Therapy, Division of Biokinesiology and Physical Therapy, Keck School of Medicine Department of Neurology, University of Southern California, Los Angeles, 1540 Alcazar Street, CHP 133 MC 9003, 90089, California, USA

Full list of author information is available at the end of the article

**Table 1** Laterality Index Using a Voxel-Count-based Method in FSL: A Comparison Across Different Stroke Lesion Profiles and Different Thresholds


Side of Stroke Lesion	Z-score	Subcortical Lesion			Cortical Lesion		
		LH	RH	LI	LH	RH	LI
Left	1	272	284	-0.022	382	22	0.891
	1.5	167	217	-0.130	101	0	1
	2.3	37	123	-0.538	1	0	1
Mean				-0.230			0.964
Right	1	335	68	0.662	509	49	0.824
	1.5	193	29	0.739	318	3	0.981
	2.3	76	1	0.974	216	0	1
Mean				0.792			0.935

brain observations (not included here), in which we examined the laterality of the action observation network in individuals with left versus right hemisphere stroke (Ito, OHBM 2016). Briefly, we show that most individuals show a left-lateralized pattern of activity during action observation, regardless of the lesion location. That is, Both individuals with left-hemisphere stroke and individuals with right-hemisphere stroke tend to activate the left, dominant hemisphere, suggesting a role of hemispheric dominance during recovery after stroke.



**Figure 1** A Comparison Across Different Stroke Lesion Profiles at Maximum Lesion. MRI scans of individuals who sustained A. subcortical left-hemisphere stroke, B. cortical left-hemisphere stroke, C. subcortical right-hemisphere stroke, D. cortical right-hemisphere stroke.

Importantly, we notice that the voxel count method is highly dependent on the threshold value: as the threshold increases in stringency, the value of the LI increases. With individuals after stroke, higher thresholds may yield 0 active voxels, leading to a potentially skewed LI (LI=1).



**Figure 2** Overlap Between Activation in ROI and Lesion. Activity in the inferior frontal gyrus, pars opercularis at  $z=1.5$  for individuals who sustained A. subcortical left-hemisphere stroke, B. cortical left-hemisphere stroke, C. subcortical right-hemisphere stroke, D. cortical right-hemisphere stroke. Images were registered to standard space.

4 Conclusions

We suggest that stroke neuroimaging might benefit from calculating an average LI across different thresholds (including more lenient thresholds such as  $z=1.0$ ), in order to provide a more robust outcome that takes into account threshold dependency. This is especially true for individuals with cortical strokes, where the ROI may overlap with the lesion and yield 0 active voxels. This issue of thresholding, specifically for stroke research, is an interesting question that remains to be addressed further. Future work may also utilize threshold-weighted overlap maps to visualize subject variability when using different thresholds to calculate a laterality index [3]. Our scripts for these calculations

may be found online at <http://chan.usc.edu/npnl/resources>.

#### Availability of Supporting Data

More information about this project can be found at:

[http://github.com/npnl/LI\\_FSL](http://github.com/npnl/LI_FSL). Further data and files supporting this project are hosted in the *GigaScience* repository REFXXX.

#### Competing interests

None

#### Author's contributions

KLI and SLL wrote the software, performed tests, and wrote the report.

#### Acknowledgements

The authors would like to thank the organizers and attendees of Brainhack LA and the developers of FSL. This project was funded in part by a Educational Research Grant from Amazon Web Services.

#### References

1. Jansen, A., Menke, R., Sommer, J., Förster, A.F., Bruchmann, S., Hempleman, J., Weber, B., Knecht, S.: The assessment of hemispheric lateralization in functional mri—robustness and reproducibility. *NeuroImage* **33**(1), 204–217 (2006). doi:[10.1016/j.neuroimage.2006.06.019](https://doi.org/10.1016/j.neuroimage.2006.06.019)
2. Wilke, M., Lidzba, K.: Li-tool: A new toolbox to assess lateralization in functional mr-data. *Journal of Neuroscience Methods* **163**(1), 128–136 (2007). doi:[10.1016/j.jneumeth.2007.01.026](https://doi.org/10.1016/j.jneumeth.2007.01.026)
3. Seghier, M.L., Price, C.J.: Visualising inter-subject variability in fmri using threshold-weighted overlap maps. *Scientific Reports* **6**(20170) (2016). doi:[10.1038/srep20170](https://doi.org/10.1038/srep20170)