

Exercise2_cc4515_cy2550_ml4404

February 11, 2020

```
[1]: import numpy as np
from scipy.misc import derivative
from scipy.stats import norm
from sympy import diff, symbols
```

Exercise 2

Chutian Chen cc4515; Congcheng Yan cy2550; Mingrui Liu ml4404

(1)

$$E(\bar{X}^2) = E\left(\frac{\sum_{i=1}^n X_i^2 + 2 \sum_{i=1}^{n-1} \sum_{j=i+1}^n X_i X_j}{n^2}\right)$$

$$= \frac{1}{n} EX^2 + \left(1 - \frac{1}{n}\right) (EX)^2$$

$$EX^2 = \text{Var}X + (EX)^2 = \lambda + \lambda^2$$

$$EX = \lambda$$

$$\text{So } E(\bar{X}^2) = \lambda^2 + \frac{\lambda}{n}$$

(2)

$$Es^2 = \frac{1}{n-1} E\left(\sum_{i=1}^n X_i^2 - n\bar{X}^2\right)$$

$$= \frac{1}{n-1} (n\lambda + n\lambda^2 - n\lambda^2 - \lambda) = \lambda$$

(3)

$$EY_i = EX_i^2 - 2\lambda EX_i + \lambda^2 - EX_i$$

$$= \lambda^2 + \lambda - 2\lambda^2 + \lambda^2 - \lambda = 0$$

$$\text{Var}Y_i = EY_i^2 - (EY_i)^2 = EY_i^2 = E(X_i^2 - (2\lambda + 1)X_i + \lambda^2)^2$$

Using moment generating function, we can get

$$EX = \lambda$$

$$EX^2 = \lambda^2 + \lambda$$

$$EX^3 = \lambda^3 + 3\lambda^2 + \lambda$$

$$EX^4 = \lambda^4 + 6\lambda^3 + 7\lambda^2 + \lambda$$

$$\text{So } \text{Var}Y_i = 2\lambda^2$$

(4)

$$\begin{aligned}
s^2 - \bar{X} &= \frac{1}{n-1} (\sum_{i=1}^n X_i^2 - n\bar{X}^2 - (n-1)\bar{X}) \\
Y_i &= (X_i - \bar{X})^2 + (\bar{X} - \lambda)^2 + 2(X_i - \bar{X})(\bar{X} - \lambda) - X_i \\
\sum_{i=1}^n Y_i &= \sum_{i=1}^n (X_i - \bar{X})^2 + n(\bar{X} - \lambda)^2 - \sum_{i=1}^n X_i \\
&= (n-1)s^2 + n(\bar{X} - \lambda)^2 - n\bar{X} \\
\text{So } s^2 - \bar{X} &= \frac{\sum_{i=1}^n Y_i - n(\bar{X} - \lambda)^2 + \bar{X}}{n-1}
\end{aligned}$$

(5)

By CLT, $\sqrt{n}\bar{Y} \Rightarrow_D N(0, 2\lambda^2)$

$$\frac{n}{n-1} \Rightarrow_P 1$$

By LLN, $(\bar{X} - \lambda) \Rightarrow_P 0$

By CLT, $\sqrt{n}(\bar{X} - \lambda) \Rightarrow_D N(0, \lambda)$

Then Slutsky's Theorem tells that

$$\sqrt{n}(\bar{X} - \lambda)(\bar{X} - \lambda) = \sqrt{n}(\bar{X} - \lambda)^2 \Rightarrow_D 0 * N(0, \lambda) = 0$$

$$\frac{\bar{X}}{\sqrt{n}} \Rightarrow_P 0$$

$$\text{So } (\sqrt{n}\bar{Y} - \sqrt{n}(\bar{X} - \lambda)^2 + \frac{\bar{X}}{\sqrt{n}}) \Rightarrow_D N(0, 2\lambda^2)$$

$$\text{Because } \frac{n}{n-1} \Rightarrow_P 1$$

$$\frac{n}{n-1}(\sqrt{n}\bar{Y} - \sqrt{n}(\bar{X} - \lambda)^2 + \frac{\bar{X}}{\sqrt{n}}) \Rightarrow_D N(0, 2\lambda^2)$$

$$\text{So } \sqrt{n}(s^2 - \bar{X}) \Rightarrow_D N(0, 2\lambda^2)$$

$$\bar{X} \Rightarrow_P \lambda$$

$$\text{So } \sqrt{\frac{n}{2} \frac{(s^2 - \bar{X})}{\bar{X}}} \Rightarrow_D N(0, 1)$$

(6)

$$\text{When } E(\bar{X}) = E(s^2), P(|\sqrt{\frac{n}{2} \frac{(s^2 - \bar{X})}{\bar{X}}} - E(s^2 - \bar{X})| \leq Z_{1-\alpha/2}) = \alpha$$

$$\text{So if } |\sqrt{\frac{n}{2} \frac{(s^2 - \bar{X})}{\bar{X}}}| \geq Z_{1-\alpha/2}, \text{ we can reject } H_0.$$

(7)

```
[9]: model_1 = np.random.poisson(5, 500*50)
     m1, v1 = np.mean(model_1), np.var(model_1)
```

```
[10]: def model2_gen(n):
      a = []
      for i in range(n):
          a = a + list(np.random.poisson(np.random.gamma(2.5, 2, 1), 50))
      return np.array(a)
```

```
[11]: model_2 = model2_gen(500)
     m2, v2 = np.mean(model_2), np.var(model_2)
```

```
[12]: print("Model A: ", np.sqrt(500/2)*(v1-m1)/m1)
      print("Model B: ", np.sqrt(500/2)*(v2-m2)/m2)
```

Model A: 0.023104027613830565

Model B: 31.934773221864305

```
[57]: norm.ppf(0.975)
```

```
[57]: 1.959963984540054
```

As we can see, model A $0.023 < 1.96$, model B $31.93 > 1.96$.

So for model A we can accept H_0 at significant level 0.05.

For model B we should reject H_0 at significant level 0.05.

(8)

```
[58]: f = [1,4,15,31,39,55,54,49,47,31,16,9,8,4,3]
      data = []
      for i,fre in enumerate(f):
          for j in range(fre):
              data.append(i)
      data = np.array(data)
```

```
[59]: m3,v3 = np.mean(data), np.var(data)
      np.sqrt(len(data)/2)*(v3-m3)/m3
```

```
[59]: 0.9390395410831514
```

$0.93 < 1.96$, so we can accept H_0 at significant level 0.05.

The test doesn't detect any overdispersions.

```
[60]: #e = np.exp(1)
      #t = symbols('x', real=True)
      #l = symbols('l', real = True)
      #diff(e**(l*(e**t-1)), t, 4).subs(t,0)
```