



Universidad de los Andes

PHYSICS DEPARTMENT

MEASUREMENT AND CHARACTERIZATION OF
GRANULATION PATTERN IN THE IAG SOLAR
FLUX SPECTRUM

BSc Physics Final Project

Author:

Claudia Alejandra Cuellar Nieto

Advisor:

Benjamin Oostra Vannoppen

Nov 2025

Abstract

Hello, this is my work :D

Acknowledgements

Thanks :b

Contents

List of Figures	3
1 Introduction: The Sun's granulation pattern	5
1.1 Convective motion	6
1.2 Solar granulation pattern	6
1.2.1 The three signatures of convection	7
1.3 IAG Solar Flux Atlas	7
1.4 IAG Spatially Resolved Quiet Sun Atlas	8
1.5 Motivation	8
2 Literature Review: Convective movement in the Sun	10
2.1 The solar interior and the solar outer atmosphere	10
2.2 The solar convection Zone	12
2.2.1 The convection movement in the sun	13
2.2.2 Dynamics of solar convection	13
2.2.3 The Schwarzschild condition	14
2.3 The Solar Photosphere	15
2.3.1 Static photosphere: Limb darkening phenomenon	15
2.3.2 Dynamic photosphere: The C-curved profile bisector.	17
2.4 The three signature of convection	18
2.4.1 Line broadening	18
2.4.2 Line profile bisector asymmetry	19

2.4.3	Line depth-dependent wavelength shifts	20
2.4.4	Solar granulation pattern	20
2.4.5	Chromodependence characterization	21
3	Methodology: The blend-free list of Fe I lines	23
3.1	Methodology	23
3.2	Blend-free Fe I line list	24
3.2.1	Selection method for Fe I lines	24
4	Results and discussion: Granulation pattern and Characterization of chromodependence	26
4.1	The third signature: Chromodependence on the granulation pattern . . .	26
4.2	The first and second signature: Detailed view of line broadening and asymmetry	30
4.3	Higher quality graphs	33
5	Conclusions	34
A	Z-score Standardization	35
B	The third derivate relation	36
C	Visualizer for outliers	37

List of Figures

2.1	The interior structure of the Sun. The convection zone is the responsible by the general movement that characterize the third signature.	11
2.2	The first clear photography of the photosphere where is visble the granulation pattern. Image taken from [1]	12
2.3	A view of granulation on the Sun's surface. The central regions exhibit blueshifts while the edges display redshifts. Image taken from [2]	15
2.4	Squared profile for the disk intensity at increasing wavelengths. Image taken from [3]	16
2.5	The image shows frame by frame the motion of few granules. Image taken from [3]	16
2.6	The C-curved profile bisector. In the infrared shown to be less pronounced than violet range.	17
2.7	In the figure is comparised the strongest and the weaker line core shapes.	18
2.8	Asymmetries on an average absorption line. Can be observed the differences in intensity redshift profile. Image taken from [4]	19
2.9	Granulation pattern for the IAG spatially resolved quiet sun atlas, is shown to have a strong chromodependece in the weaker lines. Image taken from [5].	20
2.10	Contributions on angular momentum. Image taken from [3].	22
4.1	Granulation pattern for the Solar Flux Atlas.	27
4.2	Granulation pattern for the Solar Flux Atlas with color curves.	27

4.3	Granulation pattern for the Spatially Resolved Quiet Sun Atlas at $\mu = 1$.	28
4.4	Comparision between atlases with velocity bins for the relation between wavelength and line depth.	29
4.5	Comparision between atlases with velocity bins for the relation between wavelength and line depth. The first order fit is showed for each velocity bin.	30
4.6	C bisector graph for slopes in ehich is clear the convection movement and how its affected in the profile.	31
4.7	Sharpness of the core bisector in the solar flux atlas separated by range. .	32
4.8	Sharpness of the core bisector in the solar flux atlas. Is clear that the infrared follow two different behavior due teluric lines.	33
4.9	We realized the same graphic for the comparision with the Ellwarth article to show the less scattered points.	33

Chapter 1

Introduction: The Sun’s granulation pattern

For decades, the solar spectrum has served as the fundamental reference point for spectroscopic analysis and characterization. As our nearest star, the Sun has enabled comprehensive studies of stellar composition. However, advances in optical instrumentation have recently revealed previously undetected spectral details, providing new insights into even the most basic solar properties from dynamics to surface geometry. One of this advances is the granulation pattern due to convection motion, which is revealed by three fundamental signatures of its hydrodynamics: Line broadening, line profile bisector asymmetry, and line-depth dependence of the convective blueshift.

Inspired by Gray’s foundational research [6], this project seeks to extract the granulation pattern from the IAG Solar Flux Atlas to calculate relative velocities and perform detailed analysis. A parallel focus will investigate the chromatic dependence detected in absorption line profiles (the third signature of convection), which currently obscures the universality of the pattern.

1.1 Convective motion

The stellar spectrum serves as astronomy’s primary source of information, particularly regarding a star’s composition and relative velocity. However, convective motions in the stellar photosphere complicate spectral interpretation by inducing differential velocities in individual spectral lines. Specifically, fluid movements caused by density variations from temperature fluctuations in the Sun’s outermost layer modify the spectrum, causing each spectral line to display distinct relative velocities as we can relate to different research [7, 8, 6, 9]. Persistent convective motion generates a granular structure in solar photospheric images. One of the first person to discover the pattern was Janssen in 1885, who detected a granulation movement in the photosphere. Lately, in 1901 Plaskett associated this pattern with the same of convective cells in Bernard’s experiment [3]. Where fluids heated from below representing hot rising gas elements convecting heat to surface. This characteristic configuration gives the name *granulation* to all observable signs of the convection.

1.2 Solar granulation pattern

A plot of Doppler shift against line depth is called *Granulation Pattern*; it shows that weaker lines are more blueshifted. When the Sun pushes material up through its outer layer, the spectrum exhibits a blueshift. As this material subsequently cools and falls back through the atmosphere, it produces a redshift, but emits less light, making the blueshift dominant. This phenomenon has been particularly documented by David Gray, whose work has significantly improved the precision of stellar radial velocity measurements. This improvement stems primarily from Gray’s observation that granulation patterns in stars resemble solar patterns, differing only by a scaling factor. These findings are particularly significant given the considerable challenges in obtaining such parameters for other stars, which are affected by spectral noise, stellar proper motions, and relative velocity uncertainties [6]. Furthermore, analysis of solar granulation patterns contributes to the understanding and validation of photospheric hydrodynamic mod-

els [7]. Such analysis enables improved calibration when testing dynamic atmospheric models.

1.2.1 The three signatures of convection

Due to convection motion, there's three distinct granulation signatures can be identified in stellar spectra: Line broadening, line profile bisector asymmetry, and line-depth dependent shifts in absorption lines [10]. The third signature exhibits a correlation between line depth and wavelength shift[6], for which neutral iron lines (Fe I) serve as optimal calibration references due to their high abundance, minimal thermal broadening and limited isotopic variation. This approach offers the additional advantage of deriving natural wavelengths from a single source, thereby eliminating potential discrepancies. On the way of trating th Sun as any other star, which does not have the proximity for spectra study, we search for obtain the most accurate and precise granulation pattern. Which is posible if we take the most accurate and precise solar atlas.

1.3 IAG Solar Flux Atlas

In 2016, Reiners et al.[9] published the unprecedented precision *Institut für Astrophysik Göttingen (IAG)* Solar Flux Atlas obtained with the FTS *Fourier Transform Spectrograph*, simultaneously reporting convective blueshifts for a sample of neutral iron lines. This atlas provides highly precise and accurate data, with radial velocity uncertainties on the order of $\pm 10 \text{ m/s}$ across the wavelength range of 4050 to 10650 Å. In contrast to other FTS atlases, the entire visible wavelength range was observed simultaneously using only one spectrograph setting[9].

In 2016, the resulting granulation pattern appeared notably scattered and noisy, attributable to the rudimentary line position measurement methodology employed, and the poorly curated line selection, which included numerous blended features, outdated wavelength references and incomplete spectral coverage. The exceptional quality of the IAG spectrum enables more accurate determination of convective blue-shifts when

analyzed through refined methods.

1.4 IAG Spatially Resolved Quiet Sun Atlas

In 2023, Ellwarth et al.[5] observed and published the *Institut für Astrophysik Göttingen (IAG)* Spatially Resolved Quiet Sun Atlas obtained with the FTS *Fourier Transform Spectrograph*. This atlas has the advantage of use observations from the disc center ($\mu = 1.0$) towards the solar limb ($\mu = 0$), where $\mu = \cos(\theta)$. The research had the objective of study the blueshift exhibits variations from the disc centre to the solar limb due to differing projection angles onto the solar atmosphere.

Our principal objective is to establish a characterization of the granulation pattern by treating the Sun as any other star. This approach allows us to develop methods that can be directly scaled and applied to other solar-type stars. However, to accurately analyze relative velocities and dynamics, we must account for insights that can only be corrected using the Spatially Resolved Quiet Sun Atlas.

1.5 Motivation

During the first semester of 2025, physics student on the Universidad de los Andes Manuel Fuentes implemented these improvements in his computational project, focusing specifically on the visible spectral range $4050 - 10650 \text{ \AA}$. By developing enhanced measurement techniques and employing a carefully curated line list that fully covers the IAG-VIS range with updated wavelength references [11], Fuentes achieved significantly sharper granulation patterns compared to previous analyses. This optimized approach demonstrates how proper line selection and modern wavelength standards can extract more reliable convective signatures from high-quality solar spectra.

With professor Benjamin, we continue with a measurement of granulation pattern taking into account the near infrared range. This region contains spectral lines originating from deeper layers of the solar photosphere, which are consequently weaker. These lines

represent a rich source of valuable information [12], though they require adapting the measurement methodology. Furthermore, we start a characterization of solar dynamics and line asymmetries based on granulation pattern observations.

By addressing these challenges, we aim to achieve two key outcomes: A robust characterization of solar granulation pattern and insights into the photosphere's geometry. These efforts are guided by the central research question: What are the direct consequences of solar dynamics on its spectrum?

Chapter 2

Literature Review: Convective movement in the Sun

As previously mentioned, David Gray [6, 10, 13] has significantly advanced the study of granulation patterns in the solar photosphere, with a particular focus on measuring their associated relative velocities with high precision. These developments have enabled more accurate characterizations of other stars by extrapolating the physical principles observed in the Sun. This chapter explores the three signatures of convective motion in the Sun photosphere, and how this reveal the hydrodynamic on this outermost layer. Furthermore, is given the point of view of different authors respect the reasons behind the chromodepende on the granulation patterns.

2.1 The solar interior and the solar outer atmosphere

The Sun is classified as a yellow dwarf star of spectral type G2V, title achieved for a big amount of hot hydrogen (ionised H in 90 percent) and helium (in 10 percent). What makes this star really unique is his proximity to earth and the facility to study from the planet with precision. In general, the Sun is divided in two fundamental parts: The

solar interior and the solar outer atmosphere.




Figure 2.1: The interior structure of the Sun. The convection zone is the responsible by the general movement that characterize the third signature.

As shown in the figure 2.1 the overall structure of the solar interior is core, radiative and convective zone. Across then the density and temperature falls significantly, as the energy is slowly transferred outwards by radiative diffusion. Some models of the interior structure give a core temperature of $1.6 \times 10^7 K$ and density to $1.6 \times 10^5 Kg/m^3$, high enough for thermonuclear reactions and remains the central material in plasma like a gigantic atomic reactor. This characteristic allow the collisions, absorptions and reemissions of photons that made this zone opaque. In consequence, there exists an increase of the wavelength from high-energy gamma rays to visible lighth.

For this research, the two target layers of the Sun are the convection zone and the photosphere.




Figure 2.2: The first clear photograph of the photosphere where is visible the granulation pattern. Image taken from [1]

As we shown in the figure 2.2, in 1877 Janssen took the first clear photograph of the granules in the photosphere [1]. This was the starting point for different studies across the pattern of granules. In 1930 Unsold pointed out that the layers below the photosphere should be convective unstable [3]. Later, Plasketts relationed these granules with the Bernard's laboratory measurements of fluids convection. This statement are based on fluids heated from below representing hot rising gas elements convecting heat to surface [14]. This hot rising gas elements are known as granulation, and each individual region—spanning approximately 700 km in size and lasting between five to ten minutes—is referred to as a granule. The understanding of photospheric granules as convective cells leads the thought to the existence of a zone responsible of the convective motion.

2.2 The solar convection Zone


Starting at the $0.86R_{\odot}$ lies the zone where the dynamics processes took place, the great temperature gradient across the layer allow the process of convection [15].

2.2.1 The convection movement in the sun

As we mentioned, the convective movement are based on fluids heated from below representing hot rising gas granules or *convective cells* convecting heat to the photosphere [14]. In this case, solar convection occurs in a highly compressible, stratified gas which leads to determine the conditions under which we expect convection and the dynamics of the granules [3].

2.2.2 Dynamics of solar convection

From the core, He nuclei is built from H nuclei in the proton-proton cycle as say equation (2.1).



Where from the H nuclei is liberated an considerable amount of high frequency γ -rays ($26.7MeV$) and the energy of two neutrinos ($0.5MeV$). However, the strong Coulomb repulsion between positively charged nuclei increases as the product of their nuclear charges, so only lightest elements will have appreciable reaction probabilities. As the electrons recombine with others particles the photons can be absorbed more easily. With this, decrease the radiative conductivity and increases the temperature gradient with the opacity [3]. The decrease of temperature allow a convective instability and beyond a convective turbulence. When reaches the low photosphere, some radiation scape from the sun and the material returns to the convective stability [15].

This inclines to establish an onset of convection: If T is increased to a value T' , the granule will expand rapidly to achieve a new pressure equilibrium with its surroundings. Where is suppose the granules as vertical stratified plasma in hydrostatic equilibrium with pressure ($P(r)$); density ($\rho(r)$); and temperature ($T(r)$), the same as its surroundings at radial distance r from the center of the star.

The lower-density gas experiences a buoyancy force, which will cause it to rise. The buoyancy force will disappear when the density has dropped to the same value as the new surroundings, after an element has traveled a distance l . Let be T'_n the new temperature

and that of its surroundings T_n , the adiabatic and radiative gradient for the element follows the equation (2.2).

$$T_n = T + \left(\frac{dT}{dr} \right)_R l \quad | \quad T'_n = T' + \left(\frac{dT}{dr} \right)_{ad} l \quad (2.2)$$

Where ad references for adiabatic gradient; R is for radiative gradients; and taking l small. Since we have assumed radiative equilibrium, the R denotes the gradient present in the stellar atmosphere. Two conditions can arise from here (see equation (2.3)): The radiative gradient is unstable, so convection pattern is stablish and viceversa.

$$- \left(\frac{dT}{dr} \right)_R > \left(\frac{dT}{dr} \right)_{ad} \quad (2.3)$$

If the convection pattern is stablish, the element continues to expand further and rise. Otherwise, the element begins to contract, becomes heavier, and begins to move down to its original position.

This onset of instability when the vertical temperature gradient is too large is explained by the Schwarzschild condition.

2.2.3 The Schwarzschild condition

Taking the element described before, now suppose an elementary parcel of the material is displaced so slowly that remains in horizontal pressure equilibrium. If the motion is adiabatic there is no heat exchange with surroundings, the rate between the density and pressure is constant. This generates a criterion for the presence of convection known as the Schwarzschild condition (2.4).

$$- \frac{dT}{dr} > \frac{\gamma - 1}{\gamma} \left(\frac{GM_{\odot}m}{r^2 k_B} \right) \quad (2.4)$$

Where k_B refers to Boltzman constant; G the gravitational constant; M_{\odot} the solar mass; m the mass of the granule; and γ the degree freedom of the fluid. As the presence of convection reduces the temperature gradient from the higher value it would have assumed under radiative transport alone to the essentially adiabatic value [3].

All of the material and energy generated by convection ended up to low photosphere, where the granules shows different properties examined in after sections.

2.3 The Solar Photosphere

Since 1874, when Langley gives a detailed description of granulation on the photosphere, the astronomers have been studying different motions and reactions on the outermost solar layer [15]. The distinct pattern of granules with dynamic behavior becomes apparent, where individual areas continuously emerge and disappear (see figure 2.3).




Figure 2.3: A view of granulation on the Sun's surface. The central regions exhibit blueshifts while the edges display redshifts. Image taken from [2]

The bright areas of granules correspond to regions where hot gas rises through the solar atmosphere. As this gas releases energy in the form of photons at the photosphere, it cools and subsequently descends, creating the darker regions of granules [16, 7].

2.3.1 Static photosphere: Limb darkening phenomenon

Because the temperature decreases from the higher layers, the photosphere intensity falls off towards the solar limb. Consequently, the disk intensity profile becomes more squared at increasing wavelength (see figure 2.4).




Figure 2.4: Squared profile for the disk intensity at increasing wavelengths. Image taken from [3]

This effect is known as *Limb darkening phenomenon*. The analysis of this effect provides a direct technique for determining the photosphere temperature structure along depth. Furthermore, these granules has shown to be in continual motion (see figure 2.5).




Figure 2.5: The image shows frame by frame the motion of few granules. Image taken from [3]

This perpetual motion across the layer generates asymmetries on the absorption lines.

2.3.2 Dynamic photosphere: The C-curved profile bisector.

Observing the dynamic of the granules, there appear a height dependence of the granular velocities: The velocity of an upward moving granule decays much less rapidly than its excess brightness. Changes in the granulation structure, contrast and velocity field around the spots and network have been inferred indirectly from observations of Fraunhofer line profile shapes [3]. The result is characteristically C-Curved profile bisector (see figure 2.6).




Figure 2.6: The C-curved profile bisector. In the infrared shown to be less pronounced than violet range.

The process of the creation in this c-curved profile bisector is divided in three stages. First, the line profile near its mid-depth portion is formed in the most rapidly upflowing bright material (Blueshifted). Then, the deepest portion of the line core is formed higher in the decelerated upflow (Less Blueshifted). Finally, the line wings where the opacity is least, tend to be formed deepest in the cool (Redshifted).

This dynamic process induces perturbations in spectral absorption lines, line profile asymmetries, and line depth-dependent wavelength shifts [10] known as the three signatures of convection.

2.4 The three signature of convection

The signatures of convection in stars are described by Gray in his research [6, 13, 10] about the principal characteristics in the spectra to identified and treat the asymmetries due to convective motion.

2.4.1 Line broadening

Due to the limb darkening phenomenon and atmospheric absorption, there exists two types of line cores: Strongest and weaker lines [10]. Where strongest refers to more curved than weaker lines as is shown in the figure 2.7.




Figure 2.7: In the figure is compared the strongest and the weaker line core shapes.

This effect is measurable with the second derivate in the observed wavelength, that representing the core curvature (see equation (2.5)).

$$C_c = \lambda^2 \left(\frac{d^2 p(\lambda)}{d\lambda^2} \right) \quad (2.5)$$

Where $p(\lambda)$ is the polinomial fit for the core evaluated on the observed wavelength. In the near infrared range we can denote a natural bands or groups of lines which are separated for the teluric elements, on others words, elements like oxygen and CO2 that are absorbed by the atmosphere. Tis range are invisible for the spectra, and it helps us to detected the natural bands in the near infrered curvature profile.




Figure 2.8: Asymmetries on an average absorption line. Can be observed the differences in intensity redshift profile. Image taken from [4]

2.4.2 Line profile bisector asymmetry

As demonstrated by Nieminen [4], the asymmetry occurs because the c-curved profile bisector reflects velocity variations and a bisector slope (see figure 2.8).

According to Kirchhoff's laws, absorption line formation requires lower temperature conditions, which are found precisely in the Sun's outermost atmospheric layers [16]. These regions not only provide the appropriate temperatures for absorption line formation but also exhibit comparatively higher opacity. Among all available spectral lines, those from neutral iron (Fe I) are particularly valuable for solar granulation studies due to two key characteristics: they display significant opacity and relatively low thermal broadening. These properties make Fe I lines excellent tracers of granulation patterns in the solar photosphere [4].

These asymmetries can be measured using the third derivate relation for the line bisector (see equation (2.6)).

$$\frac{c}{\lambda} \left(\frac{\frac{d^3 p(\lambda)}{d\lambda^3}}{3C_c^2} \right) \quad (2.6)$$

Where C_c refers to the core curvature in the equation (2.5).




Figure 2.9: Granulation pattern for the IAG spatially resolved quiet sun atlas, is shown to have a strong chromodependence in the weaker lines. Image taken from [5].

2.4.3 Line depth-dependent wavelength shifts

Many studies across the years have detected and observed the phenomena of wavelength shifts against the line depth, or as it is called, chromodependence in the granulation pattern. This behavior was showed to be more present in the weaker lines which are related to infrared and violet range.

2.4.4 Solar granulation pattern

The solar granulation pattern is a plot of relative velocity against line depth, as shown in the figure 2.9.

The importance of this pattern lies in stars resemble solar patterns, differing only by a scaling factor (see [10]). Furthermore, these analysis contributes to the understanding and radiation of photospheric hydrodynamic models [7, 6]. However, for the measurement is necessary the understanding of the convective blueshift.

Convective Blueshift

When the Sun pushes material up through its outer layer, the spectrum exhibits a blueshift. As this material subsequently cools and falls back through the atmosphere, it produces a redshift, but emits less light, making the blueshift dominant. That can be measure by the Doppler effect but as the velocities can be significant, the relativistic formulation of this effect must be applied.

The relativistic Doppler effect accounts for length contraction, as predicted by Einstein's theory of relativity. This introduces an additional correction term to the classical shift, which becomes particularly relevant in high-velocity scenarios or strong gravitational fields. However, the measured negative redshift resulting from convective motions is known as *convective blueshifts* which are measured by the equation (??).

$$v_r \approx c \left(\frac{\lambda_{obs} - \lambda_{rep}}{\lambda_{rep}} \right) \quad (2.7)$$

Since the strength of the convective distortions and shifts as spectral lines varies across the H-R diagram, we expect the systematic errors in radial velocities [6].

2.4.5 Chromodependence characterization

Recently, the emphasis on the measurement of granulation pattern has opened new interpretations to the line depth-dependent wavelength shifts. However, there is still no characterization for this phenomenon.

In 2018 Gray and Oostra try to stablish a standard curve determined by the second grade polynomial fit to the solar granulation pattern. Although, the authors just take into account the range of 4950\AA to 5700\AA of the spectrum, avoiding the chromodependence in the weaker lines. This motivate to search a strong characterization and give the recipe for dealing with this phenomenon.

Hamilton and Lester, give a little theory that this phenomenon is atribute to the rotation in the photosphere. The pronounced differential rotation with latitude observed at the photosphere seems to be the result of convective flows driven radially by the bouyancy force and deflected horizontally by the coriolis force due to solar rotation [3]. This




Figure 2.10: Contributions on angular momentum. Image taken from [3].

rotation made contributions on angular momentum in two different forms: Meridian circulation and Reynolds stresses.

The first contribution occurs if axysimmetric meridional circulation are present (see figure 2.10). In the absence of any other angular momentum transport, a circulation in either sense will tend to spin up the poles and the interior because the fluid carries angular momentum.

The second contribution tends to enforce solid body rotation, then the meridional circulation drive an equatorial acceleration. The reason is for equal velocity in the meridional plane, the flux of angular momentum per unit mass across the dashed line will be larger toward the equator than away from it. This mechanism depends on the existence of nonaxisymmetric convective motion because net fluxes of angular momentum in latitudinal or radials directions are produced without requering a net mass flux. Neither buoyancy forces, which are strictly radial, nor presure gradients, which must average to zero around the solar circumference, can themselves influences the suns axysimetric rotation profile[3].

Chapter 3

Methodology: The blend-free list of Fe I lines

As the spectral analysis is complicated cause the convective motion, I take an computational approach leading an observational requirement to lead the research. This leading by the experience of professor Benjamin and the previous research. This research endeavor encompass both analytical and computational components. For the last part, the programming language Python and Jupyter Notebook was used. The computational aspect focus on identifying the granulation pattern within the solar spectrum by calculating relative velocities using the wavelengths of Fe I. All analysis data and code used in the process was uploaded to a GitHub repository, allowing anyone interested to reproduce the results and verify the authenticity of the conclusions presented. Moreover, proper credit will be given to all previous work from other researchers.

3.1 Methodology

We follow the methodology established in previous studies [7, 9, 5] , which utilized a selected list of Fe I lines. These lines are ideal for this calibration due to their minimal thermal broadening and reduced susceptibility to other atmospheric affectations.

The line up for the code was identify the Fe I lines in the IAG Solar Flux Atlas and the

IAG Spatially Resolved Quiet Sun Atlas using the blend-free list of Fe I lines. Second, fit a fourth-grade polynomial fit due to the c-curved line profile bisector and find the observed wavelength. Then calculate the Doppler velocity, convective blueshift and flux with the fit. Finally, find the second derivate (core curvature) and the third derivate relation (bisector slope) for the curvature in the observed wavelength. This last is develop in detail in the appendix B.

3.2 Blend-free Fe I line list

The previous methodology implemented the Nave et. al list of laboratory measured Fe I lines [11]. This list classifies lines with a quality rating (A,B,C,D) with A be the most and best quality. However, not all the listed lines are clearly present in the solar spectrum, and within the near-infrared range, many lines are severely mixed.

In collaboration with Professor Benjamin and Manuel Fuentes, we refined this list.

3.2.1 Selection method for Fe I lines

This selection method has two different approaches: Computational and visual. First, we selected only quality A lines and then performed the next filters according to the statement of having an fourth order polynomial fit for the line core.

- The coefficient of the fourth grade term need to be positive and not too small cause it denotes a curve too big. This can be for a line, specially in the infrared, cause the big cores are related to atmospheric lines.
- The difference between extreme points of the fit need to be less than the half of the distance, that is a curve and not a slope.
- The absolute difference between the wavelength observed and the one from Nave list need to be less to 0.025\AA .

The last part was a visual inspection to discard line mixes or absent from the solar spectrum. This part was divided between porfessor Benjamin and me. I used the

visualizer (explained in detail in the appendix C) and profesor Benjamin use the great years of experience. Then I combined the list, made a check and profesor Benjamin check again. The discard part was only using the geometry of the curve and guidance of profesor Benjamin (and God). The new blend-free list improves our results, leading to cleaner graphics and characterization with less scatter in the data. Then we perform different graphics for our analysis conform we see it necessary for the characterization. In case of the IAG Spatially Resolved Quiet Sun Atlas we only use the $\mu = 1$ for the effect to eliminate the limb darkening effect in the spectrum.

Chapter 4

Results and discussion: Granulation pattern and Characterization of chromodependence

Our results can be summarized in three principal aspects: Chromodependence on the granulation pattern along the line depth; a detailed view of the characteristic curvature, asymmetries, and sharpness of spectral lines and higher-quality graphs with reduced scatter.

4.1 The third signature: Chromodependence on the granulation pattern

We obtain the granulation patterns for the IAG Solar Flux Atlas for all the wavelength.




Figure 4.1: Granulation pattern for the Solar Flux Atlas.

The behavior along the line depth is according to literature, in which is clear the chromodependence. For a different try into the characterization of this behavior we try to generate different second order polynomial fits for each color range.




Figure 4.2: Granulation pattern for the Solar Flux Atlas with color curves.

But this is not standard. So, we performed an analysis of line depth against wavelength. Our hypothesis was: “If the chromodependence is present only in the Solar Flux Atlas, then rotation could be the cause of this phenomenon” Surprisly, we observed a chromodependence in both spectral datasets. We start taking the granulation pattern of the center disk




Figure 4.3: Granulation pattern for the Spatially Resolved Quiet Sun Atlas at $\mu = 1$.

This was initially unexpected because the rotation is negligible at the disc center. For the measurement of its velocity we took a 4300-5600 Å range and sorted all lines from both atlases into 50 m/s velocity bins.




Figure 4.4: Comparison between atlases with velocity bins for the relation between wavelength and line depth.

For the measurement of rotation in the lines we made a first order polynomial fit for each one.




Figure 4.5: Comparison between atlases with velocity bins for the relation between wavelength and line depth. The first order fit is showed for each velocity bin.

The clue here is the value of slopes in the Spatially Resolved Atlas is greater than the Solar Flux, which contrindicates for complete the initial hypothesis. The rotation is not the cause.

4.2 The first and second signature: Detailed view of line broadening and asymmetry

We made two graphics corresponding to each signature related to the line bisector and core asymmetries. The slope of line bisector shows the behavior of the c curved line profile bisector.




Figure 4.6: C bisector graph for slopes in which is clear the convection movement and how its affected in the profile.

The interesting one is the core curvature but multiplied by the squared wavelength, cause it affirms the chromodependence in the weaker lines as the literature says.




Figure 4.7: Sharpness of the core bisector in the solar flux atlas separated by range.

We can see a natural division for lines in the infrared range, which is interesting due to corresponding Teluric lines of absorption in the atmosphere. There is a little division in 11400 Å who separates the lines in one that follow the mean curve and others that don't do that.




Figure 4.8: Sharpness of the core bisector in the solar flux atlas. Is clear that the infrared follow two different behavior due teluric lines.

This is the graphic which is clear that we have chromodependence in the weaker lines.

4.3 Higher quality graphs

In comparison of Ellwarth graphics, which is the study most recent, there is a better resolution in the different graphs without scattered points.




Figure 4.9: We realized the same graphic for the comparison with the Ellwarth article to show the less scattered points.

Chapter 5

Conclusions

In general, we conclude... Characterize the convective blueshift of solar absorption lines and its dependence on line depth and wavelength range.

In specific, we produce a blend-free list of Fe I absorption lines and solar granulation patterns with minimal scatter. We describe in detail how the granulation plot depends on wavelength range. We explore possible explanations of wavelength-dependence and found that () is the best. We explore ways of dealing with the wavelength-dependence when using the solar spectrum as a gauge for stellar spectra.

Appendix A

Z-score Standardization

This was for a problem present in the fit and we just use a computational standardization in the selected bins of fits to make a standardization.

Appendix B

The third derivate relation

Or called the bisector slope. It was multiplied by the relation $(\frac{c}{\lambda})$ to see each clear in the graphic.

Appendix C

Visualizer for outliers

Was an app made with Tkinter for help the visualization of outliers in the process of blendfree the Fe I list. There is in the Git hub

Bibliography

- [1] J.-M. Malherbe, “Jules janssen, the birth of solar physics, the foundation of meudon observatory and the mont blanc adventure (1875-1895),” *HAL CNRS*, Aug. 2022.
- [2] S. A. Hamouda, F. F. Alfarjani, and F. Y. Elfituri, “Sunspots production and relation to other phenomena: A review,” *International Journal of Science and Research Methodology*, 06 2018.
- [3] F. Peter, *Solar Astrophysics*. Cambridge research and instrumentation, 1990.
- [4] T. A. Nieminen, “Solar line asymmetries: Modelling the effect of granulation on the solar spectrum,” *arXiv*, Aug. 2017. arXiv:1708.06408 [astro-ph].
- [5] M. Ellwarth, B. Ehmman, S. Schäfer, and A. Reiners, “Convective characteristics of fe i lines across the solar disc,” *Astronomy and Astrophysics*, vol. 680, p. A62, Dec. 2023.
- [6] D. F. Gray, “The third signature of stellar granulation,” *The Astrophysical Journal*, vol. 697, p. 1032, May 2009.
- [7] D. Dravins, L. Lindegren, and A. Nordlund, “Solar granulation - influence of convection on spectral line asymmetries and wavelength shifts,” *Astronomy and Astrophysics*, vol. 96, p. 345–364, Mar. 1981.
- [8] D. Hamilton and J. B. Lester, “A technique for the study of stellar convection: The visible solar flux spectrum,” *Publications of the Astronomical Society of the Pacific*, vol. 111, p. 1132, Sept. 1999.

- [9] A. Reiners, N. Mrotzek, U. Lemke, J. Hinrichs, and K. Reinsch, “The iag solar flux atlas: Accurate wavelengths and absolute convective blueshift in standard solar spectra,” *Astronomy and Astrophysics*, vol. 587, p. A65, Mar. 2016.
- [10] D. F. Gray and T. Pugh, “The third signature of granulation in bright-giant and supergiant stars,” *The Astronomical Journal*, vol. 143, p. 92, Mar. 2012.
- [11] G. Nave, S. Johansson, R. C. M. Learner, A. P. Thorne, and J. W. Brault, “A new multiplet table for fe i,” *The Astrophysical Journal Supplement Series*, vol. 94, p. 221, Sept. 1994.
- [12] A. Cacciani, R. Briguglio, F. Massa, and P. Rapex, “Precise measurement of the solar gravitational red shift,” *Celestial Mechanics and Dynamical Astronomy*, vol. 95, p. 425–437, May 2006.
- [13] D. F. Gray and B. Oostra, “The solar-flux third granulation signature,” *The Astrophysical Journal*, vol. 852, p. 42, Jan. 2018. ADS Bibcode: 2018ApJ...852...42G.
- [14] H. H. Plaskett, “Solar granulation,” *Monthly Notices of the Royal Astronomical Society*, vol. 96, p. 402, Mar. 1936. ADS Bibcode: 1936MNRAS..96..402P.
- [15] P. Eric, *Solar magnetohydrodynamics*. Kluwer, 1982.
- [16] B. Carroll and D. Ostlie, *An Introduction to Modern Astrophysics*. Cambridge University Press, 2017.
- [17] J. S. Aponte, “Medición de la velocidad convectiva en la fotosfera solar,” 2017.
- [18] F. Stief, J. Löhner-Böttcher, W. Schmidt, T. Steinmetz, and R. Holzwarth, “Convective blueshifts in the solar atmosphere - ii. high-accuracy observations of the fe i 6173.3 Å line and deviations of full-disk dopplergrams,” *Astronomy & Astrophysics*, vol. 622, p. A34, Feb. 2019.