

Assignment

Section 3.4: 2(fj), 8, 11, 14, 15; Section 4.1: 10; Section 4.2: 23, 29, 30; Section 4.3: 10, 11, 12, 15

Work

3.4

2. Use Gaussian elimination to solve the following systems of linear equations.

(f)

$$\begin{aligned}x_1 + 2x_2 - x_3 + 3x_4 &= 2 \\2x_1 + 4x_2 - x_3 + 6x_4 &= 5 \\x_2 + 2x_4 &= 3\end{aligned}$$

$$\left(\begin{array}{cccc|c}1 & 2 & -1 & 3 & 2 \\2 & 4 & -1 & 6 & 5 \\0 & 1 & 0 & 2 & 3\end{array}\right) \begin{array}{c} \boxed{-2} \leftarrow + \quad \boxed{+} \leftarrow \\ \boxed{+} \leftarrow \quad \boxed{1} \leftarrow \\ \boxed{-2} \leftarrow \end{array} \rightsquigarrow \left(\begin{array}{cccc|c}1 & 0 & 9 & -4 & -3 \\0 & 1 & 0 & 2 & 3 \\0 & 0 & 1 & 0 & 1\end{array}\right) \quad (1)$$

$$x_1 = -3 + 4x_4 \quad (2)$$

$$x_2 = 3 - 2x_4 \quad (3)$$

$$x_3 = 1 \quad (4)$$

$$x_4 = x_4 \quad (5)$$

$$S = \left\{ \begin{pmatrix} -3 \\ 2 \\ 1 \\ 0 \end{pmatrix} + z \begin{pmatrix} 4 \\ -2 \\ 0 \\ 1 \end{pmatrix} : z \in \mathbb{R} \right\} \quad (6)$$

(j)

$$2x_1 + 3x_3 - 4x_5 = 5 \quad (7)$$

$$3x_1 - 4x_2 + 8x_3 + 3x_4 = 8 \quad (8)$$

$$x_1 - x_2 + 2x_3 + x_4 - x_5 = 2 \quad (9)$$

$$-2x_1 + 5x_2 - 9x_3 - 3x_4 - 5x_5 = -8 \quad (10)$$

$$\left(\begin{array}{ccccc|c}2 & 0 & 3 & 0 & -4 & 5 \\3 & -4 & 8 & 3 & 0 & 8 \\1 & -1 & 2 & 1 & -1 & 2 \\-2 & 5 & -9 & -3 & -5 & -8\end{array}\right) \rightsquigarrow \left(\begin{array}{ccccc|c}1 & 0 & 0 & 0 & -2 & 1 \\0 & 1 & 0 & 0 & -3 & 0 \\0 & 0 & 1 & 0 & 0 & -1 \\0 & 0 & 0 & 1 & -2 & -1\end{array}\right) \quad (11)$$

$$x_1 = 2x_5 + 1 \quad (12)$$

$$x_2 = 3x_5 \quad (13)$$

$$x_3 = -1 \quad (14)$$

$$x_4 = 2x_5 - 1 \quad (15)$$

$$x_5 = x_5 \quad (16)$$

$$S = \left\{ \begin{pmatrix} 1 \\ 0 \\ -1 \\ -1 \\ 0 \end{pmatrix} + z \begin{pmatrix} 2 \\ 3 \\ 0 \\ 2 \\ 1 \end{pmatrix} : z \in \mathbb{R} \right\} \quad (17)$$

8. Let W denote the subspace of \mathbb{R}^5 consisting of all vectors having coordinates that sum to zero. The vectors

$$\begin{aligned} u_1 &= (2, -3, 4, -5, 2), & u_2 &= (-6, 9, -12, 15, -6), \\ u_3 &= (3, -2, 7, -9, 1), & u_4 &= (2, -8, 2, -2, 6), \\ u_5 &= (-1, 1, 2, 1, -3), & u_6 &= (0, -3, -18, 9, 12), \\ u_7 &= (1, 0, -2, 3, -2), & u_8 &= (2, -1, 1, -9, 7) \end{aligned}$$

generate W . Find a subset $\{u_1, u_2, \dots, u_8\}$ that is a basis for W .

$$\mathbb{R}^5 = \left\{ \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} : x_1 + x_2 + x_3 + x_4 + x_5 = 0, x_1, \dots, x_5 \in \mathbb{R} \right\} \quad (18)$$

$$\begin{pmatrix} 2 & -6 & 3 & 2 & -1 & 0 & 1 & 2 \\ -3 & 9 & -2 & -8 & 1 & -3 & 0 & -1 \\ 4 & -12 & 7 & 2 & 2 & -18 & -2 & 1 \\ -5 & 15 & -9 & -2 & 1 & 9 & 3 & -9 \\ 2 & -6 & 1 & 6 & -3 & 12 & -2 & 7 \end{pmatrix} \quad (19)$$

$$\rightsquigarrow \begin{pmatrix} 1 & -3 & 0 & 4 & 0 & 1 & 0 & -1 \\ & & 1 & -2 & 0 & -2 & 0 & 1 \\ & & & & 1 & -4 & 0 & -2 \\ & & & & & & 1 & -1 \\ & & 0 & & & & & 0 \end{pmatrix} \quad (20)$$

It follows that $\{u_1, u_3, u_5, u_7\}$ is linearly independent by theorem 3.16. Therefore $\{u_1, u_3, u_5, u_7\}$ is a basis for W .

11.

14.

15.

4.1

10.

4.2

23.

29. Prove that if E is an elementary matrix, then $\det(E^t) = \det(E)$.

(a) **Types 1 & 2**

$$E^t = E \quad (\text{by HW.3.1.5}) \quad (21)$$

$$\Rightarrow \det(E^t) = \det(E) \quad (22)$$

(b) **Type 3**

E^t is an type 3 elementary matrix (by HW.3.1.5) $\det(E) = \det(I) = 1$ for any type elementary operation on I_n

$$\det(E^t) = \det(I) \text{ because } E^t \text{ is type 3} \quad (23)$$

$$\Rightarrow \det(E) = \det(E^t) = 1 \quad (24)$$

30. Let the rows of $A \in M_{n \times n}(F)$ be a_1, a_2, \dots, a_n and let B be the matrix in which the rows are a_n, a_{n-1}, \dots, a_1 . Calculate $\det(B)$ in terms of $\det(A)$.

(a) **n is even**

In A , swap

$$a_{n-1} \text{ with } a_1 \quad (25)$$

$$a_{n-2} \text{ with } a_2 \quad (26)$$

\vdots

$$a_{n-\frac{n}{2}+1} \text{ with } a_{n-\frac{n}{2}} \quad (27)$$

From the fact that $n/2$ swaps were performed it follows from Theorem 4.6 that

$$\det(B) = (-1)^{\frac{n}{2}} \det(A) \quad (28)$$

(b) **n is odd** In A , swap

$$a_{n-1} \text{ with } a_1 \quad (29)$$

$$a_{n-2} \text{ with } a_2 \quad (30)$$

\vdots

$$a_{n-\frac{n+1}{2}+1} \text{ with } a_{n-\frac{n+1}{2}} \quad (31)$$

From the fact that $n - \frac{n+1}{2}$ swaps were performed it follows from Theorem 4.6 that

$$\det(B) = (-1)^{\frac{n-1}{2}} \det(A) \quad (32)$$

4.3

10.

11.

12.