

Delphi-Stat Forecasting Methodology EW44 Update

Logan C. Brooks¹ David C. Farrow¹ Shannon Gallagher² Sangwon Hyun²
Ryan J. Tibshirani² Roni Rosenfeld¹

¹ School of Computer Science ² Department of Statistics
Carnegie Mellon University,
Pittsburgh, Pennsylvania,
United States of America

Changes to Delphi-Stat Methodology

New method added to ensemble: modified time-weighted kernel density estimation: this version changes the weighting criteria used for matching simulated data for this year to past seasons; attempts to make simulated trajectories more closely resemble past seasons' data; and considers a wider range of past data by looking at other epi weeks.

When constructing trajectories, this version weights past seasons based on the previous week's wILI value; the sum of previous wILI values in the season; a weighted sum of wILI values stressing more recent weeks; and a weighted sum of the week-to-week changes in wILI stressing more recent times. With low probability, these weights are ignored and a random change in wILI is selected from historical data. The simulated data values are also pushed towards randomly selected historical data by a small amount. When simulating data at epi week t , instead of just looking at other seasons at week t , also considers nearby weeks, unless t is a time near the end of year holidays.

Clarification: older kernel density estimation method, direct target density estimation: only weight data based on the previous wILI value.