

Introduction to Longitudinal Data Analysis

Christopher David Desjardins
Chu-Ting Chung
Jeffrey D. Long

11 January 2010

Outline

- 1 Introduction
- 2 Benefits
- 3 Synonyms
- 4 Formula and Assumptions
- 5 LMMs in R
- 6 Getting Help

What is Longitudinal Data?

- The defining characteristic of longitudinal studies is that individuals are **measured repeatedly through time**.
 - In contrast, cross-sectional studies involve collecting measurements on an individual **only once**.
- Concerned with **mean change** over time.
- Longitudinal studies allow examination of **age and cohort** effects.
 - Longitudinal studies can distinguish changes over time within individuals (aging effects) from differences among people in their baseline levels (cohort effects).

What is Longitudinal Data?

- The defining characteristic of longitudinal studies is that individuals are **measured repeatedly through time**.
 - In contrast, cross-sectional studies involve collecting measurements on an individual **only once**.
- Concerned with **mean change** over time.
- Longitudinal studies allow examination of **age and cohort** effects.
 - Longitudinal studies can distinguish changes over time within individuals (aging effects) from differences among people in their baseline levels (cohort effects).

What is Longitudinal Data?

- The defining characteristic of longitudinal studies is that individuals are **measured repeatedly through time**.
 - In contrast, cross-sectional studies involve collecting measurements on an individual **only once**.
- Concerned with **mean change** over time.
- Longitudinal studies allow examination of **age and cohort** effects.
 - Longitudinal studies can distinguish changes over time within individuals (aging effects) from differences among people in their baseline levels (cohort effects).


What is Longitudinal Data?

- The defining characteristic of longitudinal studies is that individuals are **measured repeatedly through time**.
 - In contrast, cross-sectional studies involve collecting measurements on an individual **only once**.
- Concerned with **mean change** over time.
- Longitudinal studies allow examination of **age and cohort** effects.
 - Longitudinal studies can distinguish changes over time within individuals (aging effects) from differences among people in their baseline levels (cohort effects).


What is Longitudinal Data?

- The defining characteristic of longitudinal studies is that individuals are **measured repeatedly through time**.
 - In contrast, cross-sectional studies involve collecting measurements on an individual **only once**.
- Concerned with **mean change** over time.
- Longitudinal studies allow examination of **age and cohort** effects.
 - Longitudinal studies can distinguish changes over time within individuals (aging effects) from differences among people in their baseline levels (cohort effects).


Age or cohort effects?



Age or cohort effects?



Age or cohort effects?



Benefits of Linear Mixed Models

- Fit intercepts and slopes for individuals
- Accounts for correlations between time points
- Accommodates missing data
- Accommodates measurements at unequal intervals
- Relatively few parameters required to account for dependency due to repeated measures
- Can handle both continuous and categorical covariates

Benefits of Linear Mixed Models

- Fit intercepts and slopes for individuals
- Accounts for correlations between time points
- Accommodates missing data
- Accommodates measurements at unequal intervals
- Relatively few parameters required to account for dependency due to repeated measures
- Can handle both continuous and categorical covariates

Benefits of Linear Mixed Models

- Fit intercepts and slopes for individuals
- Accounts for correlations between time points
- Accommodates missing data
- Accommodates measurements at unequal intervals
- Relatively few parameters required to account for dependency due to repeated measures
- Can handle both continuous and categorical covariates

Benefits of Linear Mixed Models

- Fit intercepts and slopes for individuals
- Accounts for correlations between time points
- Accommodates missing data
- Accommodates measurements at unequal intervals
- Relatively few parameters required to account for dependency due to repeated measures
- Can handle both continuous and categorical covariates

Benefits of Linear Mixed Models

- Fit intercepts and slopes for individuals
- Accounts for correlations between time points
- Accommodates missing data
- Accommodates measurements at unequal intervals
- Relatively few parameters required to account for dependency due to repeated measures
- Can handle both continuous and categorical covariates

Benefits of Linear Mixed Models

- Fit intercepts and slopes for individuals
- Accounts for correlations between time points
- Accommodates missing data
- Accommodates measurements at unequal intervals
- Relatively few parameters required to account for dependency due to repeated measures
- Can handle both continuous and categorical covariates

Correlations between time points

- Traditional regression does not account for this correlation
- This results in bogus standard error estimates and p -values.
- This is done in the covariance-variance matrix among the repeated measures known as Σ_i
- Recall a correlation matrix is a standardized covariance-variance matrix.
- This matrix is involved directly in estimating the parameters unlike in traditional regression.
- This matrix is broken down further and includes the random effects matrix.

Correlations between time points

- Traditional regression does not account for this correlation
- This results in bogus standard error estimates and p -values.
- This is done in the covariance-variance matrix among the repeated measures known as Σ_i
- Recall a correlation matrix is a standardized covariance-variance matrix.
- This matrix is involved directly in estimating the parameters unlike in traditional regression.
- This matrix is broken down further and includes the random effects matrix.

Correlations between time points

- Traditional regression does not account for this correlation
- This results in bogus standard error estimates and p -values.
- This is done in the covariance-variance matrix among the repeated measures known as Σ_i
- Recall a correlation matrix is a standardized covariance-variance matrix.
- This matrix is involved directly in estimating the parameters unlike in traditional regression.
- This matrix is broken down further and includes the random effects matrix.

Correlations between time points

- Traditional regression does not account for this correlation
- This results in bogus standard error estimates and p -values.
- This is done in the covariance-variance matrix among the repeated measures known as Σ_i
- Recall a correlation matrix is a standardized covariance-variance matrix.
- This matrix is involved directly in estimating the parameters unlike in traditional regression.
- This matrix is broken down further and includes the random effects matrix.

Correlations between time points

- Traditional regression does not account for this correlation
- This results in bogus standard error estimates and p -values.
- This is done in the covariance-variance matrix among the repeated measures known as Σ_i
- Recall a correlation matrix is a standardized covariance-variance matrix.
- This matrix is involved directly in estimating the parameters unlike in traditional regression.
- This matrix is broken down further and includes the random effects matrix.

Correlations between time points

- Traditional regression does not account for this correlation
- This results in bogus standard error estimates and p -values.
- This is done in the covariance-variance matrix among the repeated measures known as Σ_i
- Recall a correlation matrix is a standardized covariance-variance matrix.
- This matrix is involved directly in estimating the parameters unlike in traditional regression.
- This matrix is broken down further and includes the random effects matrix.

Accommodates missing data

- Missing data handled through robust maximum likelihood estimation
- No need for multiple imputation when using continuous variables
- Two types of Maximum Likelihood Estimates
 - Full Maximum Likelihood
 - Restricted Maximum Likelihood - provides unbiased variance estimates

Accommodates missing data

- Missing data handled through robust maximum likelihood estimation
- No need for multiple imputation when using continuous variables
- Two types of Maximum Likelihood Estimates
 - Full Maximum Likelihood
 - Restricted Maximum Likelihood - provides unbiased variance estimates

Accommodates missing data

- Missing data handled through robust maximum likelihood estimation
- No need for multiple imputation when using continuous variables
- Two types of Maximum Likelihood Estimates
 - Full Maximum Likelihood
 - Restricted Maximum Likelihood - provides unbiased variance estimates

Accommodates missing data

- Missing data handled through robust maximum likelihood estimation
- No need for multiple imputation when using continuous variables
- Two types of Maximum Likelihood Estimates
 - Full Maximum Likelihood
 - Restricted Maximum Likelihood - provides unbiased variance estimates

Accommodates missing data

- Missing data handled through robust maximum likelihood estimation
- No need for multiple imputation when using continuous variables
- Two types of Maximum Likelihood Estimates
 - Full Maximum Likelihood
 - Restricted Maximum Likelihood - provides unbiased variance estimates

Other benefits

- Individuals can be measured at different times and do not need to be measured at all times.
 - Therefore, we do not need to throw out data or impute data on individuals.
 - We throw out **cases not individuals**.
- Fewer parameters used in these models than ANOVA and MANOVA models. This leads to higher statistical power.
- Can handle continuous and categorical covariates simultaneously. No need for a separate model.
 - Categorical covariates are dummy coded.

Other benefits

- Individuals can be measured at different times and do not need to be measured at all times.
 - Therefore, we do not need to throw out data or impute data on individuals.
 - We throw out **cases not individuals**.
- Fewer parameters used in these models than ANOVA and MANOVA models. This leads to higher statistical power.
- Can handle continuous and categorical covariates simultaneously. No need for a separate model.
 - Categorical covariates are dummy coded.

Other benefits

- Individuals can be measured at different times and do not need to be measured at all times.
 - Therefore, we do not need to throw out data or impute data on individuals.
 - We throw out **cases not individuals**.
- Fewer parameters used in these models than ANOVA and MANOVA models. This leads to higher statistical power.
- Can handle continuous and categorical covariates simultaneously. No need for a separate model.
 - Categorical covariates are dummy coded.

Other benefits

- Individuals can be measured at different times and do not need to be measured at all times.
 - Therefore, we do not need to throw out data or impute data on individuals.
 - We throw out **cases not individuals**.
- Fewer parameters used in these models than ANOVA and MANOVA models. This leads to higher statistical power.
- Can handle continuous and categorical covariates simultaneously. No need for a separate model.
 - Categorical covariates are dummy coded.

Other benefits

- Individuals can be measured at different times and do not need to be measured at all times.
 - Therefore, we do not need to throw out data or impute data on individuals.
 - We throw out **cases not individuals**.
- Fewer parameters used in these models than ANOVA and MANOVA models. This leads to higher statistical power.
- Can handle continuous and categorical covariates simultaneously. No need for a separate model.
 - Categorical covariates are dummy coded.

Other benefits

- Individuals can be measured at different times and do not need to be measured at all times.
 - Therefore, we do not need to throw out data or impute data on individuals.
 - We throw out **cases not individuals**.
- Fewer parameters used in these models than ANOVA and MANOVA models. This leads to higher statistical power.
- Can handle continuous and categorical covariates simultaneously. No need for a separate model.
 - Categorical covariates are dummy coded.

Other names for these models

- Hierarchical linear models (HLM)
 - HLM are generally cross-sectional and involve multiple levels, such as school effects
 - In longitudinal models, measurements over time are **nested** within individuals rather than a school
- Linear Mixed Models
- Multilevel Models
- Mixed Effects Models

Other names for these models

- Hierarchical linear models (HLM)
 - HLM are generally cross-sectional and involve multiple levels, such as school effects
 - In longitudinal models, measurements over time are **nested** within individuals rather than a school
- Linear Mixed Models
- Multilevel Models
- Mixed Effects Models

Other names for these models

- Hierarchical linear models (HLM)
 - HLM are generally cross-sectional and involve multiple levels, such as school effects
 - In longitudinal models, measurements over time are **nested** within individuals rather than a school
- Linear Mixed Models
- Multilevel Models
- Mixed Effects Models

Other names for these models

- Hierarchical linear models (HLM)
 - HLM are generally cross-sectional and involve multiple levels, such as school effects
 - In longitudinal models, measurements over time are **nested** within individuals rather than a school
- Linear Mixed Models
- Multilevel Models
- Mixed Effects Models

Other names for these models

- Hierarchical linear models (HLM)
 - HLM are generally cross-sectional and involve multiple levels, such as school effects
 - In longitudinal models, measurements over time are **nested** within individuals rather than a school
- Linear Mixed Models
- Multilevel Models
- Mixed Effects Models

Other names for these models

- Hierarchical linear models (HLM)
 - HLM are generally cross-sectional and involve multiple levels, such as school effects
 - In longitudinal models, measurements over time are **nested** within individuals rather than a school
- Linear Mixed Models
- Multilevel Models
- Mixed Effects Models

LMM formula

Recall a simple multiple regression model is:

$$Y_i = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \dots + \beta_p X_{pi} + e_i$$

In a LMM, a regression model measured j times might become:

$$Y_{ij} = \beta_0 + b_{0i} + \beta_1 X_{1i} + (\beta_2 + b_{2i})t_{ij} + e_{ij}$$

LMM formula explained

Nomenclature	Definition
Y_{ij}	refers to the outcome variable, e.g. infant's weight, for individual i at time j
β_0	refers to the intercept
b_{0i}	refers to the random intercept for individual i . This allows individuals to vary in their initial values.
$\beta_1 X_{1i}$	the parameter estimate associated with a covariate, such as primary feeding type.
β_2	the fixed effect estimate associated with time
b_{2i}	refers to the random slope for individual i . This allows individuals to vary in their slopes.
t_{ij}	time variable, months since birth.
e_{ij}	is the residual error term.

Assumptions of LMMs

- Observations of individuals are independent
- Random effects are normally distributed
- Random residuals are normally distributed

Assumptions of LMMs

- Observations of individuals are independent
- Random effects are normally distributed
- Random residuals are normally distributed

Assumptions of LMMs

- Observations of individuals are independent
- Random effects are normally distributed
- Random residuals are normally distributed

Running LMMs

$$Y_{ij} = \beta_0 + b_{0i} + (\beta_1 + b_{1i})t_{ij} + e_{ij}$$

- Use the `{lme4}` package in **R** and the `lmer` function.
- If you can write your model as a LMM then you can program it in **R**.
- Basic **R** Formula:

```
m1 <- lmer(Outcome ~ 1 + Time + (1 + month|ID), data = data))
```

Running LMMs

$$Y_{ij} = \beta_0 + b_{0i} + (\beta_1 + b_{1i})t_{ij} + e_{ij}$$

- Use the `{lme4}` package in **R** and the `lmer` function.
- If you can write your model as a LMM then you can program it in **R**.
- Basic **R** Formula:
`m1 <- lmer(Outcome ~ 1 + Time + (1 + month|ID), data = data))`

Running LMMs

$$Y_{ij} = \beta_0 + b_{0i} + (\beta_1 + b_{1i})t_{ij} + e_{ij}$$

- Use the `{lme4}` package in **R** and the `lmer` function.
- If you can write your model as a LMM then you can program it in **R**.
- Basic **R** Formula:
`m1 <- lmer(Outcome ~ 1 + Time + (1 + month|ID), data = data))`

Running LMMs

```
> library(lme4)
> data(sleepstudy)
> m1 <- lmer(Reaction ~ 1 + Days + (1 + Days|Subject), sleepstudy)
> summary(m1)
```

Linear mixed model fit by REML

Formula: Reaction ~ 1 + Days + (1 + Days | Subject)

Data: sleepstudy

	AIC	BIC	logLik	deviance	REMLdev
	1756	1775	-871.8	1752	1744

Random effects:

Groups	Name	Variance	Std.Dev.	Corr
Subject	(Intercept)	612.090	24.7405	
	Days	35.072	5.9221	0.066
Residual		654.941	25.5918	

Number of obs: 180, groups: Subject, 18

Fixed effects:

	Estimate	Std. Error	t value
(Intercept)	251.405	6.825	36.84
Days	10.467	1.546	6.77

Correlation of Fixed Effects:

(Intr)	
Days	-0.138

Getting Help with LMMs and R

- Help with LMMs

- Fitzmaurice, G. M., Laird, N.M., & Ware, J.H. (2004). *Applied Longitudinal Analysis*. Wiley-Interscience.
- Diggle, P., Heagerty, P, Liang, K-Y, & Zeger, S. (2002). *Analysis of Longitudinal Data*. Oxford University Press.
- Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.

- Help with **R** and {lme4}.

- General R help. www.r-project.org
- Nice intro to R. <http://cran.r-project.org/doc/contrib/Verzani-SimpleR.pdf>
- {lme4} help. Pinheiro, J. C. & Bates, D. (2002). *Mixed Effects Models in S and S-Plus*. Springer Press. CAN
DOWNLOAD FROM LIBRARY!

Getting Help with LMMs and R

- Help with LMMs

- Fitzmaurice, G. M., Laird, N.M., & Ware, J.H. (2004). *Applied Longitudinal Analysis*. Wiley-Interscience.
- Diggle, P., Heagerty, P., Liang, K-Y, & Zeger, S. (2002). *Analysis of Longitudinal Data*. Oxford University Press.
- Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.

- Help with **R** and {lme4}.

- General R help. www.r-project.org
- Nice intro to R. <http://cran.r-project.org/doc/contrib/Verzani-SimpleR.pdf>
- {lme4} help. Pinheiro, J. C. & Bates, D. (2002). *Mixed Effects Models in S and S-Plus*. Springer Press. CAN
DOWNLOAD FROM LIBRARY!

Getting Help with LMMs and R

- Help with LMMs

- Fitzmaurice, G. M., Laird, N.M., & Ware, J.H. (2004). *Applied Longitudinal Analysis*. Wiley-Interscience.
- Diggle, P., Heagerty, P, Liang, K-Y, & Zeger, S. (2002). *Analysis of Longitudinal Data*. Oxford University Press.
- Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.

- Help with **R** and {lme4}.

- General R help. www.r-project.org
- Nice intro to R. <http://cran.r-project.org/doc/contrib/Verzani-SimpleR.pdf>
- {lme4} help. Pinheiro, J. C. & Bates, D. (2002). *Mixed Effects Models in S and S-Plus*. Springer Press. CAN
DOWNLOAD FROM LIBRARY!

Getting Help with LMMs and R

- Help with LMMs

- Fitzmaurice, G. M., Laird, N.M., & Ware, J.H. (2004). *Applied Longitudinal Analysis*. Wiley-Interscience.
- Diggle, P., Heagerty, P, Liang, K-Y, & Zeger, S. (2002). *Analysis of Longitudinal Data*. Oxford University Press.
- Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.

- Help with R and {lme4}.

- General R help. www.r-project.org
- Nice intro to R. <http://cran.r-project.org/doc/contrib/Verzani-SimpleR.pdf>
- {lme4} help. Pinheiro, J. C. & Bates, D. (2002). *Mixed Effects Models in S and S-Plus*. Springer Press. CAN DOWNLOAD FROM LIBRARY!

Getting Help with LMMs and R

- Help with LMMs

- Fitzmaurice, G. M., Laird, N.M., & Ware, J.H. (2004). *Applied Longitudinal Analysis*. Wiley-Interscience.
- Diggle, P., Heagerty, P, Liang, K-Y, & Zeger, S. (2002). *Analysis of Longitudinal Data*. Oxford University Press.
- Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.

- Help with **R** and {lme4}.

- General **R** help. www.r-project.org
- Nice intro to **R**. <http://cran.r-project.org/doc/contrib/Verzani-SimpleR.pdf>
- {lme4} help. Pinheiro, J. C. & Bates, D. (2002). *Mixed Effects Models in S and S-Plus*. Springer Press. CAN
DOWNLOAD FROM LIBRARY!

Getting Help with LMMs and R

- Help with LMMs
 - Fitzmaurice, G. M., Laird, N.M., & Ware, J.H. (2004). *Applied Longitudinal Analysis*. Wiley-Interscience.
 - Diggle, P., Heagerty, P, Liang, K-Y, & Zeger, S. (2002). *Analysis of Longitudinal Data*. Oxford University Press.
 - Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.
- Help with **R** and {lme4}.
 - General **R** help. www.r-project.org
 - Nice intro to **R**. <http://cran.r-project.org/doc/contrib/Verzani-SimpleR.pdf>
 - {lme4} help. Pinheiro, J. C. & Bates, D. (2002). *Mixed Effects Models in S and S-Plus*. Springer Press. CAN
DOWNLOAD FROM LIBRARY!

Getting Help with LMMs and R

- Help with LMMs

- Fitzmaurice, G. M., Laird, N.M., & Ware, J.H. (2004). *Applied Longitudinal Analysis*. Wiley-Interscience.
- Diggle, P., Heagerty, P, Liang, K-Y, & Zeger, S. (2002). *Analysis of Longitudinal Data*. Oxford University Press.
- Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.

- Help with **R** and {lme4}.

- General **R** help. www.r-project.org
- Nice intro to **R**. <http://cran.r-project.org/doc/contrib/Verzani-SimpleR.pdf>
- {lme4} help. Pinheiro, J. C. & Bates, D. (2002). *Mixed Effects Models in S and S-Plus*. Springer Press. CAN
DOWNLOAD FROM LIBRARY!

Getting Help with LMMs and R

- Help with LMMs
 - Fitzmaurice, G. M., Laird, N.M., & Ware, J.H. (2004). *Applied Longitudinal Analysis*. Wiley-Interscience.
 - Diggle, P., Heagerty, P, Liang, K-Y, & Zeger, S. (2002). *Analysis of Longitudinal Data*. Oxford University Press.
 - Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge University Press.
- Help with **R** and {lme4}.
 - General **R** help. www.r-project.org
 - Nice intro to **R**. <http://cran.r-project.org/doc/contrib/Verzani-SimpleR.pdf>
 - {lme4} help. Pinheiro, J. C. & Bates, D. (2002). *Mixed Effects Models in S and S-Plus*. Springer Press. CAN DOWNLOAD FROM LIBRARY!