

Bill shape distinguishes Antarctic penguin species

The Day-1 result — one source, a branded Typst PDF

Christophe Dervieux
Posit, PBC

2026-08-08

Table of contents

1 Introduction	1
2 Species at a glance	1
3 Bill length versus depth	2
4 Conclusion	2
References	2
Bibliography	2

1 Introduction

The base-R `penguins` dataset records body measurements for 342 penguins of three species collected at Palmer Station, Antarctica (Gorman et al., 2014). It reached the R community first through the `palmerpenguins` package (Horst et al., 2020). Here we ask a simple question: **does bill shape tell the species apart?** We render the answer as a branded PDF with **Typst**, a modern typesetting system that ships inside Quarto, so there is no LaTeX involved.

2 Species at a glance

Table 1

Penguins of the Palmer Archipelago

Mean measurements per species

Species	N	Bill (mm)		Body mass (g)
		Length	Depth	
Adelie	151	38.8	18.3	3 701
Gentoo	123	47.5	15.0	5 076
Chinstrap	68	48.8	18.4	3 733

Gentoo is the heaviest on average, at 5 076 g, the `large-bodied` species of the three.

3 Bill length versus depth




Figure 1: Bill length versus depth by species. The single heaviest penguin is labeled.

The three species separate cleanly in bill shape. This is the manuscript-ready version of the same figure from the running HTML report, now in a branded PDF.

4 Conclusion

One `.qmd`, one `_brand.yml`, and `format: typst` give a citable, typeset document, with the `RaukR` palette carried through layout, tables, and figures.

References

Bibliography

- Gorman, K. B., Williams, T. D., & Fraser, W. R. (2014). Ecological Sexual Dimorphism and Environmental Variability within a Community of Antarctic Penguins (Genus *Pygoscelis*). *PLoS ONE*, 9(3), e90081. <https://doi.org/10.1371/journal.pone.0090081>
- Horst, A. M., Hill, A. P., & Gorman, K. B. (2020). *palmerpenguins: Palmer Archipelago (Antarctica) Penguin Data*. <https://doi.org/10.5281/zenodo.3960218>