

# 2019 ADA miniHW 1

b07902064 資工二 蔡銘軒

September 21, 2019

From the definition of  $g(n) = \Theta(F(n))$ , there exist positive constants  $c_1$ ,  $c_2$ , and  $n_0$ , such that  $0 \leq c_1 F(n) \leq g(n) \leq c_2 F(n)$ , for all  $n \geq n_0$ .

And from the definition of  $h(n) = o(F(n))$ , for every positive constant  $c > 0$ , there exists a constant  $n_1 > 0$ , such that  $0 \leq h(n) < cF(n)$ , for all  $n \geq n_1$ .

By taking  $n_2 = \max(n_0, n_1)$ , we have  $0 \leq h(n) < c_1 F(n) \leq g(n) \leq c_2 F(n)$ , for all  $n \geq n_2$ . Subtracting the inequalities by  $h(n)$ , we get  $0 < c_1 F(n) - h(n) \leq f(n) \leq c_2 F(n) - h(n)$ , for all  $n \geq n_2$ .

We now focus on the term  $c_1 F(n) - h(n)$ . Since  $h(n) < cF(n)$  for all  $c > 0$ , we have  $c_1 F(n) - cF(n) = (c_1 - c)F(n) < c_1 F(n) - h(n)$  for all  $c > 0$ . The term  $(c_1 - c)$  can take any value in the interval  $(-\infty, c_1)$ . Since  $c_1 > 0$ , we can choose a constant  $c_3$  such that  $c_3 \in (-\infty, c_1)$  and  $c_3 > 0$ .

Now we arrive at  $0 < c_3 F(n) < f(n)$ , which can also be written as  $0 \leq c_3 F(n) \leq f(n)$ , for all  $n \geq n_2 = \max(n_0, n_1)$  with a positive constant  $c_3 > 0$ , which is the definition of  $f(n) = \Omega(F(n))$ . The statement is thus proven.