


**TABLE 1.1 Fourier Series for Common Repetitive Waveforms**


## 1. Half-wave rectified sine wave



$$v(t) = \frac{V}{\pi} + \frac{V}{2} \sin \omega t - \frac{2V}{\pi} \left( \frac{\cos 2\omega t}{1 \times 3} + \frac{\cos 4\omega t}{3 \times 5} + \frac{\cos 6\omega t}{5 \times 7} + \dots \right)$$


## 2. Full-wave rectified sine wave

(a) With time zero at voltage zero



$$v(t) = \frac{2V}{\pi} - \frac{4V}{\pi} \left( \frac{\cos 2\omega t}{1 \times 3} + \frac{\cos 4\omega t}{3 \times 5} + \frac{\cos 6\omega t}{5 \times 7} + \dots \right)$$


(b) With time zero at voltage peak



$$v(t) = \frac{2V}{\pi} \left( 1 + \frac{2 \cos 2\omega t}{1 \times 3} - \frac{2 \cos 4\omega t}{3 \times 5} + \frac{2 \cos 6\omega t}{5 \times 7} - \dots \right)$$

## 3. Square wave

(a) Odd function




$$v(t) = \frac{4V}{\pi} \left( \sin \omega t + \frac{1}{3} \sin 3\omega t + \frac{1}{5} \sin 5\omega t + \dots \right)$$

(continues)


TABLE 1.1 (continued)

(b) Even function




$$v(t) = \frac{4V}{\pi} \left( \cos \omega t - \frac{1}{3} \cos 3\omega t + \frac{1}{5} \cos 5\omega t - \dots \right)$$

4. Pulse train



$$v(t) = \frac{V\tau}{T} + \frac{2V\tau}{T} \left( \frac{\sin \pi\tau/T}{\pi\tau/T} \cos \omega t + \frac{\sin 2\pi\tau/T}{2\pi\tau/T} \cos 2\omega t + \frac{\sin 3\pi\tau/T}{3\pi\tau/T} \cos 3\omega t + \dots \right)$$


5. Triangle wave



$$v(t) = \frac{8V}{\pi^2} \left( \cos \omega t + \frac{1}{3^2} \cos 3\omega t + \frac{1}{5^2} \cos 5\omega t + \dots \right)$$

6. Sawtooth wave

(a) With no dc offset




$$v(t) = \frac{2V}{\pi} \left( \sin \omega t - \frac{1}{2} \sin 2\omega t + \frac{1}{3} \sin 3\omega t - \dots \right)$$

(continues)

TABLE 1.1 (continued)

(b) Positive-going



$$v(t) = \frac{V}{2} - \frac{V}{\pi} \left( \sin \omega t + \frac{1}{2} \sin 2\omega t + \frac{1}{3} \sin 3\omega t + \dots \right)$$

**SOLUTION**

A square wave is another signal with a simple Fourier representation, although not quite as simple as for a sine wave. For the signal shown in Figure 1.6(a), the frequency is 1 kHz, as before, and the peak voltage is 1 V.

According to Table 1.1, this signal has components at an infinite number of frequencies: all odd multiples of the fundamental frequency of 1 kHz. However, the amplitude decreases with frequency, so that the third harmonic has an amplitude one-third that of the fundamental, the fifth harmonic an amplitude of one-fifth that of the fundamental, and so on. Mathematically, a square wave of voltage with a rising edge at  $t = 0$  and no dc offset can be expressed as follows (see Table 1.1):

$$v(t) = \frac{4V}{\pi} \left( \sin \omega t + \frac{1}{3} \sin 3\omega t + \frac{1}{5} \sin 5\omega t + \dots \right)$$

where

$V$  = peak amplitude of the square wave

$\omega$  = radian frequency of the square wave

$t$  = time in seconds

For this example, the above equation becomes:

$$v(t) = \frac{4}{\pi} \left( \sin(2\pi \times 10^3 t) + \frac{1}{3} \sin(6\pi \times 10^3 t) + \frac{1}{5} \sin(10\pi \times 10^3 t) + \dots \right) V$$

This equation shows that the signal has frequency components at odd multiples of 1 kHz, that is, at 1 kHz, 3 kHz, 5 kHz, and so on. The 1-kHz component has a peak amplitude of

$$V_1 = \frac{4}{\pi} = 1.27 \text{ V}$$