




Math League Contest ~ Fall 2010 ~ Solutions

1.	$f(i, 2010) = \sum_{k=0}^{2010} i^k = i^0 + i^1 + i^2 + i^3 + i^4 + \dots + i^{2007} + i^{2008} + i^{2009} + i^{2010}$ $= \underbrace{(1+i-1-i)}_{k=0,1,2,3} + \underbrace{(1+i-1-i)}_{k=4,5,6,7} + \dots + \underbrace{(1+i-1-i)}_{k=2004,2005,2006,2007} + 1+i-1 = \underbrace{0+0+\dots+0}_{502 \text{ zeros}} + 1+i-1 = i$ <p>Or use the formula for a geometric series, <math>\sum_{k=0}^n r^k = \frac{r^{n+1}-1}{r-1}</math> with <math>r=i</math> and <math>n=2010</math>. <span style="border: 1px solid black; padding: 2px;">Answer: i</span></p>
2.	<p>If we let <math>y = \sqrt[4]{2010x+1}</math>, then <math>\sqrt{2010x+1} = y^2</math>. Now the equation takes on a simpler form: <math>y^2 - y = 2</math>. Solving for <math>y</math>: <math>y^2 - y - 2 = 0 \Rightarrow (y+1)(y-2) = 0 \Rightarrow y = -1</math> or <math>y = 2</math>. However, <math>y</math> must be positive. Thus, <math>y = 2 \Rightarrow \sqrt{2010x+1} = 2^2 \Rightarrow 2010x+1 = 4^2 \Rightarrow 2010x = 15</math>.</p> <p>Therefore, <math>x = \frac{15}{2010} = \frac{1}{134}</math>. <span style="border: 1px solid black; padding: 2px;">Answer: <math>\frac{15}{2010}</math> or <math>\frac{1}{134}</math></span></p>
3.	<p>Let <math>x = \sqrt{7+\sqrt{48}} + \sqrt{7-\sqrt{48}}</math>, which is certainly greater than zero. Squaring gives:</p> $x^2 = (\sqrt{7+\sqrt{48}} + \sqrt{7-\sqrt{48}})^2 \Rightarrow x^2 = (\sqrt{7+\sqrt{48}})^2 + 2(\sqrt{7+\sqrt{48}})(\sqrt{7-\sqrt{48}}) + (\sqrt{7-\sqrt{48}})^2$ $= 7 + \sqrt{48} + 2\sqrt{49-48} + 7 - \sqrt{48} = 14 + 2\sqrt{1} = 16$ <p>Taking the positive root, <math>x = \sqrt{16}</math>. <span style="border: 1px solid black; padding: 2px;">Answer: c</span></p>
4.	<p>Rich would have 4 pennies, 4 dimes, 1 quarter and 1 half-dollar. <span style="border: 1px solid black; padding: 2px;">Answer: \$1.19</span></p>
5.	<p>Let <math>x</math> be the length of the straight segment between two circles, and let <math>y</math> be the length of the arc from one straight segment to the other. <math>x = 2 \cdot \text{radius} = 2 \cdot 1 = 2</math> ft, and <math>y = (\frac{1}{3}) \cdot \text{circumference} = (\frac{1}{3}) \cdot 2\pi = (\frac{2}{3})\pi</math> ft. Thus, the total length <math>= 3x + 3y = 3 \cdot 2 + 3 \cdot (\frac{2}{3})\pi = 6 + 2\pi</math> ft.</p> <p style="text-align: center;"><span style="border: 1px solid black; padding: 2px;">Answer: <math>6 + 2\pi</math> ft</span></p> 
6.	$x^4 - y^4 = x^2 - y^2 \Rightarrow (x^4 - y^4) - (x^2 - y^2) = 0 \Rightarrow (x^2 - y^2)(x^2 + y^2) - (x^2 - y^2) = 0$ $\Rightarrow (x^2 - y^2)[(x^2 + y^2) - 1] = 0 \Rightarrow x^2 - y^2 = 0 \text{ or } x^2 + y^2 - 1 = 0$ $\Rightarrow y^2 = x^2 \Rightarrow y = \pm x \text{ or } x^2 + y^2 = 1, \text{ which gives two lines through the origin with slopes of } -1 \text{ and } 1, \text{ and a circle of radius } 1 \text{ centered at the origin.}$ <span style="border: 1px solid black; padding: 2px;">Answer: d</span>
7.	<p><math>a^2 - b^4 = 345 \Rightarrow (a-b^2)(a+b^2) = 3 \cdot 5 \cdot 23</math>, grouping the three factors of 345 in pairs give <math>(3 \cdot 5) \cdot 23</math>, <math>3 \cdot (5 \cdot 23)</math>, and <math>5 \cdot (3 \cdot 23) \Rightarrow 15 \cdot 23</math>, <math>3 \cdot 115</math>, and <math>5 \cdot 69</math>. Thus, either</p> <p>① <math>a-b^2=15</math> and <math>a+b^2=23</math>, ② <math>a-b^2=3</math> and <math>a+b^2=115</math>, or ③ <math>a-b^2=5</math> and <math>a+b^2=69</math></p> <p>Only equations ① yield an integer for both <math>a</math> and <math>b</math>, with <math>a=19</math> and <math>b=2</math>. <span style="border: 1px solid black; padding: 2px;">Answer: <math>a+b=21</math></span></p>

8.	 <p>Letting <math>a</math> be the length of one side of the square, we get:  <math>a^2 = 3^2 + 7^2 \Rightarrow a^2 = 9 + 49 = 58 = \text{Area of the Square}</math></p>	Answer: 58
9.	 <p>The area of the shaded region is: <math>\pi R^2 - \pi r^2 = \pi(R^2 - r^2)</math>.  From the diagram, we get the relation: <math>R^2 = r^2 + 1 \Rightarrow R^2 - r^2 = 1</math>.  Thus, the area of the shaded region is <math>\pi \cdot 1 = \pi</math>.</p>	Answer: $\pi$
10.	Three tickets can be selected in $3!$ ways, or 6 ways. Only 1 of the 6 permutations would have the tickets in increasing order.	Answer: $\frac{1}{6}$
11.	Since the maximum value of $\sin(x)$ is 1, the line will not intersect the sine curve for $x > 100\pi$ . Thus, from $x = 0$ to $x = 100\pi$ , there will be 2 points of intersection per period. The period of $y = \sin(x)$ is $2\pi$ , which means there are 50 periods from $x = 0$ to $x = 100\pi$ . Hence, there will be $50 \cdot 2 = 100$ points of intersection from $x = 0$ to $x = 100\pi$ . Similarly from $x = -100\pi$ to $x = 0$ , but that would count the origin again. Therefore, there are $100 + 100 - 1$ points of intersection.	Answer: 199
12.	If there is only one such number between 2 and $2^{100}$ , then it must be the smallest. Hence, it must be $2^n$ , where $n$ is the least common multiple of 2, 3, 4, 5 and 6 (so the 2 <sup>nd</sup> , 3 <sup>rd</sup> , 4 <sup>th</sup> , 5 <sup>th</sup> and 6 <sup>th</sup> roots are whole numbers). Thus, $n = 3 \cdot 4 \cdot 5 = 60$ and the number is $2^{60}$ . <u>Note:</u> The next such number is $2^{2 \cdot 60} = 2^{120} > 2^{100}$ .	Answer: $2^{60}$
13.	$\log_{2011}(2010) - \frac{1}{\log_{2010}(2011)} = \frac{\log(2010)}{\log(2011)} - \frac{1}{\left(\frac{\log(2011)}{\log(2010)}\right)} = \frac{\log(2010)}{\log(2011)} - \frac{\log(2010)}{\log(2011)} = 0$	Answer: b
14.	Ray works at the rate of $\frac{1}{9}$ of a job/hr. Let $r$ be the rate at which Tim works. Together, they work at $\left(\frac{1}{9} + r\right)$ job/hr, and since Work = Rate $\cdot$ Time, after 4 hours they complete $4\left(\frac{1}{9} + r\right)$ of the job. However, it takes Ray 2 more hours to finish. Thus, there was only $2\left(\frac{1}{9}\right) = \frac{2}{9}$ of the job remaining. Therefore, together they must have completed $\frac{7}{9}$ of the job. From this we get: $4\left(\frac{1}{9} + r\right) = \frac{7}{9}$ . Solving for $r$ , gives $r = \frac{1}{12}$ job/hr. Hence, working alone, Tim would need 12 hours.	Answer: 12 hours
15.	Let $p$ = the probability of obtaining <i>heads up</i> on one toss of the coin. Then $1 - p$ = the probability of obtaining <i>tails up</i> on one toss of the coin. The probability of getting two heads on two tosses is $p^2$ , which is equal to $1 - p$ . Thus, $p^2 = 1 - p \Rightarrow p^2 + p - 1 = 0 \Rightarrow p = \frac{-1 \pm \sqrt{1^2 - 4(1)(-1)}}{2(1)} = \frac{-1 \pm \sqrt{5}}{2}$ . However, only $\frac{-1 + \sqrt{5}}{2}$ is positive (and between 0 and 1).	Answer: $\frac{\sqrt{5} - 1}{2}$

16.	I must pick up 7 shirts to hold me over until the following Monday. Hence, I must drop off 7 shirts each Monday. Counting the shirt I wear on Monday, the required total is $7+7+1=15$ . Note: I cannot get by with only 14 shirts, as I would not have a clean shirt to wear the following Monday. <b>Answer: d</b>
17.	When Sophia completed half the race, Ida was 3.5 km ahead. Thus, by the time Sophia completed the entire race, Ida would have been 7 km ahead (if she continued at the same rate). Hence, Sophia completed the race at 11:52 AM plus the time it would have taken Ida to go an additional 7 km. At 42 km/hr, it would have taken Ida $(7 \div 42)$ hour = $\frac{1}{6}$ hour = 10 minutes. Therefore, Sophia completed the race at 12:02 PM. <b>Answer: 12:02 PM</b>
18.	Since each match eliminates one competitor, and 2009 competitors must be eliminated so that only one person (the winner) remains, 2009 matches must be played. <b>Answer: b</b>
19.	Let $w$ represent the hourly wage. Then the tax rate is $(2w)\%$ and the tax on $w$ will be $\frac{2w}{100} \cdot w = \frac{w^2}{50}$ . Thus, the after taxes income (per hour) is $w - \frac{w^2}{50}$ . This quadratic is maximized along the axis of symmetry, i.e. for $w = \frac{-b}{2a} = \frac{(-1)}{2\left(-\frac{1}{50}\right)} = 25$ . <b>Answer: \$25 per hour</b>
20.	Look at each clue, knowing <i>exactly</i> one of each person's statements is true. Artie's: If <i>it was Barbara</i> is true, then we know the other statement is false, therefore it was Edward. This is a contradiction. Hence we now know it wasn't Barbara, nor Edward (as <i>it wasn't Edward</i> must be the true statement). Looking at Carmine's statements, we can similarly determine that it wasn't Artie. Since we know it wasn't Barbara, Darci's statements tell us <i>it was Carmine</i> is true. This also checks against the other clues. <b>Answer: Carmine</b>

