

Newton-Raphson Method for Convex Optimization

Conner DiPaolo, Jeffrey Rutledge, Colin Adams

Harvey Mudd College

February, 2016

Finding Roots of Functions

Given a differentiable function $f : \mathbb{R} \mapsto \mathbb{R}$ we want to find the instances when $f(x) = 0$ (not generally solvable in closed form).


Newton's Method:

1. Take a starting position x_0
2. Find where the tangent line $y = f(x_n) + (x_{n+1} - x_n)f'(x_n)$ is 0 and iterate:

$$y = f(x_n) + (x_{n+1} - x_n)f'(x_n) = 0$$
$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Finding Roots of Functions


Initial $x_0 = 0.4$, $f(x) = (x - 1)(x - 3)^2 + e^{\frac{1}{3x}} - \cos\left(\frac{x}{2}\right) - 1.5$



$$x_0 = 0.4 \quad \Delta = 0.9805$$

Finding Roots of Functions


Initial $x_0 = 0.4$, $f(x) = (x - 1)(x - 3)^2 + e^{\frac{1}{3x}} - \cos\left(\frac{x}{2}\right) - 1.5$



$$x_1 = 0.6866 \quad \Delta = 0.6938$$

Finding Roots of Functions


Initial $x_0 = 0.4$, $f(x) = (x - 1)(x - 3)^2 + e^{\frac{1}{3x}} - \cos\left(\frac{x}{2}\right) - 1.5$



$$x_2 = 0.9938 \quad \Delta = 0.3866$$

Finding Roots of Functions

Initial $x_0 = 0.4$, $f(x) = (x - 1)(x - 3)^2 + e^{\frac{1}{3x}} - \cos\left(\frac{x}{2}\right) - 1.5$



$$x_3 = 1.2050 \quad \Delta = 0.1754$$

Converges in 28 iterations.

Optimizing Function Using Newton's Method

Instead of finding where $f(x) = 0$, how can we find where $f'(x) = 0$?

Optimizing Function Using Newton's Method

Instead of finding where $f(x) = 0$, how can we find where $f'(x) = 0$?

Instead of:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

Use derivatives:

$$x_{n+1} = x_n - \frac{f'(x_n)}{f''(x_n)}$$

Optimizing $f : \mathbb{R}^n \mapsto \mathbb{R}$

How do we find the derivatives of functions where $f : \mathbb{R}^n \mapsto \mathbb{R}$?

Optimizing $f : \mathbb{R}^n \mapsto \mathbb{R}$

How do we find the derivatives of functions where $f : \mathbb{R}^n \mapsto \mathbb{R}$?

$$x_{n+1} = x_n - \frac{f'(x_n)}{f''(x_n)}$$

becomes,

$$x_{n+1} = x_n - H_f^{-1}(x_n) \nabla f(x_n)$$

Optimizing $f : \mathbb{R}^n \mapsto \mathbb{R}$

How do we find the derivatives of functions where $f : \mathbb{R}^n \mapsto \mathbb{R}$?

$$x_{n+1} = x_n - \frac{f'(x_n)}{f''(x_n)}$$

becomes,

$$x_{n+1} = x_n - H_f^{-1}(x_n) \nabla f(x_n)$$

Def: The gradient of $f : \mathbb{R}^n \mapsto \mathbb{R}$ is

$$\nabla f = \begin{bmatrix} \frac{\partial}{\partial x_1} f & \frac{\partial}{\partial x_2} f & \dots & \frac{\partial}{\partial x_n} f \end{bmatrix}$$

Optimizing $f : \mathbb{R}^n \mapsto \mathbb{R}$

How do we find the derivatives of functions where $f : \mathbb{R}^n \mapsto \mathbb{R}$?

$$x_{n+1} = x_n - \frac{f'(x_n)}{f''(x_n)}$$

becomes,

$$x_{n+1} = x_n - H_f^{-1}(x_n) \nabla f(x_n)$$

Def: The gradient of $f : \mathbb{R}^n \mapsto \mathbb{R}$ is



$$\nabla f = \begin{bmatrix} \frac{\partial}{\partial x_1} f & \frac{\partial}{\partial x_2} f & \cdots & \frac{\partial}{\partial x_n} f \end{bmatrix}$$

Def: The Hessian of $f : \mathbb{R}^n \mapsto \mathbb{R}$ is

$$H_f = \begin{bmatrix} \frac{\partial^2 f}{\partial x_1 \partial x_1} & \frac{\partial^2 f}{\partial x_1 \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_1 \partial x_n} \\ \frac{\partial^2 f}{\partial x_2 \partial x_1} & \ddots & & \frac{\partial^2 f}{\partial x_2 \partial x_n} \\ \vdots & & \ddots & \vdots \\ \frac{\partial^2 f}{\partial x_n \partial x_1} & \frac{\partial^2 f}{\partial x_n \partial x_2} & \cdots & \frac{\partial^2 f}{\partial x_n \partial x_n} \end{bmatrix}$$



Optimizing $f : \mathbb{R}^n \mapsto \mathbb{R}$

Initial $x_0 = [1 \quad 0.6]$, $f(x) = -\cos(x^2 + y^2) - e^{-(x^2+y^2)}$



Optimizing $f : \mathbb{R}^n \mapsto \mathbb{R}$

Initial $x_0 = [1 \ 0.6]$, $f(x) = -\cos(x^2 + y^2) - e^{-(x^2+y^2)}$





$$x_0 = [1 \ 0.6]$$

$$\Delta = 1.1661$$

Optimizing $f : \mathbb{R}^n \mapsto \mathbb{R}$

Initial $x_0 = [1 \quad 0.6]$, $f(x) = -\cos(x^2 + y^2) - e^{-(x^2+y^2)}$





$$x_1 = [-0.1167 \quad -0.0700]$$

$$\Delta = 0.1361$$

Optimizing $f : \mathbb{R}^n \mapsto \mathbb{R}$

Initial $x_0 = [1 \quad 0.6]$, $f(x) = -\cos(x^2 + y^2) - e^{-(x^2+y^2)}$



$$x_2 = [-7.8 \times 10^{-5} \quad -4.7 \times 10^{-5}]$$

$$\Delta = 9.15 \times 10^{-5}$$

Converges in 4 iterations.