

Integrating over an Infinite Interval
○○○○●○○○○○○○○

Integrating a Discontinuous Function
○○○○○○○○○

Comparison Theorem
○○○○○○○○○

Solution

Then we have

$$A = \int_1^{\infty} \frac{1}{x} dx$$

$= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx$ Rewrite the improper integral as a limit.

$= \lim_{t \rightarrow \infty} \ln|x| \Big|_1^t$ Find the antiderivative.

$= \lim_{t \rightarrow \infty} (\ln|t| - \ln(1))$ Evaluate the antiderivative.

$= \infty$ Evaluate the limit.

Since the improper integral diverges to ∞ , the area of the region is infinite.

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

8 / 31

◀ ◻ ▶ ◀ ⏪ ▶ ◀ ⏩ ▶ ◀ ⏴ ▶ ◀ ⏵ ▶ ◀ ⏶ ▶ ◀ ⏷ ▶ ⌂ 🔍 ↺

Probability Theory:

- If accidents occur at a rate of one every 3 months, then the probability that the time between accidents is between a and b is given by

$$P(a \leq x \leq b) = \int_a^b f(x) \, dx,$$

where

$$f(x) = \begin{cases} 0 & \text{if } x < 0 \\ 3e^{-3x} & \text{if } x \geq 0 \end{cases}$$

Chapter Opener: Traffic Accidents in a City

Solution:

- We need to calculate the probability as an improper integral:

$$P(X \geq 8) = \int_8^{\infty} 3e^{-3x} dx$$

$$= \lim_{t \rightarrow \infty} \int_8^t 3e^{-3x} dx$$

$$= \lim_{t \rightarrow \infty} (-e^{-3x})$$

$$= \lim_{t \rightarrow \infty} (-e^{-3t} + e^{-24})$$

$$= e^{-24} \approx 3.8 \times 10^{-11}.$$

- The value 3.8×10^{-11} represents the probability of no accidents in 8

Integrating over an Infinite Interval
○○○○○○○○●○○○○

Integrating a Discontinuous Function
○○○○○○○○○

Comparison Theorem
○○○○○○○○○

Evaluating an Improper Integral over an Infinite Interval

Evaluate $\int_{-\infty}^0 \frac{1}{x^2+4} dx$. State whether the improper integral converges or diverges.

$$\int_{-\infty}^0 \frac{1}{x^2+4} dx = \lim_{t \rightarrow -\infty} \int_t^0 \frac{1}{x^2+4} dx$$

Rewrite as a limit.

$$= \lim_{t \rightarrow -\infty} \frac{1}{2} \tan^{-1} \left(\frac{x}{2} \right) \Big|_t^0$$

Find the antiderivative.

$$= \frac{1}{2} \lim_{t \rightarrow -\infty} \left(\tan^{-1}(0) - \tan^{-1} \left(\frac{t}{2} \right) \right)$$

Evaluate the antiderivative.

$$= \frac{\pi}{4}.$$

Evaluate the limit and simplify.

● The improper integral converges to $\frac{\pi}{4}$.

Math 1700 (University of Manitoba)3.7 Improper IntegralsMarch 15, 202413 / 31

Evaluate $\int_{-3}^{\infty} e^{-x} dx$. State whether the improper integral converges or diverges.

Answer: e^3 , converges

Hint: $\int_{-3}^{\infty} e^{-x} dx = \lim_{t \rightarrow \infty} \int_{-3}^t e^{-x} dx$

Integrating over an Infinite Interval

○○○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function

●○○○○○○○○

Comparison Theorem

○○○○○○○○

Understanding Integrals with Infinite Discontinuities

In mathematical analysis, the concept of integration is vital for understanding the accumulation of quantities over intervals. However, when dealing with functions that exhibit infinite discontinuities within the interval of integration, a nuanced approach is required.


● Consider an integral of the form: $\int_a^b f(x) \, dx$

● $f(x)$ is continuous over $[a, b]$ but discontinuous at b .

● Let's examine the behavior of this integral as t , the upper limit of integration, approaches b .

● Since $f(x)$ remains continuous over $[a, t]$ for all t satisfying $a < t < b$, the integral $\int_a^t f(x) \, dx$ is well-defined for such values of t .

● We define: $\int_a^b f(x) \, dx = \lim_{t \rightarrow b^-} \int_a^t f(x) \, dx$ provided this limit exists.



Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

17 / 31

Integrating over an Infinite Interval
○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function
○○○○○○○○○○

Comparison Theorem
○○○○○○○○○○

Definition

Let $f(x)$ be continuous over $[a, b)$. Then,

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx.$$

Let $f(x)$ be continuous over $(a, b]$. Then,

$$\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx.$$

If the limit exists, then the improper integral is said to converge. If the limit does not exist, then the improper integral is said to diverge.

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

18 / 31

◀ ◻ ▶ ⏪ ⏩ 🔍 ↺

Integrating over an Infinite Interval
○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function
○○○○○○○○○

Comparison Theorem
○○○○○○○○○

Definition

If $f(x)$ is continuous over $[a, b]$ except at a point c in (a, b) , then we define $\int_a^b f(x) \, dx$ as

$$\int_a^b f(x) \, dx = \int_a^c f(x) \, dx + \int_c^b f(x) \, dx,$$

provided both $\int_a^c f(x) \, dx$ and $\int_c^b f(x) \, dx$ converge. If either of these two integrals diverges, then $\int_a^b f(x) \, dx$ diverges.

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

19 / 31

◀ ◻ ▶ ⏪ ⏩ 🔍 ↺

Integrating over an Infinite Interval
○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function
○○●○○○○○

Comparison Theorem
○○○○○○○○○

Integrating a Discontinuous Integrand

Evaluate $\int_0^4 \frac{1}{\sqrt{4-x}} dx$, if possible. State whether the integral converges or diverges.

Solution: The function $f(x) = \frac{1}{\sqrt{4-x}}$ is continuous over $[0, 4)$ and discontinuous at 4. Using the above definition, we rewrite $\int_0^4 \frac{1}{\sqrt{4-x}} dx$ as a:

$\int_0^4 \frac{1}{\sqrt{4-x}} dx = \lim_{t \rightarrow 4^-} \int_0^t \frac{1}{\sqrt{4-x}} dx$

Rewrite as a limit.

$= \lim_{t \rightarrow 4^-} (-2\sqrt{4-x}) \Big|_0^t$

Find the antiderivative.

$= \lim_{t \rightarrow 4^-} (-2\sqrt{4-t} + 4)$

Evaluate the antiderivative.

$= 4.$

Evaluate the limit.

Because the limit exists, the improper integral converges.

Math 1700 (University of Manitoba)
3.7 Improper Integrals
March 15, 2024
20 / 31

Integrating over an Infinite Interval
○○○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function
○○○○●○○○

Comparison Theorem
○○○○○○○○

Integrating a Discontinuous Integrand

Evaluate $\int_0^2 x \ln(x) \, dx$. State whether the integral converges or diverges.

Solution: Since $f(x) = x \ln(x)$ is continuous over $(0, 2]$ and is discontinuous at zero, we can rewrite :

$$\int_0^2 x \ln(x) \, dx = \lim_{t \rightarrow 0^+} \int_t^2 x \ln(x) \, dx$$

Rewrite as a limit.

$$= \lim_{t \rightarrow 0^+} \left(\frac{1}{2} x^2 \ln(x) - \frac{1}{4} x^2 \right) \Big|_t^2$$

Evaluate using integr

$$= \lim_{t \rightarrow 0^+} \left(2 \ln(2) - 1 - \frac{1}{2} t^2 \ln(t) + \frac{1}{4} t^2 \right).$$

Evaluate the antideri

$$= 2 \ln(2) - 1.$$

The improper integral converges.

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

21 / 31

Integrating over an Infinite Interval
○○○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function
○○○○○○○○○●○○

Comparison Theorem
○○○○○○○○○

Integrating a Discontinuous Integrand

Evaluate $\int_{-1}^1 \frac{1}{x^3} dx$. State whether the improper integral converges or diverges.

Solution: Since $f(x) = \frac{1}{x^3}$ is continuous at every point of $[-1, 1]$ except zero, we use the corresponding definition to write
$$\int_{-1}^1 \frac{1}{x^3} dx = \int_{-1}^0 \frac{1}{x^3} dx + \int_0^1 \frac{1}{x^3} dx.$$

Our integral converges if both integrals on the right converge. If either of the two integrals on the right diverges, then the original integral diverges as well. Begin with $\int_{-1}^0 \frac{1}{x^3} dx$:

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

22 / 31

◀ ◻ ▶ ◀ ⏮ ▶ ▶ ⏭ ▶ ▶ ⏪ ▶ ▶ ⏩ ▶ ▶ ⏫ ▶ ▶ 🔍 ↺

Integrating over an Infinite Interval

○○○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function

○○○○○○○○●○

Comparison Theorem

○○○○○○○○○

Solution

$$\int_{-1}^0 \frac{1}{x^3} dx = \lim_{t \rightarrow 0^-} \int_{-1}^t \frac{1}{x^3} dx$$

Rewrite as a limit.

$$= \lim_{t \rightarrow 0^-} \left(-\frac{1}{2x^2} \right) \Big|_{-1}^t$$

Find the antiderivative.

$$= \lim_{t \rightarrow 0^-} \left(-\frac{1}{2t^2} + \frac{1}{2} \right)$$

Evaluate the antiderivative.

$$= -\infty.$$

Evaluate the limit.

Therefore, $\int_{-1}^0 \frac{1}{x^3} dx$ diverges, and hence $\int_{-1}^1 \frac{1}{x^3} dx$ diverges regardless of the behavior of $\int_0^1 \frac{1}{x^3} dx$.

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

23 / 31

Integrating over an Infinite Interval
○○○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function
○○○○○○○○○○○○○○○○○○○○

Comparison Theorem
●○○○○○○○○

Comparison Property for Integrals

To see this, consider two continuous functions $f(x)$ and $g(x)$ satisfying $0 \leq f(x) \leq g(x)$ for $x \geq a$. In this case, we may view integrals of these functions over intervals of the form $[a, t]$ as areas. By the comparison property for definite integrals, we have the

relationship: $0 \leq \int_a^t f(x) \, dx \leq \int_a^t g(x) \, dx$ for $t \geq a$.

If $0 \leq f(x) \leq g(x)$ for $x \geq a$, then for $t \geq a$,

$$\int_a^\infty f(x) \, dx \leq \int_a^\infty g(x) \, dx.$$

Thus, if $\int_a^\infty f(x) \, dx = \lim_{t \rightarrow \infty} \int_a^t f(x) \, dx = \infty$, then

$$\int_a^\infty g(x) \, dx = \lim_{t \rightarrow \infty} \int_a^t g(x) \, dx \geq \lim_{t \rightarrow \infty} \int_a^t f(x) \, dx = \infty$$


as well.

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

25 / 31



Comparison Theorem

Let $f(x)$ and $g(x)$ be continuous over $[a, \infty)$. Assume that $0 \leq f(x) \leq g(x)$ for $x \geq a$.

- If $\int_a^\infty f(x) \, dx = \lim_{t \rightarrow \infty} \int_a^t f(x) \, dx = \infty$, then $\int_a^\infty g(x) \, dx = \lim_{t \rightarrow \infty} \int_a^t g(x) \, dx = \infty$.
- If $\int_a^\infty g(x) \, dx = \lim_{t \rightarrow \infty} \int_a^t g(x) \, dx = L$, where L is a real number, then $\int_a^\infty f(x) \, dx = \lim_{t \rightarrow \infty} \int_a^t f(x) \, dx = M$ for some real number $M \leq L$.

Integrating over an Infinite Interval
0000000000000000

Integrating a Discontinuous Function
000000000

Comparison Theorem
00000000

Applying the Comparison Theorem

Use the comparison theorem to show that $\int_1^\infty \frac{1}{xe^x} dx$ converges.

Solution:

- The integrand is continuous over $[1, \infty)$ and for $x > 1$:

$$0 \leq \frac{1}{xe^x} \leq \frac{1}{e^x} = e^{-x}$$

- So if $\int_1^\infty e^{-x} dx$ converges, then so does $\int_1^\infty \frac{1}{xe^x} dx$.
- To evaluate $\int_1^\infty e^{-x} dx$, first rewrite it as a limit:

$$\begin{aligned} \int_1^\infty e^{-x} dx &= \lim_{t \rightarrow \infty} \int_1^t e^{-x} dx = \lim_{t \rightarrow \infty} \left(-e^{-x} \right) \Big|_1^t \\ &= \lim_{t \rightarrow \infty} (-e^{-t} + e^1) = e. \end{aligned}$$

- Since the limit is finite, $\int_1^\infty e^{-x} dx$ converges, and hence, by the comparison theorem, so does $\int_1^\infty \frac{1}{xe^x} dx$.

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

27 / 31

Integrating over an Infinite Interval
○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function
○○○○○○○○○○

Comparison Theorem
○○○○○○●○

Applying the Comparison Theorem

Use the comparison theorem to show that $\int_1^\infty \frac{1}{x^p} dx$ diverges for all $p < 1$.

Solution:

- First we note that $\frac{1}{x^p}$ is continuous over $[1, \infty)$. If $p < 1$, then $\frac{1}{x} \leq \frac{1}{x^p}$ for all $x \in [1, \infty)$.
- We already showed that $\int_1^\infty \frac{1}{x} dx = \infty$. Therefore, by the comparison theorem, $\int_1^\infty \frac{1}{x^p} dx$ diverges for all $p < 1$.

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

30 / 31

Integrating over an Infinite Interval
○○○○○○○○○○○○○○○○○○○○

Integrating a Discontinuous Function
○○○○○○○○○○○○○○○○○○○○

Comparison Theorem
○○○○○○○○●

Key Concepts and Equations

Key Concepts

- Integrals of functions over infinite intervals are defined in terms of limits.
- Integrals of functions over an interval for which the function has a discontinuity at an endpoint may be defined in terms of limits.
- The convergence or divergence of an improper integral may be determined by comparing it with the value of an improper integral for which the convergence or divergence is known.

Key Equations

$$\int_a^\infty f(x) \, dx = \lim_{t \rightarrow \infty} \int_a^t f(x) \, dx$$
$$\int_{-\infty}^b f(x) \, dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) \, dx$$
$$\int_{-\infty}^\infty f(x) \, dx = \int_{-\infty}^0 f(x) \, dx + \int_0^\infty f(x) \, dx$$

Math 1700 (University of Manitoba)

3.7 Improper Integrals

March 15, 2024

31 / 31