

## Homework 1

Instructor: Nate Eldredge

Costantino Dufort Moraites

## 18.1 : 1, 2, 3, 4, 5

**Problem 1.** Let  $X$  be a random variable with  $X \geq 0$  almost surely. Prove: if  $\mathbb{E}[X] = 0$  then  $X = 0$  almost surely.

*Proof.* Say  $X \neq 0$ , but  $\mathbb{E}[X] = 0$  and let  $X_n$  be a sequence of simple random variables converging monotonically pointwise to  $X$ . Then by the monotone convergence theorem we have  $\mathbb{E}[X] = \lim \mathbb{E}[X_n]$ . Let  $A_{X_n} = \cup_{\{A_i | a_i > 0\}} A_i$  where  $\{a_i\}, \{A_i\}$  are the corresponding finite sets defining  $X_n$ . We have  $X \neq 0$  so there must be some  $A \subseteq \omega$ ,  $N$ , for which  $\mathbb{P}[A] > 0$  and  $X_j(\omega) > 0$  a.s. for all  $j \geq N$ ,  $\omega \in A$ , otherwise  $X = 0$ , a.s. for all  $n$ . Let  $a_M = \min\{a_i | a_i > 0\}$ , then  $\mathbb{E}[X_N] \geq a_M \cdot \mathbb{P}[A]$ . But then for  $j \geq N$ , and for all  $x \in A_{X_N}$ ,  $X_j(x) \geq a_M$  because  $\{X_n\}$  is a monotone sequence, thus  $\mathbb{E}[X_j] \geq a_M \cdot \mathbb{P}[A_{X_N}] > 0$  for all  $j \geq N$  and  $\mathbb{E}[X] = \lim \mathbb{E}[X_n] \neq 0$ , a contradiction.  $\square$

**Problem 2.** Let  $X \geq 0$  and let  $f : [0, \infty) \rightarrow [0, \infty)$  be continuous differentiable and monotone increasing.

- (a) Show that  $\mathbb{E}[f(X)] = \int_0^\infty f'(t)P(X \geq t)dt$ . Hint: Write  $P(X \geq t)$  as  $\mathbb{E}[1_{\{X \geq t\}}]$ . Now you have two integrals, one over  $[0, \infty)$  and the other over  $\Omega$ . Use Fubini/Tonelli's theorem.
- (b) In particular,  $\mathbb{E}[X] = \int_0^\infty P(X \geq t)dt$ .

**Solution.**

(a) First let's see that  $\mathbb{P}[X \geq t] = \mathbb{E}[1_{X \geq t}]$ . By definition  $\mathbb{P}[X \geq t]$  is the probability of the event that  $X \geq t$ , i.e.  $\int 1_{\{X \geq t\}} d\mu = \mathbb{E}[1_{\{X \geq t\}}]$ . This justifies writing:

$$\int_0^\infty f'(t)\mathbb{P}[X \geq t]dt = \int_0^\infty f'(t)\mathbb{E}[1_{\{X \geq t\}}]dt = \int_0^\infty f'(t) \int_\Omega 1_{\{X \geq t\}} d\mu dt = \int_0^\infty \int_\Omega f'(t) 1_{\{X \geq t\}} d\mu dt$$

We know that  $1_{X \geq t} \geq 0$  and because we are assuming  $f$  continuously differentiable and monotone,  $f'(t) \geq 0$  and we may apply Tonelli's theorem and interchange the order of integration to get:

$$\int_0^\infty \int_\Omega f'(t) 1_{\{X \geq t\}} d\mu dt = \int_\Omega \int_0^t f'(t) 1_{\{X \geq t\}} dt d\mu$$

Next we notice that  $\int_0^\infty f'(t) 1_{\{X \geq t\}} dt = \int_0^{X(\omega)} f'(t) dt$  as for  $t > X(\omega)$ ,  $1_{\{X \geq t\}}(\omega) = 0$ . This allows us to apply the fundamental theorem of calculus and we have:

$$\int_\Omega \int_0^t f'(t) 1_{\{X \geq t\}} dt d\mu = \int_\Omega f(X(\omega)) d\mu = \int_\Omega f(X) d\mu = \mathbb{E}[f(X)]$$

thus we have shown  $\mathbb{E}[f(X)] = \int_0^\infty f'(t)\mathbb{P}[X \geq t]dt$ , as desired.

(b)

This is immediate from part (a). Namely, the function  $f(t) = t$  is continuous differentiable and monotone increasing as a map  $[0, \infty) \rightarrow [0, \infty)$ , thus  $\mathbb{E}[f(X)] = \int_0^\infty f'(t)\mathbb{P}[X \geq t]dt = \int_0^\infty \mathbb{P}[X \geq t]dt$ , as desired.

**Problem 3.** Let  $X$  be a random variable with cumulative distribution function  $F$  (i.e.  $F(x) := P(X \leq x)$ ). Show that if  $F$  is continuous then  $Y = F(X)$  has a uniform distribution on  $[0, 1]$ , i.e.  $P(Y \leq t) = t$  for  $t \in [0, 1]$ . Give an example to show that this need not be true if  $F$  is not continuous. This is a sort of converse to Theorem 1.2.2.

*Proof.*

*The general result.*

We want to calculate  $P(Y \leq t) = P(F(X) \leq t)$ . We know that  $F$  is continuous and  $t \in [0, 1]$ , so assuming  $F$  assumes the values  $0, 1$  we can apply the intermediate value theorem to take an  $x$  so that  $F(x) = t$ . Further, we know the set of all such values  $y$  where  $F(y) = t$  is an interval because  $F$  is monotone increasing (this follows from the fact that  $F$  is defined by a probability and as  $x$  increases, the event  $\{X \leq x\}$  is a superset of the event  $\{X \leq x'\}$  for  $x' \leq x$ ). We know it is a closed interval because if it was an open interval  $(a, b)$  then we would have  $F(b - \epsilon) = x$  and  $F(b) = x'$  and  $F$  would not be continuous at  $b$ . This justifies letting  $b$  be the maximal number so that  $F(b) = t$ .

Now  $P(F(X) \leq t) = P(X \leq b) = F(b) = t$ , and  $Y$  has a uniform distribution on  $[0, 1]$  as desired.

*A counterexample when  $F$  is not continuous.*

Let  $X$  be the random variable corresponding to the flip of a fair coin. I.e.  $X = 1$  with probability  $1/2$  and  $X = 0$  with probability  $1/2$ . Then  $F(a) = P(X \leq a) = 1/2$  for  $a < 1$  and  $F(1) = P(X \leq 1) = 1$ , so  $F$  is not continuous at  $1$ . Now looking at  $P(Y \leq 3/4) = P(F(X) \leq 3/4) = P(X \leq 0) = 1/2 \neq 3/4$  and we have our desired counterexample.  $\square$

**Problem 4.** Let  $(A_i)$  be a sequence of events. Define:

$$\limsup_{n \rightarrow \infty} A_n := \bigcap_{m=1}^{\infty} \bigcup_{n=m}^{\infty} A_n$$

$\limsup A_n$  is also sometimes denoted  $\{A_n \text{ i.o.}\}$  (for “infinitely often”); it is the event that “infinitely many of the events  $A_n$  occur”.

- Show that  $1_{\limsup A_n} = \limsup 1_{A_n}$ .
- Show that  $P(\limsup A_n) \geq \limsup P(A_n)$ .
- Give an example to show that equality need not hold in the previous part. Indeed, try to find an example where  $P(\limsup A_n) = 1$  but  $\limsup P(A_n) = 0$ .
- Define  $\liminf A_n := \bigcup_{m=1}^{\infty} \bigcap_{n=m}^{\infty} A_n$ . This is also denoted  $\{A_n \text{ a.a.}\}$  and is the event that “all but finitely many of the events  $A_n$  occur”. By taking complements (or directly) show that  $1_{\liminf A_n} = \liminf 1_{A_n}$ ,  $P(\liminf A_n) \leq \liminf P(A_n)$  and that equality need not hold.

**Solution.**

- $1_{\limsup A_n} = \limsup 1_{A_n}$

*Proof.* Let's say  $1_{\limsup A_n}(\omega) = 1$ . Then  $\omega \in \limsup A_n$ . But this means that for every  $A_n$  such that  $\omega \in A_n$ ,  $1_{A_n}(\omega) = 1$  and so  $1_{A_n}(\omega) = 1$  infinitely often, i.e.  $\limsup 1_{A_n}(\omega) = 1$ . On the other hand, say  $1_{\limsup A_n}(\omega) = 0$ , then  $\omega \notin \limsup A_n$  and so there must be some last  $N$  so that  $\omega \notin A_j$  for all  $j \geq N$ . But then for all  $j \geq N$ ,  $1_{A_j}(\omega) = 0$  and so  $\limsup 1_{A_n}(\omega) = 0$ . So  $1_{\limsup A_n}, \limsup 1_{A_n}$  agree on all values, thus we have  $1_{\limsup A_n} = \limsup 1_{A_n}$ .  $\square$

(b)

*Proof.* First notice that  $\limsup P(A_n)$  is a  $\limsup$  of a sequence of real numbers and so we may choose a subsequence  $B_n$  of  $A_n$  so that  $\lim P(B_n) = \limsup P(A_n)$ . Let  $a(B_n)$  denote the index  $m$  of  $(A_n)$  so that  $B_n = A_m$ . Notice  $a(B_n) \geq n$ .

Now, if we write out:

$$\begin{aligned} P(\limsup A_n) &= \\ &= \int 1_{\limsup A_n} d\mu \end{aligned}$$

Which in particular makes us notice the monotone decreasing sequence of functions  $1_{\cup_{m=n}^{\infty} A_n} \rightarrow 1_{\limsup A_n}$ . So we may apply the monotone convergence theorem (well at least a 'standard' corollary of it) and write:

$$\begin{aligned} P(\limsup A_n) &= \\ &= \int 1_{\limsup A_n} d\mu \\ &= \lim_{n \rightarrow \infty} \int 1_{\cup_{m=n}^{\infty} A_n} d\mu \\ &= \lim_{n \rightarrow \infty} P(\cup_{m=n}^{\infty} A_n) \end{aligned}$$

Letting  $C_n = P(\cup_{m=n}^{\infty} A_n)$  we have  $C_n \geq P(B_n)$  for all  $n$  as  $a(B_n) \geq n$  and so  $B_n \subseteq \cup_{m=n}^{\infty} A_n$ . Thus we must have

$$\lim_{n \rightarrow \infty} P(\cup_{m=n}^{\infty} A_n) \geq \lim P(B_n)$$

So  $P(\limsup A_n) \geq \limsup P(A_n)$ , as desired.  $\square$

(c)

*Proof.* Let our probability space be the unit interval with Lebesgue measure. Let  $A_{(i,j)}$  be the  $j$ -th partition of  $[0, 1]$  into  $i$  equal sized pieces, let  $A_n$  denote the sequence  $A_{(1,1)}, A_{(2,1)}, A_{(2,2)}, A_{(3,1)}, \dots$ . Then  $\limsup P(A_n) = 0$  as the length of the interval  $A_n$  goes to 0 as  $n \rightarrow \infty$ , but for any finite  $m$ ,  $\cup_{n=m}^{\infty} A_n$  is the whole unit interval so  $\cap_{m=1}^{\infty} \cup_{n=m}^{\infty} A_n = \cap_{m=1}^{\infty} [0, 1] = [0, 1]$  and so  $P(\limsup A_n) = 1$ .  $\square$

(d)

*Proof.* For the first two parts, we redo the proofs directly and for the last part we appeal to 'taking complements' to get our counterexample.

Let's say  $1_{\liminf A_n}(\omega) = 1$ . Then  $\omega \in \liminf A_n$ . But this means there exists an  $N$  so that for all  $j \geq N$ ,  $\omega \in A_j$ , so  $1_{A_j}(\omega) = 1$  for all  $j \geq N$  and so  $\liminf 1_{A_n}(\omega) = 1$ . On the other hand, say  $1_{\limsup A_n}(\omega) = 0$ , then  $\omega \notin \limsup A_n$  and so for all  $N$  there exists a  $j > N$  such that  $\omega \notin A_j$ . But then for all  $N$ ,  $1_{A_j}(\omega) = 0$  and so  $\limsup 1_{A_n}(\omega) = 0$  thus  $1_{\limsup A_n}, \limsup 1_{A_n}$  agree on all values and we have  $1_{\limsup A_n} = \limsup 1_{A_n}$ .

For (b) I think it is unnecessary to rewrite the whole proof, I will just point out the places where the two proofs are different. We still choose a subsequence  $(B_n)$  so that  $\lim P(B_n) = \liminf P(A_n)$ , and when we rewrite  $P(\liminf A_n)$  as an integral we get a monotone increasing sequence of functions  $1_{\cap_{n=m}^{\infty} A_n}$  increasing to  $1_{\liminf A_n}$  and so we can use the monotone convergence theorem directly to interchange the integral and the limit. Now though, we find that  $\cap_{n=m}^{\infty} A_n \subseteq B_n$  for all  $n$  because  $a(B_n) \geq n$  and so  $P(B_n) \geq P(\cap_{n=m}^{\infty} A_n)$  for all  $n$ , thus we get the inequality from part b with the direction reversed, i.e.  $\liminf P(A_n) \geq P(\liminf A_n)$ .

Now for our counterexample, let our probability space be the unit interval with Lebesgue measure as before. Let  $A_{(i,j)}$  be the complement of the  $j$ -th partition of  $[0, 1]$  into  $i$  equal sized pieces, let  $A_n$  denote the sequence  $A_{(1,1)}, A_{(2,1)}, A_{(2,2)}, A_{(3,1)}, \dots$ . Then  $\liminf P(A_n) = 1$  as the length of the interval  $A_n$  goes to 1 as  $n \rightarrow \infty$  (it is the complement of an interval whose length goes to 0), but for any finite  $m$ ,  $\cap_{n=m}^{\infty} A_n = \emptyset$  because we can always pick a  $j \geq m$  so that way the next  $k$   $A_l$  are the complements of the partition of  $[0, 1]$  into  $k$  pieces. Each of these  $A_l$  are missing a different piece of the unit interval so their intersection is  $\emptyset$  and so  $P(\liminf A_n) = 0$ .  $\square$

**Problem 5.** Let  $1 \leq p \leq \infty$ . Suppose  $X_n$  is a sequence of random variables,  $X_n \rightarrow X$  a.s., and  $Y$  is another random variable such that  $|X_n| \leq Y$  and  $\mathbb{E}[Y^p] < \infty$ . Show that  $\mathbb{E}[|X|^p] \leq \infty$  and that  $\mathbb{E}[|X_n - X|^p] \rightarrow 0$ . This is a strengthening of the dominated convergence theorem.

*Proof.* Notice that  $X \leq Y$  almost everywhere. This follows from the fact that  $X_n \rightarrow X$  and  $X_n \leq Y$ . Thus we have  $X \leq Y$  so  $X^p \leq Y^p$  and so  $\mathbb{E}[X^p] \leq \mathbb{E}[Y^p]$ , say by theorem 1.4.7 in Durrett.

Now because  $X_n \rightarrow X$  a.e.,  $Z_n = |X_n - X| \rightarrow 0$  a.e., and similarly  $Z_n^p \rightarrow 0$  a.e. To be able to apply the dominated convergence theorem, it suffices to show that  $|Z_n|^p \leq h$  for  $h$  integrable, but we already know that

$$Z_n^p = |X_n - X|^p \leq |X_n|^p + |X|^p \leq 2^p Y^p$$

So we have that  $|Z_n^p| \leq 2^p Y^p$  and by assumption  $\mathbb{E}[Y^p] < \infty$  so  $\mathbb{E}[2^p Y^p] = 2^p \mathbb{E}[Y^p] < \infty$  and the dominated convergence theorem applies. Thus we have  $\mathbb{E}[Z_n^p] = \mathbb{E}[|X_n - X|^p] \rightarrow \mathbb{E}[0] = 0$ , as desired.  $\square$