

Grow with the flow: a latitudinal cline in physiology is associated with more variable precipitation in *Erythranthe cardinalis*

Christopher D. Muir^{1,*} and Amy L. Angert¹

¹ Biodiversity Research Centre, University of British Columbia, Vancouver, BC, Canada

*corresponding author: Chris Muir, cdmuir@biodiversity.ubc.ca

Running Head: Latitudinal cline and climate in *Erythranthe*

Key words: local adaptation, cline, photosynthesis, growth rate, [Erythranthe](#)

Data will be archived on Dryad upon publication.

Acknowledgements

Erin Warkman and Lisa Lin helped collect data. CDM was supported by a Biodiversity Postdoctoral Fellowship funded by the NSERC CREATE program. ALA was supported by an NSERC Discovery Grant and a grant from the National Science Foundation (DEB 0950171).

Grow with the flow: a latitudinal cline in physiology is associated with more variable precipitation in *Erythranthe cardinalis*

Abstract

Local adaptation is commonly observed in nature: organisms perform well in their natal environment, but poorly outside it. Correlation between traits and latitude, or latitudinal clines, are among the most common pieces of evidence for local adaptation, but identifying the traits under selection and the selective agents are challenging. Here, we investigated a latitudinal cline in growth and photosynthesis across 16 populations of the perennial herb *Erythranthe cardinalis* (Phrymaceae). Using machine learning methods, we identify interannual variation in precipitation as a likely selective agent: Southern populations from more variable environments had higher photosynthetic rates and grew faster. We hypothesize that selection may favor a more annualized life history – grow now rather than save for next year – in environments where severe droughts occur more often. Thus our study provides insight into how species may adapt if Mediterranean climates become more variable due to climate change.

Introduction

1 Local adaptation ~~within species~~ has been documented in numerous species; populations
2 generally have higher fitness in their native environment, but perform poorly outside it
3 (Schluter, 2000; Leimu and Fischer, 2008; Hereford, 2009). However, the prevalence of local
4 adaptation remains difficult to assess because researchers rarely test for local adaptation
5 unless there are obvious phenotypic or environmental differences (but see Hereford and
6 Winn 2008). When local adaptation occurs, it frequently leads to clines in both phenotypes

7 and allele frequencies when selection varies over environmental gradients (Huxley, 1938;
8 Endler, 1977; Barton, 1999). Phenotypic differences between populations along a cline
9 ~~most~~ often have a genetic basis and can be studied in a common garden (Turesson, 1922;
10 Clausen et al., 1940; Hiesey et al., 1942). Despite a long history of studying local adaptation
11 and clines, it remains challenging to identify exactly which traits are under selection and
12 which differ for nonadaptive reasons. In particular, the role that physiological differences
13 play in local adaptation is poorly understood, despite the fact that physiology is frequently
14 assumed to explain adaptation to the abiotic environment. A related problem is identifying
15 which of the myriad and often covarying aspects of the environment causes spatially varying
16 selective pressures.

17 When populations are locally adapted, reaction norms for fitness will cross, such that local
18 genotypes have higher fitness than foreign genotypes and rank orders change across envi-
19 ronments (Kawecki and Ebert, 2004). The traits that underlie local adaptation, however,
20 need not mirror this pattern. Populations can have fixed genetic differences conferring
21 trait values that are adaptive at home but neutral or maladaptive away. Alternatively,
22 the ability to plastically respond to a particular environment or the magnitude of response
23 to an environment could be adaptive. We distinguish between these patterns of adaptive
24 trait differences by referring to ‘intrinsic’ and ‘plastic’ trait variation, respectively. Both
25 intrinsic and plastic trait variation can be explained by genetic differences and both are
26 involved in adaptation. For example, intrinsic differences in photoperiod responses (Black-
27 man et al., 2011) and developmental rate (Stinchcombe et al., 2004) allow organisms to
28 properly time their life history with the local environment. Conversely, sun and shade
29 plants do not have intrinsically higher or lower rates of carbon assimilation, but rather,
30 genotype-by-environment interactions cause sun plants to assimilate more under high light
31 and shade plants under low light (Givnish, 1988). In plants especially, we know little about
32 the prevalence and adaptive significance of variation in fundamental physiological traits like
33 photosynthesis and their impact on plant performance.

34 A basic approach to identify candidate traits underlying local adaptation is to find asso-
35 ciations between traits and environments. Either intrinsic and/or plastic variation should
36 vary clinally along environmental gradients. Indeed, clines in ecologically important traits
37 are widespread in nature (Endler, 1977) and often adaptive, but in most cases the selective
38 agent is unknown. For example, in *Drosophila* numerous latitudinal clines exist for traits
39 like thermal tolerance (Hoffmann et al., 2002), body size (Coyne and Beecham (1987) and
40 references therein), and life history (Schmidt et al., 2005). Some *Drosophila* clines have
41 evolved multiple times (Oakeshott et al. (1982); Huey et al. (2000), see also Bradshaw and
42 Holzapfel (2001)) or shifted in response to climate change (Umina et al., 2005), evincing
43 climatic adaptation. Similarly, plant species exhibit latitudinal clines in traits like flowering
44 time (Stinchcombe et al., 2004), cyanogenesis (Kooyers and Olsen, 2012), leaf morphology
45 (Hopkins et al., 2008; Stock et al., 2014), and drought response (Kooyers et al., 2015) that
46 likely relate to climatic variation.

47 Despite the fact that latitudinal clines have been studied for a long time, latitude *per se*
48 cannot be a selective agent. Latitude may be strongly correlated with one or two key
49 climatic variables, such as temperature, precipitation, or growing degree-days. Latitude
50 may also correlate with the strength of biotic interactions (Schemske et al., 2009) or other
51 nonclimatic aspects of the environment, though as we explain below, we do not yet have
52 compelling data that these are important in our study system. Hence, we focus on whether
53 latitude could be an effective proxy for an underlying climatic driver, in which case we
54 would expect a yet stronger relationship between traits and the key climatic variable(s)
55 driving selection. Alternatively, latitude may be more strongly related to traits than any
56 single climatic variable for at least two reasons. First, latitude may be correlated with
57 several climatic agents of selection that are individually weak, but add up to a strong
58 latitudinal cline. Alternatively, gene flow among neighbouring populations could smooth
59 out local climatic effects, since alleles will experience selection across populations linked
60 by migration (Slatkin, 1978; Paul et al., 2011; Hadfield, 2016). We refer to this as the

61 ‘climatic neighborhood’. For example, in mountainous regions average temperature at
62 a given latitude varies widely, but in aggregate, a lower latitude set of populations will
63 experience warmer climate than a higher latitude one. Thus, any particular low latitude
64 population would be warm-adapted, even if it was located in a cooler (e.g. high elevation)
65 site. Because many climatic factors vary latitudinally, and which climatic factors vary
66 latitudinally changes over the earth’s surface (e.g. coastal vs. continental), dissecting the
67 evolution of latitudinal clines across many species will help identify generalities, such as
68 whether thermal tolerance maxima or seasonal timing is more important (Bradshaw and
69 Holzapfel, 2008), or whether local versus regional climate shape selective pressures.

70 In this study, we investigated two major questions: 1) whether intrinsic or plastic physiolog-
71 ical trait variation corresponds with latitude; and 2) what climatic factor(s) could plausibly
72 be responsible for latitudinal clines. Within question 2, we tested three hypotheses outlined
73 in the previous paragraph: latitudinal clines are explained by a single dominant climatic
74 factor, multiple climatic factors, or the climatic neighborhood experienced by nearby pop-
75 ulation connected through gene flow. These hypotheses are not mutually exclusive since,
76 for example, single or multiple factors in a climatic neighborhood may lead to latitudinal
77 clines. We focused on climate because climate often determines where species are found
78 and also can exert strong selection on populations within species [CITATION?]. There is
79 also a compelling need to know how populations are (or are not) locally adapted to cli-
80 mate so as to predict how they’ll respond to climate change [CITATION?]. We did not
81 test for local adaptation to other environmental characteristics, such as edaphic or biotic
82 (e.g. competitors, pollinators) factors, because we did not have compelling observations
83 from which to make testable hypotheses about the latter in our focal species (described
84 below).

85 We examined these questions in *Erythranthe cardinalis* (formerly *Mimulus cardinalis* [Ne-
86 som 2014]) because linking physiological traits to potentially complex patterns of local
87 adaptation requires integrating multiple lines of evidence from comparative, experimental,

88 and genomic studies under both lab and field conditions. Many classic and contemporary
89 studies of local adaptation use *Mimulus sensu lato* species because of ~~its~~^{its} natural history,
90 easy propagation, and genetic/genomic resources (Clausen et al., 1940; Hiesey et al., 1971;
91 Bradshaw and Schemske, 2003; Wu et al., 2008; Lowry and Willis, 2010; Wright et al.,
92 2013). Yet, there is a conspicuous deficiency of links between local adaptation and physio-
93 logical mechanisms (Angert (2006); Angert et al. (2008); Wu et al. (2010), **but see** Wright
94 et al. (2013) on mechanisms of copper tolerance). We measured genetic and genotype-
95 by-environment variation in response to temperature and drought among 16 populations
96 distributed over 10.7° of latitude. We found a latitudinal cline of intrinsic variation in pho-
97 tosynthesis and growth, but little evidence for variation in plasticity. Interannual variation
98 in precipitation and temperature are associated with this axis of variation, suggesting that
99 climatic variance rather than mean may be an important driver of local adaptation in
100 *E. cardinalis*. The climatic neighborhoods around populations explained trait variation
101 better than local climate, indicating that latitudinal clines may be common because lati-
102 tude integrates effects of selection on populations connected through gene flow. We place
103 these findings in the context of life history theory and consider future directions in the
104 Discussion.

105 **Material and Methods**

106 **Population Selection**

107 We used 16 populations from throughout the range of *E. cardinalis* (Table 1). These
108 populations were intentionally chosen to span much of the climatic range of the species
109 based on all known occurrences (see below). Seeds were collected in the field from mature,
110 undehisced fruit left open for 2-4 weeks to dry, then stored at room temperature. We used
111 seeds from 154 families, 4-12 (mean = 9.6, median = 12) families per population.

Table 1: Geographic region, latitude, longitude, and elevation (mas = meters above seal level) of 16 focal populations used in this study.

Name	Region	Latitude	Longtiude	Elevation (mas)
Hauser Creek	South Margin	32.657	-116.532	799
Cottonwood Creek	South Margin	32.609	-116.7	267
Sweetwater River	South Margin	32.9	-116.585	1180
Grade Road Palomar	South Margin	33.314	-116.871	1577
Whitewater Canyon	Transverse	33.994	-116.665	705
Mill Creek	Transverse	34.077	-116.873	2050
West Fork Mojave River	Transverse	34.284	-117.378	1120
North Fork Middle Tule River	South Sierras	36.201	-118.651	1314
Paradise Creek	South Sierras	36.518	-118.759	926
Redwood Creek	South Sierras	36.691	-118.91	1727
Wawona	Central Sierras	37.541	-119.649	1224
Rainbow Creek	Central Sierras	37.819	-120.007	876
Middle Yuba River	North Sierras	39.397	-121.082	455
Little Jamison Creek	North Sierras	39.743	-120.704	1603
Deep Creek	North Coast	41.668	-123.11	707
Rock Creek	North Margin	43.374	-122.957	326

Plant propagation

On 14 April, 2014, 3-5 seeds per family were sown directly on sand (Quikrete Play Sand, Georgia, USA) watered to field capacity in RLC4 Ray Leach cone-tainers placed in RL98 98-well trays (Stuewe & Sons, Inc., Oregon, USA). We used pure sand because *E. cardinalis* typically grows in sandy, riparian soils (A. Angert, pers. obs.). Two jumbo-sized cotton balls at the bottom of cone-tainers prevented sand from washing out. Cone-tainers sat in medium-sized flow trays (FLOWTMD, Stuewe & Sons, Inc., Oregon, USA) to continuously bottom-water plants during germination in greenhouses at the University British Columbia campus in Vancouver, Canada (49°15' N, 123°15' W). Mistlers thoroughly wetted the top of the sand every two hours during the day. Most seeds germinated between 1 and 2 weeks, but we allowed 3 weeks before transferring seedlings to growth chambers. We recorded germination daily between one to two weeks after sowing, and every 2-3 days thereafter.

124 On 5 May (21 days after sowing), we transferred seedlings to one of two growth chambers
125 (Conviron, Manitoba, Canada). We thinned seedlings to one plant per cone-tainer, leaving
126 the center-most plant. 702 of 768 (91.4%) had plants that could be used in the experiment.
127 We allowed one week at constant, non stressful conditions (day: 20°C, night: 16°C) for plants
128 to acclimate to growth chambers before starting treatments. The initial size of seedlings,
129 measured as the length of the first true leaves, did not differ between populations, families,
130 or treatments (Table S1).

131 **Temperature and drought treatments**

132 We imposed four treatments, a fully-factorial cross of two temperature levels and two
133 watering levels. The temperature levels closely simulated an average growing season at the
134 thermal extremes of the species range, which we designate as Hot and Cool treatments.
135 Watering levels contrasted a perennial and seasonal stream, which we refer to as Well-
136 watered and Drought treatments. A detailed description of treatments is provided in the
137 Supplemental Materials and Methods and summarized in Fig 1. Because growth chambers
138 cannot be subdivided, one chamber was assigned to the Hot treatment level and another
139 to the Cool treatment level. Within each chamber, there were two Well-watered blocks
140 and two Drought blocks. The photosynthetically active radiation in both chambers was
141 approximately 400 $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$. The growth chambers did not control humidity,
142 but because of watering and high plant transpiration rates, the relative humidity was quite
143 high in both temperature levels (data not shown). Lower humidity would have made the
144 drought more severe, but low soil moisture is stressful in and of itself. The total number of
145 plants in each treatment was: $n_{\text{cool,dry}} = 169$; $n_{\text{cool,ww}} = 174$; $n_{\text{hot,dry}} = 176$; $n_{\text{hot,ww}} = 183$.
146 Each population had 8–12 individuals per treatment level (mean = 11, median = 11).




Figure 1: Overview of experimental treatments and timing of key trait measurements. All plants germinated within 21 days of sowing. At that time, we began temperature treatments (left axis), simulating a typical June-August weather pattern at Hot (red) and Cool (blue) sites. The bold lines track the average daily temperatures. Within each day, there was a maximum daytime temperature (top of translucent polygons) and minimum nighttime temperature (bottom of translucent polygons). The drought treatment commenced later by ramping down the frequency of bottom-watering episodes (dashed black line; right axis), while watering frequency was maintained in the control treatment (solid black line). Grey boxes on the bottom of the plot outline the period of key measurements described in the Material and Methods.

Trait measurements

We measured five traits in response to temperature and watering treatments (Table 2).

Days to germination We tested for population variation in germination rate, measured as Days to Germination, using a lognormal survival model fit using the `survreg` function in the R package **survival** version 2.38 (Therneau, 2015). We treated Population as a fixed effect and Family as random effect using a Γ frailty function. Statistical significance of the

Table 2: Key traits measured in this study.

Trait	Units
Days to germination	day
Leaf expansion rate	mm day ⁻¹
Stem elongation rate	cm day ⁻¹
Photosynthetic rate	$\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$
Mortality	probability of death

Population effect was determined using analysis of deviance. Note that, unlike other traits discussed below, we did not include Block, Treatment, or Population \times Treatment interactions because during germination plants had not been placed into blocks and treatments had not yet been applied.

Growth rate: leaf expansion and stem elongation We measured growth rate during two phases: leaf expansion and stem elongation. Growth measurements were taken during the early vegetative stage. We censused leaf length twice per week shortly after the emergence of true leaves from 12 May – 12 June (28–59 days after sowing), resulting in 10 measurements. We ceased measuring leaf length once it appeared to asymptote and growth shifted to stem elongation. We also censused plant height on 7 occasions (twice per week) between 29 May and 20 June (45 to 67 days after sowing) until plants began to initiate floral buds. Thus all growth measurements occurred during the vegetative, pre-reproductive phase. Both leaf expansion and stem elongation were modelled separately as second-order polynomials. We used empirical Bayes’ estimates of growth for each individual plant from linear mixed-effects models fit with the R package **lme4** version 1.1-12 (Bates et al., 2015).

Photosynthesis During the week of 10 to 16 June (57 to 63 days after sowing), we measured daytime photosynthetic rate on a subset of 329 plants evenly spread between

171 treatments and families within populations. The youngest, fully-expanded leaf acclimated
 172 for 3 minutes to reach steady state in a 6-cm² chamber of a LI-COR 6400XT Portable Pho-
 173 tosynthesis System (LI-COR Biosciences, Lincoln, Nebraska). We made all measurements
 174 at ambient light (400 $\mu\text{mol m}^{-2} \text{ s}^{-1}$ of photosynthetically active radiation), atmospheric
 175 CO₂ (400 ppm), temperature, and moderate relative humidity. During this period, we sus-
 176 pended normal day-to-day temperature fluctuations and set daytime temperatures to the
 177 average for that period (Cool: 26.5°; Hot: 36.1°) so that all plants within a temperature
 178 level could be measured under the same conditions.

179 **Mortality** We assayed mortality during twice-weekly growth measurements. We ana-
 180 lyzed the probability of surviving until the end of the experiment as a function of popula-
 181 tion, treatment, and their interactions using a Generalized Linear Mixed Model (GLMM)
 182 assuming binomially distributed errors. We included Family and Block as random effects.
 183 We assessed significance of fixed effects using Type-II Analysis of Deviance with Wald χ^2
 184 tests in the R package **car** (Fox and Weisberg, 2011).

185 **Intrinsic variation and plasticity**

186 For all traits (Table 2) except germination (see above), we tested for Population, Treat-
 187 ment (Temperature, Water, and Temperature \times Water), and Population \times Treatment
 188 interactions (Population \times Temperature, Population \times Water, and Population \times Temper-
 189 ature \times Water). We interpreted significant Population effects to indicate intrinsic variation
 190 and Population \times Treatment interactions to indicate variation in plasticity. As mentioned
 191 above, we used survival and GLMM models for germination rate and mortality, respec-
 192 tively. For all other traits, we used mixed model ANOVAs with Family and Block included
 193 as random factors. We fit models using restricted maximum likelihood in **lmer**, a function
 194 in the R package **lme4** (Bates et al., 2015). We determined significant fixed effect terms us-

ing a step-wise backward elimination procedure implemented with the step function in the R package **lmerTest** version 2.0-32 (Kuznetsova et al., 2016). This package uses Satterthwaite’s approximation to calculate denominator degrees of freedom for F -tests. We also included days to germination as a covariate in growth analyses. To ensure that Population and Treatment effects were specific to a particular growth phase, we included germination day as a covariate in leaf expansion and stem elongation analyses.

Principal components of germination, growth, and photosynthesis

For each single-trait model above, we extracted the Population coefficient (factoring out Treatment and other effects). The multivariate distribution of these coefficients was then summarized using principal components analysis. The first principal component of these traits (TraitPC1) loaded positively with germination, growth, and photosynthetic rate, therefore we define this as a phenotypic axis delineating fast to slow growth.

Identifying putative selective agents

Latitudinal clines are common, but it is often difficult to ascribe this variation to a particular selective agent. To reiterate, we tested three non-mutually exclusive hypotheses about how such latitudinal clines emerge: 1) one or two climatic variable explains latitudinal trait variation; 2) latitude is a proxy for multiple climatic factors that together shape trait variation; and 3) latitude integrates selection in a broader climatic neighborhood. We found that a population’s position along TraitPC1 correlated strongly with the latitude of origin (see Results) and next used Random Forest regression (Liaw and Wiener, 2002) to identify putative climatic factors underlying trait-latitude associations in *E. cardinalis*. We reasoned that if we identified a single climatic factor that explained more trait variation than latitude, then this would suggest that factor is a key selective agent underlying the latitudinal cline (Hypothesis 1). On the other hand, if multiple climatic factors together

219 are necessary to explain trait variation, then this would suggest that many climatic factors
 220 together have imposed selection for the latitudinal cline (Hypothesis 2). We hereafter refer
 221 to factors identified in this analysis as ‘Climate-TraitPC1’ variables. To test Hypothesis
 222 3 about climatic neighborhoods driving selection, we directly compared local with neigh-
 223 borhood climate. We used the immediate collection location for local climate. For climate
 224 neighborhoods, we sampled climate at 1000 random points (at 90-m resolution) within a
 225 62-km radius buffer around the collection and took the average. We chose this buffer size
 226 because neutral genetic differentiation increases slowly with geographic distance, indicating
 227 significant gene flow between nearby populations (Paul et al., In review). Significant spatial
 228 autocorrelation persisted for approximately 62 km. Since *E. cardinalis* is found exclusively
 229 in riparian areas, we only selected points along streams using the National Hydrography
 230 Dataset (United States Geological Survey, 2015). Climatic means and variances (see below)
 231 were weighted by their climatic suitability as determined using a multimodel ensemble av-
 232 erage of ecological niche models (Angert et al., 2016). In addition to competing local and
 233 neighborhood climate, we compared the univariate correlation between local and neigh-
 234 borhood climate with TraitPC1 and Latitude using paired *t*-tests. We adjusted degrees
 235 of freedom to account for the fact that many climatic factors are highly correlated and
 236 not independent. Specifically, we calculated the effective number of independent climatic
 237 factors (M_{eff}) using the formula $M_{\text{eff}} = 1 + (M - 1)(1 - \text{Var}(\lambda)/M)$ (Chevrud, 2001), where
 238 M is the original number of climatic factors and λ are the eigenvalues of the correlation
 239 matrix of all climatic factors.

240 To help eliminate potentially spurious correlations between TraitPC1 and climate, we tested
 241 for overlap between climatic variables that best predict latitude of all *E. cardinalis* occur-
 242 rence records (see detail below), not just the 16 focal populations. We refer to these climatic
 243 factors as ‘Climate-Latitude’ variables. The logic is that climatic factors associated with
 244 both TraitPC1 and latitude for all populations are more likely to be important selective
 245 agents than climatic factors that happen to correlate with TraitPC1 but do not covary

with latitude throughout the *E. cardinalis* range. Therefore, we did not consider Climate-TraitPC1 variables to be candidate selective agents unless the same or very similar variable was found in the Climate-Latitude analysis.

We selected Climate-Latitude and Climate-TraitPC1 variables independently using Variable Selection Using Random Forest (VSURF) algorithm in the R package **VSURF** version 1.0.3 (Genuer et al., 2016). Random Forest regression is useful for cases like ours when the number of potential predictors is similar to or greater than the number of observations ('high p , low n ' problem). VSURF is a multistep algorithm that progressively retains or eliminates variables based on their importance over regression trees in the forest. Variable importance is defined as the average amount a climate variable reduces mean-squared error in the predicted response (TraitPC1 or Latitude), compared to a randomly permuted dataset, across all trees in the random forest (see Genuer et al. [2015] for further detail). Hence, VSURF automatically eliminates unimportant and redundant variables based on the data without having to arbitrarily choose among colinear climate variables before the analysis. We kept only variables selected for prediction, the most stringent criterion. A visual overview of how we selected climatic variables is depicted in Fig 2.

For Climate-Latitude analyses, we compiled a representative set of 356 recent (since 2000) known *E. cardinalis* occurrences. These occurrences were thinned by 50% to correct for uneven sampling from a comprehensive set of herbarium records and an exhaustive field survey in 2010-11 (Angert et al., 2016). For both Climate-TraitPC1 analyses (16 focal populations) and Climate-Latitude (many populations), we used a 90-m digital elevation model from HydroSHEDS (Lehner et al., 2006) to extract elevation. Monthly interpolated climate layers were calculated using ClimateWNA version 5.30 (Wang et al., 2012), which accurately downscales climate data specifically for the rugged topography of western North America. For each occurrence, we calculated bioclimatic variables using the biovars function in the R package **dismo** version 1.1-1 (Hijmans et al., 2016). We included 24 climatic factors, 9 from ClimateWNA and 15 bioclimatic variables (Table S2). The bioclimatic

variables included all permutations of two climatic factors, temperature and precipitation, and six temporal scales (annual average, coldest quarter, warmest quarter, wettest quarter, driest quarter, or seasonality) as well as mean diurnal range, isothermality, annual temperature range. For each variable, we calculated both a 30-year normal by averaging annual values between 1981 and 2010 and 30-year coefficient of variation, a standardized metric of interannual climatic variation. Temperatures were converted to Kelvin to be on a ratio scale appropriate for calculating the coefficient of variation (CV). In total, the VSURF algorithm selected among 96 climate variables: 24 climatic factors \times 2 types (30-year average and CV) \times 2 spatial scales (local and neighborhood).

Results

A coordinated latitudinal cline in germination, growth, and photosynthesis

There are strong genetically-based trait differences in time to germination, growth, and photosynthetic rate among populations of *E. cardinalis*, as evidenced by large and significant population effects for these traits (Table 3). A single principal component captured 71.6 % of the trait variation among populations, defining an axis of variation from fast to slow growth. A population's position along this axis strongly covaried with its latitude of origin; southern populations grew faster than northern populations (Fig 3). ~~As we explain below, intrinsic differences between populations in terms of plant function (photosynthesis) and performance (growth) contrasted with the low amount of variation in plasticity.~~ There were similar latitudinal clines for individual traits underlying PC1 (Figures S1 to S4).

Table 3: Summary of Population, Treatment, and Population \times Treatment effects. We used different statistical modeling for the diverse traits assayed – glmer: generalized linear mixed model using the R package **lme4** (Bates et al., 2015); lmer: linear mixed model using the R package **lme4** (Bates et al., 2015); survreg: survival regression using the R package **survival** (Therneau, 2015). Note that temperature and water treatments were imposed after germination, hence are not applicable to this trait. Complete analysis of variance/deviance tables for each trait are available in the Supporting Information. Key to statistical significance: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

Trait	Germination	Leaf expansion	Stem elongation	Photosynthesis	Mortality
Statistical model	survreg	lmer	lmer	lmer	glmer
Population	***	***	***	***	
Temperature	NA	***	***	**	***
Water	NA	*			***
Pop \times Temp	NA			*	
Pop \times Water	NA	*			
Temp \times Water	NA				***
Pop \times Temp \times Water	NA				

Little evidence for variation in plasticity

Genotype \times environment (G \times E) interactions are also a common signature of local adaptation. We found little evidence of G \times E in *E. cardinalis*. There were only two statistically significant Population \times Treatment interactions (Table 3, Fig. S5), but these were not strong compared to Population and Temperature effects. Otherwise, populations responded similarly to treatments: faster growth in the hot treatment, slower growth in the dry treatment, and high mortality in the hot, dry treatment (Table 3). Note that interactions were calculated after factoring out intrinsic trait differences, necessarily reducing statistical power to detect significant interactions relative to main effects. However, the fact that the Population and Temperature effects were often highly significant ($P \ll 0.001$ in most cases) suggests that statistical power alone cannot explain why we failed to detect Population \times Treatment interactions. Complete ANOVA tables are available in the Supporting Information (Tables S3 to S6)

Neighborhood climatic variability best explains latitudinal cline

Interannual variation in climate averaged over ~~62-km radius~~ climatic neighborhoods ~~around~~ ~~populations~~ correlated most strongly with trait variation and latitude of *E. cardinalis* occurrences (Fig. 4, Table S7). All 16 Climate-Latitude and 3 Climate-TraitPC1 variables were neighborhood rather than local variables (Fig. 4). In fact, neighborhood climate almost always correlated better with TraitPC1 and Latitude than local climate (Fig. S6). On average, neighborhood Climate-TraitPC1 correlation coefficients were 0.16 higher than correlations with local-scale climate variables (paired *t*-test, $t = 7.87$, d.f. = 33.6, $P = 3.94 \times 10^{-9}$). Likewise, neighborhood Climate-Latitude correlation coefficients were 0.13 higher than those for local-scale climate (paired *t*-test, $t = 6.71$, d.f. = 36.8, $P = 7.22 \times 10^{-8}$).

The most important climatic variables related to the interannual variation rather than average climate. Among Climate-Latitude and Climate-TraitPC1 variables, neighborhood climatic variability over 30 years (1981–2010) in either winter precipitation (bio16_σ) and/or temperature (bio11_σ) are the strongest candidates driving the latitudinal cline in *E. cardinalis* (see Table S2 for a key to climate variable abbreviations). Note that the coefficient of variation of a climatic factor is subscripted with σ whereas the mean is subscripted with μ . More specifically, greater winter precipitation variability and lower winter temperature variability are associated with Southern latitudes and higher TraitPC1 values (Fig. 5A,B). Neighborhood interannual variation in winter precipitation (bio16_σ) was the most important Climate-Latitude variable (Fig. 4A). However, neighborhood bio16_σ did not overlap with Climate-TraitPC1 variables (Fig. 4B). We nevertheless consider it a plausible candidate for two reasons. First, neighborhood bio16_σ correlated strongly with TraitPC1 (Fig. 5A). Second, one of the most important Climate-TraitPC1 variables (neighborhood bio15_σ ; Fig. 5B,C) is very similar to bio16_σ . In Mediterranean climates like California, most precipitation occurs in the wettest quarter (winter), so years with low winter precipitation

also have low precipitation seasonality. Hence, highly variable year-to-year winter precipitation at lower latitude (Fig. 5D) is closely associated with large swings in precipitation seasonality (Fig. 5C).

Interannual variation in temperature of the coldest quarter (neighborhood bio11_σ) is another plausible candidate because it was only variable in both Climate-Latitude and Climate-TraitPC1 analyses (Fig. 4). Neighborhood bio11_σ explained more variation in TraitPC1 than latitude (latitude $r^2 = 0.55$ vs. bio11_σ $r^2 = 0.6$; Fig. S7), whereas neighborhood bio16_σ did slightly worse (bio16_σ $r^2 = 0.49$). Models using bio15_σ or bio11_σ to predict TraitPC1 also had significantly lower Akaike Information Criteria (AIC) than the latitude model (AIC of different models – bio15_σ : 48.5; bio11_σ : 52.4; latitude: 54.5). The best two-factor model including both neighborhood bio15_σ and bio11_σ did not significantly improve explanatory power ($r^2 = 0.71$, AIC= 49.2). In summary, either variation in precipitation or temperature seasonality may be important selective agents, but there is no strong evidence that they are both important. The most important Climate-TraitPC1 variable, neighborhood variation in mean diurnal range (bio2_σ ; Fig. 4B) did not have any obvious similarity to Climate-Latitude variables. Given the large number of potential associations, we therefore think this may be a spuriously strong relationship ~~and do not consider it further~~.

Discussion

We found evidence for one of two common signatures of local adaptation in the perennial herb *Erythranthe cardinalis*. Latitudinal clines in germination rate, photosynthesis, and growth suggest adaptive differentiation in important physiological traits of the species. However, we found little evidence that populations respond differently to temperature or drought. Due to low replication within families, we did not have power to assess within population genotype-by-environment interactions, which may be present. As we discuss

below, low variation in plasticity among populations ~~result~~ may indicate that some dimensions of the fundamental abiotic niche are relatively conserved. Finally, our results suggest that neighborhood-scale climate and interannual variation are more important selective agents than local averages. In the paragraphs that follow, we tie these results into the broader threads of evolutionary theory that might help explain why intrinsic variation in physiology changes clinally, whereas plastic responses to temperature and drought are relatively similar. One caveat to bear in mind is that we are limited by the size of the climate grid ($\approx 90 \text{ m}^2$) and therefore unable to detect very fine-scale local adaptation.

Evolutionary theory indicates that the shape of fitness tradeoffs, demography, and gene flow can constrain adaptation (Levins, 1968; Ronce and Kirkpatrick, 2001; Lenormand, 2002) and hence the type of variation maintained within species. Specifically, adaptive variation cannot be maintained by spatially varying selection if tradeoffs are too strong, demography is strongly asymmetric, and/or maladaptive gene flow is too high. To clarify, strong tradeoffs can prevent local adaptation in variable environments because selection favors habitat specialists that track a specific habitat regardless of its frequency in the environment (Levins, 1968). For example, a riparian specialist may experience similar selection in rivers of highfall regions and deserts, even though the habitat is much rarer in the latter. In *E. cardinalis* we found substantial genetically based variation among populations along a phenotypic axis from fast to slow growth that varied over a large spatial scale (Fig. 3). If this variation is adaptive, it suggests one of several possibilities to investigate in the future: fitness tradeoff between low versus high latitude environments is not too strong nor swamped by demographic asymmetry or maladaptive gene flow. That is, alleles favoured at one latitude are not strongly selected against when they flow to another population, allowing locally adaptive genetic variation to be maintained by spatially heterogeneous selection. We also know from previous work that population size does not vary strongly with latitude. Gene flow appears to be high, but attenuates at broad spatial scales, especially between Southern ($< 35^\circ\text{N}$) and Northern portions of the



385 range (Paul et al., In review).

386 Nevertheless, local gene flow from similar environments may shape how selection varies
387 with latitude. Theory predicts that populations will not be perfectly adapted to their
388 immediate habitat when there is gene flow from surrounding populations with different
389 optima (Lenormand, 2002). With spatial heterogeneity and gene flow, traits will not covary
390 perfectly with the local optimum (Slatkin, 1978; Paul et al., 2011; Hadfield, 2016), but
391 should instead better match the average environment experienced by nearby populations
392 connected through gene flow, which we refer to as the climatic neighborhood. Gene flow
393 and spatial heterogeneity may therefore be important in maintaining genetic variation
394 (Yeaman and Jarvis, 2006). As this hypothesis predicts, climatic neighborhoods (62-km
395 buffer around populations) correlated with traits and latitude of occurrences better than
396 local climate (Fig. 4). We interpret this as suggestive evidence that gene flow between
397 neighboring *E. cardinalis* populations shapes selection – populations are locally adapted to
398 prevailing climate in their neighborhood, but perhaps not perfectly adapted to their local
399 climate. This may not greatly constrain local adaptation because local and neighborhood
400 climate values ~~at the resolution of ClimateWNA (90 m²)~~ were generally similar in *E.*
401 *cardinalis* populations (Fig. S6). Therefore, we would predict in reciprocal transplants
402 that populations whose local climate is farther from their neighborhood average would be
403 less well adapted than those close to their neighborhood average.

404 It is reasonable to predict that southern populations, which appear to experience more
405 frequent drought years (see below), might have physiological adaptation to survive and
406 grow in drier soil. We found no evidence for this type of drought tolerance; all popula-
407 tions responded to drought and temperature similarly (Table 3). Plants grew faster in
408 the Hot treatment, but there was little effect of drought on growth. Rather, the effects
409 of drought took longer to materialize but resulted in high mortality, especially in the Hot
410 treatment. However, there was no differential mortality among populations in this treat-
411 ment. Although our results indicate that this axis of the species niche may be constrained,

412 plants have multiple ways to resist drought through both tolerance and escape (Ludlow,
413 1989; Kooyers, 2015). Next, we consider why drought tolerance may be less important in local
414 adaptation than a form of escape for this species.


415 We hypothesize that tolerance to dry soil may be constrained by a combination of strong
416 fitness tradeoffs, demographic asymmetry, and gene flow. Soil moisture in riparian habitats
417 where *E. cardinalis* live are highly heterogeneous at very small spatial scales (several me-
418 ters). Plants in the stream never have to tolerate drought whereas plants only a few meters
419 away may experience extreme drought since there is little direct precipitation during the
420 growing season in Mediterranean climates of western North America. We hypothesize alle-
421 les that confer greater drought tolerance may be quite costly in well-watered soils, and *vice*
422 *versa*, leading to strong fitness tradeoffs. Such tradeoffs would promote specialization to
423 one soil moisture or another, thereby inhibiting the evolution of broad environmental tol-
424 erance within a population. Demography and gene flow may reinforce niche conservatism.
425 A new mutant with increased drought tolerance that could survive at the resource-poor
426 margin of a population would likely be demographically overwhelmed by the larger census
427 populations that can be maintained in higher-resource environments. Infrequent wet years
428 may also produce most seeds, so selection is weighted towards alleles that have high fitness
429 in the wet environment, even if dry years are more frequent (Templeton and Levin, 1979;
430 Brown and Venable, 1986). Finally, gene flow, which is generally high among *E. cardinalis*
431 populations within the same ecoregion (Paul et al., In review), will thwart local adapta-
432 tion and reinforce specialization. Thus, the spatial grain of the environment, demographic
433 asymmetry, and gene flow may conspire to constrain local adaptation along this environ-
434 mental axis. Consistent with this hypothesis, recent record-setting droughts have caused
435 the decline or even local extinction of some natural populations of *E. cardinalis* (Sheth and
436 Angert, 2017).

437 In sum, these results indicate that differences in physiology and growth, but not responses
438 to temperature and drought, mediate local adaptation to climate in *E. cardinalis*. Next,

we would like to understand why variation in these particular traits may be adaptive. We argue that temporally more variable environments, as experienced by southern populations, select for a more ‘annualized’ life-history strategy, a form of drought escape. Demographic observations in natural populations of *E. cardinalis* reveal that southern populations tend to flower earlier at a smaller size, while northern populations invest more in vegetative growth (Sheth and Angert, 2017). The association between position along the ‘fast-slow’ continuum and associated traits in *E. cardinalis* is similar to interspecific relationships between growth, functional traits, and life history (Adler et al., 2014; Salguero-Gómez et al., 2016). However, we cannot exclude unexplored factors (e.g. edaphic conditions, competitors, pollinators, etc.) which may drive the latitudinal cline.

Greater investment in aboveground growth, as opposed to belowground storage for future seasons, may be favoured in climates with more frequent drought years, but maladaptive in climates with more consistent precipitation. This is a form of drought escape in that plants are investing more reproduction in the present to avoid possible drought in subsequent years. Suppose plants that grow quickly and allocate new resources to continued growth rather than storage have higher fitness over a single growing season. However, by not allocating resources to storage, these fast-growing plants begin future seasons at a deficit. Therefore, in a stable environment where winter survivorship is assured in most years, failure to store resources may reduce lifetime fitness. But ~~the environment is not constant.~~ For perennial herbs in Mediterranean climates, a dry winter (rainy season) can kill the rhizomes (underground stems that store nutrients for future growth) before emergence or aboveground stems before flowering. If drought years occur frequently enough, selection may favour the fast-growing strategy because there is no advantage to storage if drought kills plants before flowering. Considering life-history strategy as a continuum from no storage (annual) to lots of storage (perennial), we hypothesize that the optimal allocation to aboveground growth is more ‘annualized’ in southern climates that have greater interannual variation in precipitation. This scenario differs from classic drought escape syndromes in

466 which plants speed up development early in the season before the onset of drought.

467 The hypothesis that greater precipitation variability selects for an annualized life history
468 is tentative, but consistent with theory and data from other species. Life history theory
469 shows that less variable environments are one factor that favours the evolution of perenni-
470 ality (Stearns, 1976; Iwasa and Cohen, 1989; Friedman and Rubin, 2015). Populations of
471 the perennial *Plantago asiatica* show a similar latitudinal cline in growth and allocation to
472 storage (Sawada et al., 1994), though these authors attribute the cline to variation in grow-
473 ing season length. There are also life history clines in the closely related species *E. guttata*,
474 but the underlying traits and climatic drivers are quite different. Annual *E. guttata* flower
475 sooner and produce fewer stolons in response to climates with shorter seasons and more
476 intense summer drought (Lowry and Willis, 2010; Friedman et al., 2015; Kooyers et al.,
477 2015). In contrast, there are no truly annual (monocarpic and semelparous) populations of
478 *E. cardinalis*. Rather, our hypothesis states that climatic variability selects on quantitative
479 variation in allocation to growth versus storage. This hypothesis makes several indepen-
480 dent, testable predictions. The allocation tradeoff predicts that northern populations will
481 provision more **assimilate** to rhizomes compared with southern populations. If southern
482 populations are indeed more ‘annualized’ because more frequent droughts cause mortality,
483 then we predict that species distribution models using recent climate would best predict
484 occurrences in the south, whereas longer term climate would be a better predictor in the
485 north. Finally, we predict that southern populations would show greater variation in the
486 size of recruits and higher maximum population growth rates. **Exploratory analysis** also 
487 pointed to variation in winter temperature as a potential selective agent. Specifically, win-
488 ter temperature is more variable in the north, which may lead to less predictability in the
489 start of the growing season or late frost. This could affect selection on phenology, which we
490 did not study here, but it is not obviously connected to the latitudinal cline in physiological
491 traits.

492 In summary, we found evidence for a coordinated latitudinal cline in germination rate,

493 photosynthesis, and growth, suggesting local adaptation. We therefore predict to find
 494 different optima for these traits in different climates. We did not find evidence that the
 495 relative performance of populations shifts with temperature or watering regime, suggesting
 496 relatively little variation in plasticity. Exploratory analysis implicate that more variable
 497 precipitation regimes at lower latitude drive much of the latitudinal cline, though other
 498 climatic factors could also contribute. Interestingly, the climatic neighborhood may shape
 499 selective pressures more than local climate. In the future, we will use field experiments to
 500 test whether greater variation in precipitation selects for faster growth and ~~that~~ selection
 501 on temperature/drought responses does not vary among populations. By doing so, we aim
 502 to understand why certain physiological and developmental mechanisms, but not others,
 503 contribute to local adaptation.

504 References

- 505 Adler, P. B., R. Salguero-Gómez, A. Compagnoni, J. S. Hsu, J. Ray-Mukherjee, C. Mbeau-
 506 Ache, and M. Franco, 2014. Functional traits explain variation in plant life history
 507 strategies. *Proceedings of the National Academy of Sciences of the United States of*
 508 *America* 111:740–745.
- 509 Angert, A. L., 2006. Demography of central and marginal populations of monkeyflowers
 510 (*Mimulus cardinalis* and *M. lewisii*). *Ecology* 87:2014–2025.
- 511 Angert, A. L., M. J. Bayly, S. N. Sheth, and J. R. Paul, 2016. Testing range-limit hypothe-
 512 ses using range-wide habitat suitability and occupancy for the scarlet monkeyflower
 513 (*Mimulus cardinalis*) .
- 514 Angert, A. L., H. Bradshaw Jr, and D. W. Schemske, 2008. Using experimental evolution
 515 to investigate geographic range limits in monkeyflowers. *Evolution* 62:2660–2675.
- 516 Barton, N. H., 1999. Clines in polygenic traits. *Genetics Research* 74:223–236.

- 517 Bates, D., M. Mächler, B. Bolker, and S. Walker, 2015. Fitting linear mixed-effects
518 models using lme4. Journal of Statistical Software 67:1–48. URL [http://CRAN.R-](http://CRAN.R-project.org/package=lme4)
519 [project.org/package=lme4](http://CRAN.R-project.org/package=lme4). R package version 1.1-7.
- 520 Blackman, B. K., S. D. Michaels, and L. H. Rieseberg, 2011. Connecting the sun to
521 flowering in sunflower adaptation. Molecular Ecology 20:3503–3512.
- 522 Bradshaw, H. and D. W. Schemske, 2003. Allele substitution at a flower colour locus
523 produces a pollinator shift in monkeyflowers. Nature 426:176–178.
- 524 Bradshaw, W. and C. Holzapfel, 2008. Genetic response to rapid climate change: it’s
525 seasonal timing that matters. Molecular ecology 17:157–166.
- 526 Bradshaw, W. E. and C. M. Holzapfel, 2001. Genetic shift in photoperiodic response corre-
527 lated with global warming. Proceedings of the National Academy of Sciences 98:14509–
528 14511.
- 529 Brown, J. S. and D. L. Venable, 1986. Evolutionary ecology of seed-bank annuals in
530 temporally varying environments. The American Naturalist 127:31–47.
- 531 Chevrud, J., 2001. A simple correction for multiple comparisons in interval mapping genome
532 scans. Heredity 87:52–58.
- 533 Clausen, J., D. Keck, and W. Hiesey, 1940. Experimental studies on the nature of species.
534 I. The effects of varied environments on western American plants, vol. 520. Carnegie
535 Institution of Washington, Washington, D.C.
- 536 Cleveland, W. S., E. Grosse, and W. M. Shyu, 1992. Local regression models. Statistical
537 models in S Pp. 309–376.
- 538 Coyne, J. A. and E. Beecham, 1987. Heritability of two morphological characters within
539 and among natural populations of *Drosophila melanogaster*. Genetics 117:727–737. URL
540 <http://www.genetics.org/content/117/4/727>.

541 Endler, J., 1977. Geographic variation, clines, and speciation. Princeton University Press,
542 Princeton, NJ.

543 Fox, J. and S. Weisberg, 2011. An R Companion to Applied Regression. Second ed. Sage,
544 Thousand Oaks CA.

545 Friedman, J. and M. J. Rubin, 2015. All in good time: Understanding annual and perennial
546 strategies in plants. *American Journal of Botany* 102:497–499.

547 Friedman, J., A. D. Twyford, J. H. Willis, and B. K. Blackman, 2015. The extent and
548 genetic basis of phenotypic divergence in life history traits in *Mimulus guttatus*. *Molecular*
549 *Ecology* 24:111–122.

550 Genuer, R., J.-M. Poggi, and C. Tuleau-Malot, 2015. VSURF: An R package for variable
551 selection using random forests. *The R Journal* 7:19–33.

552 ———, 2016. VSURF: Variable Selection Using Random Forests. URL [http://CRAN.R-](http://CRAN.R-project.org/package=VSURF)
553 [project.org/package=VSURF](http://CRAN.R-project.org/package=VSURF). R package version 1.0.3.

554 Gilbert, P., 2014. Brief User’s Guide: Dynamic Systems Estimation. URL [http://cran.r-](http://cran.r-project.org/web/packages/dse/vignettes/dse-guide.pdf)
555 [project.org/web/packages/dse/vignettes/dse-guide.pdf](http://cran.r-project.org/web/packages/dse/vignettes/dse-guide.pdf).

556 Givnish, T. J., 1988. Adaptation to sun and shade: a whole-plant perspective. *Functional*
557 *Plant Biology* 15:63–92.

558 Hadfield, J. D., 2010. Mcmc methods for multi-response generalized linear mixed mod-
559 els: The MCMCglmm R package. *Journal of Statistical Software* 33:1–22. URL
560 <http://www.jstatsoft.org/v33/i02/>.

561 ———, 2016. The spatial scale of local adaptation in a stochastic environment. *Ecology*
562 *Letters* 19:780–788.

563 Hereford, J., 2009. A quantitative survey of local adaptation and fitness trade-offs. *The*
564 *American Naturalist* 173:579–588.

565 Hereford, J. and A. A. Winn, 2008. Limits to local adaptation in six populations of the
 566 annual plant *Diodia teres*. *New Phytologist* 178:888–896.

567 Hiesey, W., M. Nobs, and O. Björkman, 1971. Experimental studies on the nature of
 568 species. V. Biosystematics, genetics, and physiological ecology of the Erythranthe section
 569 of *Mimulus*, vol. 628. Carnegie Institution of Washington, Washington, D.C.

570 Hiesey, W. M., J. Clausen, and D. D. Keck, 1942. Relations between climate and intraspe-
 571 cific variation in plants. *American Naturalist* Pp. 5–22.

572 Hijmans, R. J., S. Phillips, J. Leathwick, and J. Elith, 2016. dismo: Species distribution
 573 modeling. URL <http://CRAN.R-project.org/package=dismo>. R package version 1.1-1.

574 Hoffmann, A. A., A. Anderson, and R. Hallas, 2002. Opposing clines for high and low
 575 temperature resistance in *Drosophila melanogaster*. *Ecology Letters* 5:614–618.

576 Hopkins, R., J. Schmitt, and J. R. Stinchcombe, 2008. A latitudinal cline and response to
 577 vernalization in leaf angle and morphology in *Arabidopsis thaliana* (Brassicaceae). *New*
 578 *Phytologist* 179:155–164.

579 Huey, R. B., G. W. Gilchrist, M. L. Carlson, D. Berrigan, and L. Serra, 2000. Rapid
 580 evolution of a geographic cline in size in an introduced fly. *Science* 287:308–309.

581 Huxley, J. S., 1938. Clines: an auxiliary taxonomic principle. *Nature* 142:219–220.

582 Iwasa, Y. and D. Cohen, 1989. Optimal growth schedule of a perennial plant. *American*
 583 *Naturalist* 133:480–505.

584 Kawecki, T. J. and D. Ebert, 2004. Conceptual issues in local adaptation. *Ecology Letters*
 585 7:1225–1241.

586 Kooyers, N. J., 2015. The evolution of drought escape and avoidance in natural herbaceous
 587 populations. *Plant Science* 234:155–162.

588 Kooyers, N. J., A. B. Greenlee, J. M. Colicchio, M. Oh, and B. K. Blackman, 2015. Repli-

cate altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress in annual *Mimulus guttatus*. *New Phytologist* 206:152–165.

Kooyers, N. J. and K. M. Olsen, 2012. Rapid evolution of an adaptive cyanogenesis cline in introduced north american white clover (*Trifolium repens* L.). *Molecular Ecology* 21:2455–2468.

Kuznetsova, A., P. Bruun Brockhoff, and R. Haubo Bojesen Christensen, 2016. lmerTest: Tests in Linear Mixed Effects Models. URL <http://CRAN.R-project.org/package=lmerTest>. R package version 2.0-32.

Lehner, B., K. Verdin, and A. Jarvis, 2006. HydroSHEDS technical documentation. World Wildlife Fund, Washington, D.C. URL www.worldwildlife.org/hydrosheds.

Leimu, R. and M. Fischer, 2008. A meta-analysis of local adaptation in plants. *PLoS One* 3:e4010–e4010.

Lenormand, T., 2002. Gene flow and the limits to natural selection. *Trends in Ecology & Evolution* 17:183–189.

Levins, R., 1968. *Evolution in changing environments: some theoretical explorations*. Princeton University Press, Princeton, New Jersey.

Liaw, A. and M. Wiener, 2002. Classification and regression by randomforest. *R News* 2:18–22. URL <http://CRAN.R-project.org/doc/Rnews/>.

Lowry, D. B. and J. H. Willis, 2010. A widespread chromosomal inversion polymorphism contributes to a major life-history transition, local adaptation, and reproductive isolation. *PLoS biology* 8:2227.

Ludlow, M. M., 1989. Strategies of response to water stress. Pp. 269–281, in K. H. Kreeb, H. Richter, and T. M. Hinckley, eds. *Structural and functional responses to environmental stresses*. SPB Academic, The Hague, The Netherlands.

613 Nesom, G. L., 2014. Taxonomy of *Erythranthe* sect. *Erythranthe* (phrymaceae). *Phytoneu-*
614 *ron* 31:1–41.

615 Oakeshott, J., J. Gibson, P. Anderson, W. Knibb, D. Anderson, and G. Chambers, 1982.
616 Alcohol dehydrogenase and glycerol-3-phosphate dehydrogenase clines in *Drosophila*
617 *melanogaster* on different continents. *Evolution* Pp. 86–96.

618 Paul, J. R., T. L. Parchman, B. Econopouly, C. A. Buerkle, and A. L. Angert, In review.
619 Population genomics and range limits: diversity and differentiation across the geographic
620 range of *Mimulus cardinalis* (Phrymaceae) .

621 Paul, J. R., S. N. Sheth, and A. L. Angert, 2011. Quantifying the impact of gene flow
622 on phenotype-environment mismatch: A demonstration with the scarlet monkeyflower
623 *Mimulus cardinalis*. *The American Naturalist* 178:S62–S79.

624 Pfaff, B., 2008. VAR, SVAR and SVEC models: Implementation within R package vars.
625 *Journal of Statistical Software* 27:1–32.

626 Ronce, O. and M. Kirkpatrick, 2001. When sources become sinks: migrational meltdown
627 in heterogeneous habitats. *Evolution* 55:1520–1531.

628 Salguero-Gómez, R., O. R. Jones, E. Jongejans, S. P. Blomberg, D. J. Hodgson, C. Mbeau-
629 Ache, P. A. Zuidema, H. de Kroon, and Y. M. Buckley, 2016. Fast–slow continuum and
630 reproductive strategies structure plant life-history variation worldwide. *Proceedings of*
631 *the National Academy of Sciences of the United States of America* 113:230–235.

632 Sawada, S., Y. Nakajima, M. Tsukuda, K. Sasaki, Y. Hazama, M. Futatsuya, and
633 A. Watanabe, 1994. Ecotypic differentiation of dry matter production processes in re-
634 lation to survivorship and reproductive potential in *Plantago asiatica* populations along
635 climatic gradients. *Functional Ecology* 8:400–409.

636 Schemske, D. W., G. G. Mittelbach, H. V. Cornell, J. M. Sobel, and K. Roy, 2009. Is
637 there a latitudinal gradient in the importance of biotic interactions? *Annual Review of*

638 Ecology, Evolution, and Systematics 40:245–269.

639 Schluter, D., 2000. The Ecology of Adaptive Radiation. Oxford University Press.

640 Schmidt, P. S., L. Matzkin, M. Ippolito, and W. F. Eanes, 2005. Geographic variation in di-
641 apause incidence, life-history traits, and climatic adaptation in *Drosophila melanogaster*.
642 Evolution 59:1721–1732.

643 Sheth, S. N. and A. L. Angert, 2017. Demographic compensation does not rescue *Mimulus*
644 *cardinalis* populations at the southern edge of the species range. bioRxiv .

645 Slatkin, M., 1978. Spatial patterns in the distributions of polygenic characters. Journal of
646 Theoretical Biology 70:213–28.

647 Stearns, S., 1976. Life history tactics: a review of the ideas. Quarterly Review of Biology
648 51.

649 Stinchcombe, J. R., C. Weinig, M. Ungerer, K. M. Olsen, C. Mays, S. S. Halldorsdottir,
650 M. D. Purugganan, and J. Schmitt, 2004. A latitudinal cline in flowering time in *Ara-*
651 *bidopsis thaliana* modulated by the flowering time gene FRIGIDA. Proceedings of the
652 National Academy of Sciences of the United States of America 101:4712–4717.

653 Stock, A. J., B. E. Campitelli, and J. R. Stinchcombe, 2014. Quantitative genetic variance
654 and multivariate clines in the ivyleaf morning glory, *Ipomoea hederacea*. Philosophical
655 Transactions of the Royal Society B: Biological Sciences 369:20130259.

656 Templeton, A. R. and D. A. Levin, 1979. Evolutionary consequences of seed pools. The
657 American Naturalist 114:232–249.

658 Therneau, T. M., 2015. survival: A Package for Survival Analysis in S. URL
659 <http://CRAN.R-project.org/package=survival>. R package version 2.38.

660 Turesson, G., 1922. The genotypic response of the plant species to habitat. Hereditas
661 3:211–350.

662 Umina, P., A. Weeks, M. Kearney, S. McKechnie, and A. Hoffmann, 2005. A rapid shift
 663 in a classic clinal pattern in *Drosophila* reflecting climate change. *Science* 308:691–693.
 664 United States Geological Survey, 2015. National Hydrography Dataset. United States
 665 Geological Survey, Washington, D.C. URL <http://nhd.usgs.gov/index.html>.
 666 Wang, T., A. Hamann, D. L. Spittlehouse, and T. Q. Murdock, 2012. ClimateWNA –
 667 high-resolution spatial climate data for western North America. *Journal of Applied*
 668 *Meteorology and Climatology* 51:16–29.
 669 Wright, K. M., D. Lloyd, D. B. Lowry, M. R. Macnair, and J. H. Willis, 2013. Indirect
 670 evolution of hybrid lethality due to linkage with selected locus in *Mimulus guttatus*. *PLoS*
 671 *Biol* 11:e1001497.
 672 Wu, C., D. Lowry, A. Cooley, K. Wright, Y. Lee, and J. Willis, 2008. *Mimulus* is an
 673 emerging model system for the integration of ecological and genomic studies. *Heredity*
 674 100:220–230.
 675 Wu, C. A., D. B. Lowry, L. I. Nutter, and J. H. Willis, 2010. Natural variation for drought-
 676 response traits in the *Mimulus guttatus* species complex. *Oecologia* 162:23–33.
 677 Yeaman, S. and A. Jarvis, 2006. Regional heterogeneity and gene flow maintain variance
 678 in a quantitative trait within populations of lodgepole pine. *Proceedings of the Royal*
 679 *Society of London B* 273:1587–1593.




Figure 2: Overview of method for identifying putative climatic selective agents underlying latitudinal cline. We looked for climate variables that explained both the latitude of 356 *E. cardinalis* occurrences ('Climate-Latitude variables') and with traits ('Climate-Trait variables'). For Climate-Latitude variables we extracted climate data from recent occurrences located throughout California and Oregon, USA (shown in map). For Climate-Trait variables, we extracted climatic data for the 16 focal populations. For both analyses, we extracted local and neighborhood climate. Local climate refers to climate only from where a population was collected ($x_{i,0}$). Neighborhood climate was calculated as the average over 1000 points in a 62-km radius climatic neighborhood ($x_{i,1}, x_{i,2}, \dots$), but only along stream habitats as *E. cardinalis* is riparian. We identified climatic factors that most strongly predicted latitude of occurrences (Climate-Latitude variables) and traits (Climate-Trait variables), as shown for hypothetical data in plots at the bottom of the figure.




Figure 3: Trait variation, from fast to slow growth, is closely associated with latitude. Each point is a population's latitude of origin (x-axis) and position along the slow to fast growth axis (y-axis), defined as Principal Component 1 of four traits (see Material and Methods). The line and 95% confidence intervals were estimated using linear regression.




Figure 4: Climatic variation integrated over climatic neighborhood is closely correlated with latitude of *E. cardinalis* and trait variation. A. Using Random Forest regression, we identified 16 climatic variables significantly (high importance) associated with latitude of *E. cardinalis* occurrences. B. Only one of the most important Climate-Latitude variables (in bold) was among the most important Climate-TraitPC1 variables. Variable importance is defined as the average amount a climate variable reduces mean-squared error in the predicted response (TraitPC1 or Latitude), compared to a randomly permuted dataset, across all trees in the random forest (see Genuer et al. [2015] for further detail). Note that the Importance values in A and B are not comparable because the dependent variables (Latitude and Trait PC1, respectively) are on different scales. Climatic variables (left of A; right of B) are defined by four qualities: Climatic factor – Temperature (Temp), Precipitation (Prec), Heating degree-days (Heating), Snow (precipitation as snow); Temporal scale – Annual, Coldest quarter (Cold Quar), Warmest Quarter (Warm Quar), Wettest quarter (Wet Quar), Driest Quarter (Dry Quar), or Seasonality; Type – 30-year average (Avg.) or coefficient of variation (Var.); Spatial scale – local or 62-km radius climatic neighborhood.




Figure 5: Variation in precipitation is correlated with TraitPC1 and latitude. A. Greater values of TraitPC1 are associated with greater interannual variation in precipitation of the wettest quarter. This was the most important Climate-Latitude variable, but not among the most important Climate-TraitPC1 variables. B. However, a closely related parameter, interannual variation in precipitation seasonality, was among the most important Climate-TraitPC1 variables. C. Across focal populations, variation in precipitation of the wettest quarter and seasonality are closely correlated. D. Southern populations of *E. cardinalis* experience much greater interannual variation in precipitation. In all panels, we report climatic neighborhood values (see Material and Methods). Regression lines, 95% confidence intervals, and coefficients of determination (R^2) were calculated using linear regression.

680 **Supporting Information**

681 **Supporting Tables**

Table S1: Initial size of seedlings did not vary among Populations, Families, or Treatments. We used a censored Gaussian model of initial size at the outset of the experiment (longest leaf length of the first true leaves). The model was censored because we could not accurately measure leaves less than 0.25 mm with digital callipers (217 of 702, 30.9%, were too small). We fit models using a Bayesian MCMC method implemented using the MCMCglmm function with default priors in the R package **MCMCglmm** version 2.17 (Hadfield, 2010). We estimated the posterior distribution from 1000 samples of an MCMC chain run for 10^5 steps after a 10^4 step burn-in. We used step-wise backward elimination procedure to find the best-supported model according to Deviance Information Criterion (DIC).

Model	Random	DIC
Population + Water + Temperature + Population:Water + Population:Temperature + Water:Temperature + Population:Water:Temperature	Family	1638
Population + Water + Temperature + Population:Water + Population:Temperature + Water:Temperature	Family	1605.2
Population + Water + Temperature + Population:Water + Population:Temperature	Family	1603.4
Population + Water + Temperature + Population:Water + Water:Temperature	Family	1577.5
Population + Water + Temperature + Population:Temperature + Water:Temperature	Family	1579.9
Population + Water + Temperature + Population:Water	Family	1577.3
Population + Water + Temperature + Water:Temperature	Family	1550.5
Population + Water + Temperature	Family	1549.3
Population + Water	Family	1541.7
Population + Temperature	Family	1546.8
Water + Temperature	Family	1551.1
Population	Family	1541.9
Water	Family	1543.9
-	Family	1541.7
-	-	1538.3

Table S2: Climatic variables used

Abbreviation	Climate variable
DD_0	degree-days below 0°C(chilling degree-days)
DD5	degree-days above 5°C(growing degree-days)
DD_18	degree-days below 18°C(heating degree-days)
DD18	degree-days above 18°C(cooling degree-days)
NFFD	number of frost-free days
PAS	precipitation as snow (mm) between August in previous year and July in current
Eref	Hargreaves reference evaporation (mm)
CMD	Hargreaves climatic moisture deficit (mm)
RH	mean annual relative humidity
bio1	annual mean temperature
bio2	mean diurnal range (mean of monthly (max temp - min temp))
bio3	isothermality (bio2/bio7) (* 100)
bio4	temperature seasonality (standard deviation *100)
bio5	max temperature of warmest month
bio6	min temperature of coldest month
bio7	temperature annual range (bio5-bio6)
bio8	mean temperature of wettest quarter
bio9	mean temperature of driest quarter
bio10	mean temperature of warmest quarter
bio11	mean temperature of coldest quarter
bio12	annual precipitation
bio15	precipitation seasonality (coefficient of variation)
bio16	precipitation of wettest quarter
bio17	precipitation of driest quarter
bio18	precipitation of warmest quarter
bio19	precipitation of coldest quarter

Table S3: Analysis of variance (ANOVA) table on leaf expansion rate (LER) using **lmerTest** (Kuznetsova et al., 2016). Family and Block were included as random effects. Abbreviations: SS = sum of squares; MS = mean sum of squares (SS / df1); df1 = numerator degrees of freedom; df2 = denominator degrees of freedom.

	SS	MS	df1	df2	<i>F-value</i>	<i>P-value</i>
Day to Germination	12.12	12.12	1	637	35.21	4.9×10^{-9}
Population	22.22	1.48	15	118	4.3	2.5×10^{-6}
Temperature	80.42	80.42	1	5	233.61	2.6×10^{-5}
Water	4.1	4.1	1	5	11.92	0.019
Temperature \times Water	0.03	0.03	1	4	0.07	0.801
Population \times Temperature	2.76	0.18	15	547	0.53	0.925
Population \times Water	9.66	0.64	15	562	1.87	0.024
Population \times Temperature \times Water	4.11	0.27	15	530	0.78	0.700

Table S4: Analysis of variance (ANOVA) table on stem elongation rate (SER) using **lmerTest** (Kuznetsova et al., 2016). Family and Block were included as random effects. Abbreviations: SS = sum of squares; MS = mean sum of squares (SS / df1); df1 = numerator degrees of freedom; df2 = denominator degrees of freedom.

	SS	MS	df1	df2	<i>F-value</i>	<i>P-value</i>
Day to Germination	3.6	3.6	1	662	21.1	5.1×10^{-6}
Population	12	0.8	15	113	4.7	5.8×10^{-7}
Temperature	12.4	12.4	1	6	72.8	1.5×10^{-4}
Water	0.6	0.6	1	5	3.7	0.113
Temperature \times Water	0.9	0.9	1	4	5.2	0.093
Population \times Temperature	3.6	0.2	15	549	1.4	0.126
Population \times Water	2.8	0.2	15	536	1.1	0.330
Population \times Temperature \times Water	1.5	0.1	15	518	0.6	0.874

Table S5: Analysis of variance (ANOVA) table on photosynthetic rate using **lmerTest** (Kuznetsova et al., 2016). Family and Block were included as random effects. Abbreviations: SS = sum of squares; MS = mean sum of squares (SS / df1); df1 = numerator degrees of freedom; df2 = denominator degrees of freedom.

	SS	MS	df1	df2	<i>F-value</i>	<i>P-value</i>
Population	347.7	23.2	15	78	3.02	7.5×10^{-4}
Temperature	134.1	134.1	1	6	17.46	6.4×10^{-3}
Water	51	51	1	4	6.64	0.066
Temperature \times Water	0.7	0.7	1	3	0.09	0.781
Population \times Temperature	218.6	14.6	15	263	1.9	0.024
Population \times Water	87.7	5.8	15	233	0.76	0.724
Population \times Temperature \times Water	91.4	6.1	15	208	0.79	0.686

Table S6: Analysis of deviance table on the probability of mortality by the end of the experiment using Type-II Wald χ^2 tests in the R package **car** (Fox and Weisberg, 2011). Family and Block were included as random effects. Abbreviations: df = degrees of freedom

	χ^2	df	<i>P-value</i>
Population	32	31	0.419
Temperature	31.8	6	1.8×10^{-5}
Water	69.2	12	4.6×10^{-10}
Temperature \times Water	20.7	1	5.3×10^{-6}
Population \times Temperature	5.6	15	0.985
Population \times Water	8.6	15	0.897
Population \times Temperature \times Water	0.2	15	1.000

Table S7: Important climatic variables predicting latitude of *E. cardinalis* populations ('Climate-Latitude') and the first principal component of traits measured in a common garden ('Climate-TraitPC1'). Local climatic variables were measured from the exact location of collection; neighborhood climatic variables were averaged from a 62-km neighborhood around population (see Material and Methods). Importance and significance were determined using the variable selection using random forests (VSURF) algorithm (see Material and Methods). Climatic variables are described in Table S2. μ signifies the mean of the climate variables from 1981–2010; σ indicates coefficient of variation among years.

Climate-Latitude variables	Climate-TraitPC1 variables
Precipitation of wettest quarter (σ , neighborhood)	Mean diurnal range (σ , neighborhood)
Annual precipitation (μ , neighborhood)	Precipitation seasonality (σ , neighborhood)
Precipitation of wettest quarter (μ , neighborhood)	Mean temperature of coldest quarter (σ , neighborhood)
Mean temperature of coldest quarter (σ , neighborhood)	
Annual precipitation (σ , neighborhood)	
Precipitation of driest quarter (μ , neighborhood)	
Precipitation of coldest quarter (σ , neighborhood)	
Hargreaves climatic moisture deficit (μ , neighborhood)	
Precipitation of warmest quarter (μ , neighborhood)	
Precipitation seasonality (μ , neighborhood)	
Precipitation of warmest quarter (σ , neighborhood)	
Heating degree-days (σ , neighborhood)	
Precipitation of driest quarter (σ , neighborhood)	
Number of frost-free days (σ , neighborhood)	
Mean temperature of wettest quarter (σ , neighborhood)	
Precipitation as snow (σ , neighborhood)	




Figure S1: Southern populations germinate faster. Each point is a population of *E. cardinalis* showing its latitude of origin (x-axis) and model-predicted days to germination in days under growth chamber conditions (see Material and Methods). Bars around each point are 95% confidence intervals. Predicted time to germination and confidence intervals are based on survival regression (see Materials and Materials). The line is the linear regression of $\log(\text{model-predicted days to germination}) \sim \text{latitude}$. The P -value of the regression is given in the upper left corner.




Figure S2: Southern populations grow faster. Each point is a population of *E. cardinalis* showing its latitude of origin (x-axis) and model-predicted leaf expansion rate during the rosette phase. Bars around each point are 95% confidence intervals. Predicted leaf expansion rate based least-square mean estimates and confidence intervals were calculated from linear mixed-effects models (see Materials and Materials). The line is the linear regression of model-predicted leaf expansion rate \sim latitude. The P -value of the regression is given in the upper right corner.




Figure S3: Southern populations grow faster. Each point is a population of *E. cardinalis* showing its latitude of origin (x-axis) and model-predicted stem elongation rate. Bars around each point are 95% confidence intervals. Predicted stem elongation rate based least-square mean estimates and confidence intervals were calculated from linear mixed-effects models (see Materials and Materials). The line is the linear regression of model-predicated stem elongation rate \sim latitude. The P -value of the regression is given in the upper right corner.




Figure S4: Southern populations photosynthesize faster. Each point is a population of *E. cardinalis* showing its latitude of origin (x-axis) and model-predicted instantaneous photosynthetic rate. Bars around each point are 95% confidence intervals. Predicted photosynthetic rates based least-square mean estimates and confidence intervals were calculated from linear mixed-effects models (see Materials and Materials). The line is the linear regression of model-predicted photosynthetic rate \sim latitude. The P -value of the regression is given in the upper right corner.




Figure S5: Reaction norms signify little Population \times Treatment interactions. For all panels, black lines represent population-level reaction norms from Wet to Dry in the Cool temperature treatment (dashed black lines) and Hot temperature treatment (solid black lines); gray lines represent reaction norms from Cool to Hot in the Wet treatment (solid gray lines) and Dry treatment (dashed gray lines). The responses shown are (A) leaf expansion rate, (B) stem elongation rate, (C) photosynthesis, and (D) survivorship (= 1 - mortality).




Figure S6: Neighborhood climate predicts TraitPC1 ('Climate-trait', panel A) and Latitude of occurrences ('Climate-latitude', panel B) better than local climate. Each point is the absolute value of the Pearson correlation coefficient ($|r|$) between TraitPC1 (A) or latitude (B) for 24 climatic factors, for which we used both the 30-year mean (closed circles) and coefficient of variation (open circles). Most points lie above the 1:1 line, indicating stronger correlations with neighborhood compared to local climate. Neighborhood climate was integrated over a 62-km radius around focal populations (see text for further detail).




Figure S7: Trait variation, from fast to slow growth, is closely associated with neighborhood variation in temperature of the coldest quarter (bio11_σ). Each point is a population coefficient of variation in bio11 averaged over a 62-km climatic neighborhood (x-axis) and position along the slow to fast growth axis (y-axis), defined as Principal Component 1 of four traits (see Material and Methods). The line and 95% confidence intervals were estimated using linear regression.

683 Supporting Material and Methods

684 Temperature treatments

685 We simulated typical growing season (June 1 - August 15) air temperatures at the two most
686 thermally divergent focal sites in our study, Whitewater Canyon (WWC, Hot) and Little
687 Jameson (LIJ, Cool). We downloaded daily interpolated mean, minimum, and maximum
688 air temperature from 13 years (2000-2012) at both sites from ClimateWNA (Wang et al.,
689 2012). This range was chosen because seeds used in the experiment were collected around
690 2012, thus their presence in that location at that time suggests that populations were able
691 to persist there for at least some years before collection. Monthly temperatures from Cli-
692 mateWNA are highly correlated with the air temperature recorded from data loggers in
693 the field at these sites (A. Angert, unpub. data). Hence, the ClimateWNA temperature
694 profiles are similar to actual thermal regimes experienced by *E. cardinalis* in nature. We
695 simulated realistic temperature regimes by calculating the mean temperature trend from
696 June to August using LOESS (Cleveland et al., 1992). The residuals were highly autocor-
697 related at both sites (warmer than average days are typically followed by more warm days)
698 and there was strong correlation ($r = 0.65$) between sites (warm days in WWC were also
699 warm in LIJ). The ‘VARselect’ function in the **vars** package for R (Pfaff, 2008) indicated
700 that a lag two Vector Autoregression (VAR(2)) model best captured the within-site auto-
701 correlation as well as between-site correlation in residuals. We fit and simulated from the
702 VAR(2) model using the package **dse** (Gilbert, 2014) in R. Simulated data closely resem-
703 bled the autocorrelation and between-site correlation of the actual data. From simulated
704 mean temperature, we next selected minimum and maximum daily temperatures. Mean,
705 min, and max temperature were highly correlated at both sites. We chose min and max
706 temperatures using site-specific fitted linear models between mean, max, and min tem-
707 perature, with additional variation given by normally distributed random deviates with
708 variance equal to the residual variance of the linear models. For each day, the nighttime

709 (22:00 - 6:00) chamber temperature was set to the simulated minimum temperature. Dur-
710 ing the middle of the day, temperature was set to the simulated maximum temperature,
711 with a variable period of transition between min and max so that the average temperature
712 was equal the simulated mean temperature.

713 **Watering treatments**

714 For watering treatments, we simulated two extreme types of streams where *E. cardinalis*
715 grows. In the well-watered treatment, we simulated a large stream that never goes dry
716 during the summer growing season. In the drought treatment, we simulated a small stream
717 that has ample flow at the beginning of the season due to rain and snow melt, but gradually
718 dries down through the summer. In both treatments, plants were bottom-watered using
719 water chilled to 7.5°C. Plants in the well-watered treatment were fully saturated every two
720 hours during the day. Watering in the drought treatment gradually declined from every
721 two hours to every day between May 20 (36 days after sowing) and 10 June (57 days after
722 sowing). Simultaneously, the amount of bottom-watering per flood decreased, such that
723 only the bottom of the cone-tainers were wetted by the end of the experiment.