

Light and growth form interact to shape stomatal ratio among British angiosperms

Christopher D. Muir¹

¹ Biodiversity Research Centre and Botany Department, University of British Columbia, Vancouver, British Columbia V6T 1Z4, Canada

Author for correspondence:

Christopher D. Muir

Tel: +17782284851

Email: chrisdmuir@gmail.com

University of British Columbia

6270 University Blvd.

Vancouver, BC, Canada

V6T 1Z4

Short title: Shedding light on stomatal evolution

Word count:

Summary: 190

Introduction: 1179

Materials and Methods: 1176

Results: 433

Discussion: 1504

Acknowledgement: 24

3 Figures and 1 Table, 1 Supplemental Figure

1 Summary

- 2 • In most plants, stomata are located only on the abaxial leaf surface (hypos-
3 tomy), but many plants have stomata on both surfaces (amphistomy). High
4 light and herbaceous growth form have been hypothesized to favor amphis-
5 tomy, but these hypotheses have not been rigourously tested together using
6 phylogenetic comparative methods.
- 7 • I leveraged a large dataset including stomatal ratio, Ellenberg light indica-
8 tor value, Raunkiær lifeform, and phylogenetic relationships for 372 species of
9 British angiosperms. I used phylogenetic comparative methods to test how
10 light and/or growth form influence stomatal ratio.
- 11 • High light and herbaceous growth form are correlated with amphistomy, as
12 predicted, but they also interact; the effect of light is pronounced in therophytes
13 (annuals) and perennial herbs, but muted in phanerophytes (mostly trees).
14 Interestingly, amphistomy and stomatal density evolve together in response to
15 light, suggesting coordinated selection on this trait combination.
- 16 • I show for the first time that light and growth form interact to shape variation
17 in stomatal ratio; amphistomy is advantageous in high light, but mostly for
18 herbs. These results improve our understanding of the adaptive significance of
19 stomatal ratio as well as its use as functional trait for paleoecology and crop
20 improvement.

21 **Keywords**

22 Adaptation, amphistomy, Ellenberg light indicator value, growth form, phylogenetic
23 comparative methods, Raunkiær lifeform, stomata, stomatal ratio

24 **Introduction**

25 Natural selection shapes leaf anatomy in order to optimize its photosynthetic function
26 in a given environment (Haberlandt, 1914; Givnish, 1987; Smith et al., 1997). By
27 understanding the adaptive significance of leaf anatomical variation we can learn
28 about natural history, find targets for crop improvement, and identify anatomical
29 proxies for paleoclimates preserved in the fossil record (e.g. Wolfe, 1971; Royer, 2001;
30 McElwain and Steinthorsdottir, 2017). The size, density, and distribution of stomata
31 on a leaf vary widely and impact the flux of CO₂ and water vapour (recently reviewed
32 in Sack and Buckley, 2016), as well as susceptibility to foliar pathogens that infect
33 through stomata (McKown et al., 2014; Melotto et al., 2017). Hence, stomata have
34 been especially useful in understanding plastic and evolutionary response to climate
35 change and domestication (Woodward, 1987; Beerling and Royer, 2011; Milla et al.,
36 2013).

37 While the density and size of stomata have been researched extensively (Sack and
38 Buckley, 2016, and references therein), the adaptive significance of stomatal distri-
39 bution is less well understood. Stomata are most often found only on the lower
40 leaf surface (hypostomy) but occur on both surfaces (amphistomy) in many species
41 (Metcalfe and Chalk, 1950; Parkhurst, 1978; Mott et al., 1984). Theory and ex-
42 periments demonstrate that amphistomy increases photosynthetic rates under many

43 conditions. By creating a second parallel pathway for CO₂ diffusion within the meso-
44 phyll, amphistomy optimally supplies CO₂ (Parkhurst, 1978; Gutschick, 1984; Jones,
45 1985). Amphistomy is correlated with greater CO₂ diffusion (Beerling and Kelly,
46 1996) and higher photosynthetic rates (McKown et al., 2014). These observations
47 are corroborated by experiments demonstrating that amphistomy increases maxi-
48 mum photosynthetic rates by up to 20% (Parkhurst and Mott, 1990). On the other
49 hand, amphistomy can increase transpiration (Jones, 1985; Foster and Smith, 1986;
50 Buckley et al., 2015). While transition to amphistomy is thus thought to increase
51 transpiration, empirical studies suggest greater water-use efficiency in amphistoma-
52 tous species (Bucher et al., 2017). Hence, amphistomy appears to benefit a plant's
53 carbon use relative to water loss and should be favored when CO₂ limits photo-
54 synthetic rate. The open questions are under what ecological conditions does CO₂
55 supply most strongly limit photosynthetic rate (Peat and Fitter, 1994) and when is
56 photosynthetic rate most important to fitness?

57 The leading, nonmutually exclusive hypotheses are that 1) open habitats favour
58 amphistomy because CO₂ diffusion most strongly limits photosynthetic rate under
59 high light and 2) herbaceous growth form favours amphistomy because traits that
60 maximize photosynthetic rate are often under stronger selection in herbs. Salisbury
61 (1927) first noted that amphistomy is most common in herbaceous plants from open
62 habitats (i.e., with high light) of the British flora. These observations have been
63 replicated in other studies (Mott et al., 1984; Peat and Fitter, 1994; Jordan et al.,
64 2014; Muir, 2015) and may support physiological and ecological hypotheses that CO₂
65 most strongly limits photosynthesis in high light and/or photosynthesis contributes
66 most to fitness in herbaceous plants. Under high light, CO₂ can strongly limit max-
67 imum photosynthetic rates, especially in thick leaves (Jones, 1985). Hence, having

68 stomata on both surfaces relieves this limitation by adding a second parallel pathway
69 for CO₂ diffusion. Parkhurst (1978) argued that greater leaf thickness *per se* selected
70 for amphistomy, but there is little evidence for correlations between leaf thickness
71 and stomatal ratio independent of light (Mott et al., 1984; Gibson, 1996; Muir, 2015).
72 Amphistomy is correlated with open habitat in warm desert plants of western North
73 America (Mott et al., 1984; Gibson, 1996), among the Proteaceae (Jordan et al.,
74 2014), and in continental European herbs (Bucher et al., 2017).

75 Stomatal ratio is also associated with growth form. In the British flora, Salisbury
76 (1927) found that trees and shrubs are nearly always hypostomatous, whereas herbs
77 from open habitats are amphistomatous. This pattern holds when data are averaged
78 by family to coarsely control for phylogenetic nonindependence (Peat and Fitter,
79 1994) or when using alternative classification schemes, such as Raunkiaer life form
80 (Peat and Fitter, 1994). Across plants from ~ 90 families worldwide, growth form
81 is the strongest predictor of stomatal ratio when multiple factors are estimated si-
82 multaneously and controlling for phylogenetic nonindependence (Muir, 2015). These
83 patterns are consistent with other data indicating that many herbaceous plants are
84 under strong selection for high maximum photosynthetic rates (Bazzaz, 1979; Körner
85 et al., 1989; Wullschleger, 1993).

86 Although previous comparative studies have tested whether open habitat and growth
87 form influence stomatal ratio, we do not know if these effects are independent of one
88 another. Open habitat and growth form may not be independent because open
89 habitats generally consist of more short-statured, herbaceous plants. Some authors
90 have attempted to disentangle light and growth form by contrasting herbs from
91 open and understory habitats (Salisbury, 1927). However, this is problematic if
92 phylogenetic relationships are not controlled for, because shade species may share

93 traits simply because they are more closely related to each other than they are to
94 high light species. Finally, open habitat and growth form may also interact with one
95 another. For example, amphistomy may only be favored when CO₂ strongly limits
96 photosynthetic rate (e.g. in high light) *and* photosynthetic rate strongly limits fitness
97 (e.g. in herbs).

98 To better understand the adaptive significance of stomatal ratio, I asked three main
99 questions:

- 100 1. Are light habitat and growth form correlated?
- 101 2. Do light habitat and growth form influence stomatal ratio additively, or do
102 their effects interact?
- 103 3. Is evolution of stomatal ratio mediated by changes in stomatal density on the
104 adaxial (upper) surface, abaxial (lower) surface, or both?

105 The final question is important for addressing whether amphistomy is part of a co-
106 ordinated syndrome of traits that promote higher photosynthetic rate, as both the
107 light and growth form hypotheses assume. If evolved increases in stomatal ratio are
108 mediated by shifting abaxial stomata to the adaxial surface, holding total stomatal
109 density constant, then the overall increase in CO₂ diffusion would be small. In con-
110 trast, if amphistomy evolves by increasing adaxial stomatal density while holding
111 abaxial density constant, then *total* stomatal density must increase as well. Evolu-
112 tionary coordination of amphistomy and high stomatal density would reinforce one
113 another, increasing CO₂ supply to chloroplasts more than changes in either trait
114 would in isolation.

115 To address these questions, I reanalyzed existing data on stomatal ratio, light habi-
116 tat, and growth form in British angiosperms (Salisbury, 1927; Fitter and Peat, 1994,

2017) using phylogenetic comparative methods. The British angiosperm flora is well suited for these questions because this flora has been comprehensively surveyed for many ecologically important traits, meaning it is probably the least biased survey of stomatal trait variation. Salisbury's observations on stomata and ecology in the British flora have heavily influenced plant ecophysiology, but many of his and subsequent authors' analyses have significant limitations because of inadequate statistical methods. For example, few analyses until recently account for phylogenetic nonindependence (Felsenstein, 1985), which can strongly influence inferences on stomatal traits and growth form (Kelly and Beerling, 1995, this study did not consider light). A species-level phylogeny of the entire British flora (Lim et al., 2014) now allows for the first time rigorous analysis of evolutionary relationships among stomatal ratio, light, and growth form.

Materials and Methods

Data and annotated source code to generate this manuscript are available on GitHub (<https://github.com/cdmuir/britstom>) and Dryad (Muir, 2017).

Data on stomatal ratio, light habitat, growth form, and phylogenetic relationships

I obtained data on ab- and adaxial stomatal density on 395 species from British Ecological Flora (Salisbury, 1927; Fitter and Peat, 1994, 2017). Following recent comparative analyses (e.g. Niinemets and Valladares, 2006; Bartelheimer and Poschlod, 2016; Shipley et al.), I used Ellenberg light indicator values (Ellenberg, 1974) as mea-

138 sures of light habitat. Hence, I am assuming that the species' light habitat is closely
139 related to the type of habitat (open versus closed) where that species is found. Ellen-
140 berg light indicator values, hereafter abbreviated L-value, have been recently updated
141 by taxonomic experts of the British flora (PLANTATT, Hill et al. (2004)).

142 There is no universally adopted scientific classification scheme for plant growth form,
143 therefore I statistically compared two widely used schemes. First, I used PLAN-
144 TATT data on perennation, woodiness, and height to classify species' growth form
145 as herbaceous (annual, biennial, or perennial) or woody (shrub or tree). Following
146 Muir (2015), 'biennial' includes true biennials as well as species that have a mix
147 of perennation forms (e.g. a species with both annual and perennial forms would
148 be classified as a biennial here). Woody species are shrubs (plant height less than
149 4 m) or trees (plant height greater than 4 m). Next, I compared this scheme to
150 PLANTATT data on Raunkiær life form (Raunkiær, 1934), which is another way to
151 classify growth form in comparative ecology (e.g. Peat and Fitter, 1994; Salguero-
152 Gómez et al., 2016). I retained phanerophytes, geophytes, chamaephytes, hemicryp-
153 tophytes, and therophytes, but excluded data on hydrophytes (14 species) because
154 many of these species are hyperstomatous (Fig. S1) due to the fact that leaves may
155 rest on the water's surface, selecting for stomata to be present on the upper surface
156 only. The two main differences between these growth form classifications are that
157 1) most shrubs and trees are lumped together as phanerophytes and 2) many geo-
158 phytes and chamaephytes are lumped together with hemicryptophytes as perennials
159 (Fig. S2).

160 I used a dated molecular phylogeny of the British flora (Lim et al., 2014) available
161 from TreeBASE (<http://treebase.org/>; accession number 15105). 14 species (3.5%)
162 in the dataset were not present in the phylogeny. For 8 of these species, I used the

163 position of a congeneric species as a proxy for the focal species (following Pennell
164 et al., 2016). When multiple congeneric species were present, I consulted the phy-
165 logenetic literature to identify the most closely related proxy species (Scheen et al.,
166 2004; Salmaki et al., 2013). For the remaining 6 missing species, I positioned them
167 in the tree based on phylogenetic relationships to other genera or families present in
168 the tree (Fior et al., 2006). Because many phylogenetic comparative methods do not
169 allow polytomies, zero-length branches, and non-ultrametric trees, I made several
170 small adjustments to the tree. I resolved polytomies randomly using the ‘multi2di’
171 function in **phytools** version 0.6-00 (Revell, 2012). I added 0.02 my to all zero-length
172 branches, as this was approximately the length of the shortest nonzero branch length
173 in the tree. After these changes, I slightly altered terminal branch lengths to make
174 the tree precisely ultrametric.

175 I excluded C₄ (3 species) and CAM (2 species) plants. I limited this investigation to
176 angiosperms because only 4 non-angiosperms had stomata data. The final dataset
177 contained 372 species (Fig. 1). The R code accompanying this paper documents
178 these decisions in greater detail and citations to the relevant literature.

179 Following Muir (2015), I calculated stomatal ratio in two different ways depending
180 on what was most appropriate for the question:

$$SR_{propAd} = \frac{SD_{ad}}{SD_{total}} \quad (1)$$

$$SR_{even} = \frac{\min\{SD_{ab}, SD_{ad}\}}{\max\{SD_{ab}, SD_{ad}\}} \quad (2)$$

181 SD_{ab} and SD_{ad} are the stomatal densities on abaxial or adaxial surface, respectively.

182 $SD_{total} = SD_{ab} + SD_{ad}$. SR_{propAd} is the proportion of stomata density on the adaxial
 183 surface, which is useful for discriminating among hypostomatous ($SR_{propAd} = 0$),
 184 amphistomatous ($0 < SR_{propAd} < 1$), and hyperstomatous species ($SR_{propAd} = 1$).
 185 SR_{even} indicates how evenly stomatal densities are distributed across both leaf sur-
 186 faces. This expression is useful because several hypotheses are based on the fact that
 187 a more even distribution should optimize leaf CO_2 diffusion.

188 **Testing for an association between open habitat and growth** 189 **form**

190 I tested whether growth form, under either classification scheme, was associated
 191 with L-value among British angiosperms. I predicted that species with faster life
 192 histories, especially therophytes (annuals), would have greater L-values than species
 193 with slower life histories, especially phanerophytes (shrubs and trees). I first used
 194 a phylogenetic ANOVA assuming an Ornstein-Uhlenbeck process model fit using
 195 **phyloIm** version 2.5 (Ho and Ané, 2014). However, this analysis indicated no phylo-
 196 genetic signal in the regression (See the R code accompanying this paper for further
 197 detail). Specifically, the estimated α parameter was extremely high. In the Ornstein-
 198 Uhlenbeck model, α is proportional to the inverse of the phylogenetic half-life (i.e.
 199 phylogenetic signal). When there is no phylogenetic signal (i.e. high α), regular and
 200 phylogenetic ANOVA converge on the same parameters estimates. Furthermore, sta-
 201 tistical tests assuming there is phylogenetic signal when in fact none exists performs
 202 worse than nonphylogenetic tests (Revell, 2010). Therefore, I used a regular ANOVA
 203 with Type-2 sum of squares.

204 **Open habitat, growth form, and stomatal ratio**

205 I compared phylogenetic linear models to test whether growth form, L-value, or
206 interactions between them predicted SR_{even} . Unlike the analysis above, there was
207 significant phylogenetic signal in this comparison (see R code). I used SR_{even} rather
208 than SR_{propAd} as the response variable because the hypothesis is that faster life his-
209 tory and/or high light favor more even stomatal densities on each surface. I fit
210 models using **phylolm** and extracted Akaike Information Criteria (AIC). For these
211 and subsequent analyses, I assumed an Ornstein-Uhlenbeck process model for the
212 residuals with the root character state integrated over the stationary distribution.
213 I used a 10^4 parametric bootstrap samples of the full model (including main ef-
214 fects and interactions) to calculate parameter confidence intervals (Boettiger et al.,
215 2012).

216 **Does ab- or adaxial stomatal density contribute more to stom-** 217 **atal ratio evolution?**

218 I used two related phylogenetic methods, variance decomposition and structural equa-
219 tion modeling (SEM), to assess the relative contribution of ab- versus adaxial stom-
220 atal density to light-mediated stomatal ratio evolution. First, the contribution of ab-
221 versus adaxial stomatal density can be calculated using phylogenetic variance de-
222 composition methods as derived below. Because stomatal density is highly skewed,
223 I log-transformed values for normality:

$$SR_{\text{even}} = \frac{SD_{\text{ad}}}{SD_{\text{ab}}} \quad (3)$$

$$\log(\text{SR}_{\text{even}}) = \log(\text{SD}_{\text{ad}}) - \log(\text{SD}_{\text{ab}}) \quad (4)$$

$$\text{sr}_{\text{even}} = \text{sd}_{\text{ab}} - \text{sd}_{\text{ad}} \quad (5)$$

224 Lowercase variables (sr, sd) indicate log-transformed values. Because some species
 225 had zero adaxial stomata, I added one to all values prior to log-transformation. To
 226 make the variance decomposition calculations tractable, I have defined SR_{even} here
 227 as the ratio of ad- to abaxial stomatal density because in most cases adaxial stomatal
 228 density is lower than abaxial (see Eq. 2). This differs from analyses described above
 229 because in those I wanted to test what factors influenced the evenness of stomatal
 230 densities, regardless of which surface had higher density. With this modified form,
 231 the variance in sr_{even} can readily be decomposed into contributions of sd_{ad} , sd_{ab} , and
 232 their covariance:

$$\text{Var}(\text{sr}_{\text{even}}) = \text{Var}(\text{sd}_{\text{ad}}) + \text{Var}(\text{sd}_{\text{ab}}) - 2\text{Cov}(\text{sd}_{\text{ad}}, \text{sd}_{\text{ab}}) \quad (6)$$

233 I did not use the raw covariance, but rather estimated the phylogenetic covariance
 234 matrix between L-value, sd_{ab} , and sd_{ad} using a multivariate Ornstein-Uhlenbeck
 235 model fit in **Rphylopars** version 0.2.9 (Goolsby et al., 2016, 2017). From the co-
 236 variance matrix, I estimated the contribution of abaxial density, adaxial density, and
 237 their covariance as:

$$\text{Contribution of } \text{sd}_{\text{ad}} = \frac{\text{Var}(\text{sd}_{\text{ad}})}{\text{Var}(\text{sr}_{\text{even}})} \quad (7)$$

$$\text{Contribution of } sd_{ab} = \frac{\text{Var}(sd_{ab})}{\text{Var}(sr_{\text{even}})} \quad (8)$$

$$\text{Contribution of } \text{Cov}(sd_{ad}, sd_{ab}) = \frac{\text{Cov}(sd_{ad}, sd_{ab})}{\text{Var}(sr_{\text{even}})} \quad (9)$$

238 respectively. Note that when ab- and adaxial densities positively covary, the contri-
 239 bution will be negative because this reduces the variance in stomatal ratio.

240 I also wanted to test whether light-mediated evolution of stomatal ratio acted mostly
 241 by 1) increasing adaxial stomatal density while maintaining abaxial density, or 2)
 242 keeping total stomatal density the same, but shifting a greater proportion to the adax-
 243 ial surface. The first scenario predicts that the phylogenetic regression of L-value on
 244 sd_{ad} is stronger than that for sd_{ab} . The second scenario predicts that L-value acts sim-
 245 ilarly on both and that there is a negative covariance ($\text{Cov}(sd_{ad}, sd_{ab}) < 0$). I tested
 246 these competing predictions by fitting a very simple phylogenetic SEM (see Mason
 247 et al., 2016, for a similar approach). The model uses the phylogenetic covariance
 248 matrix, as described above, to simultaneously estimate regressions of L-value on sd_{ad}
 249 and sd_{ab} while allowing covariance between them (i.e. estimating $\text{Cov}(sd_{ad}, sd_{ab})$).
 250 To fit the SEM, I used the R package **lavaan** version 0.5-23.1097 (Rosseel, 2012).
 251 I tested whether parameter estimates were significantly different from zero using
 252 z -scores.

253 Results

254 Light tolerance varies among Raunkiær life forms

255 Ellenberg light indicator values (L-value) differed significantly among life forms (Fig. 2; ANOVA
256 - $F_{4,367} = 18.3$, $P = 1.1 \times 10^{-13}$). Therophytes (annuals), hemicryptophytes (peren-
257 nial herbs with buds near the soil surface), and chamaephytes (subshrubs) had greater
258 L-values than phanerophytes (large woody plants) and geophytes (perennial herbs
259 with storage organs) (Fig. 2).

260 Interactions between light and Raunkiær life form determine 261 stomatal ratio

262 Overall, SR_{even} increased with L-value, but there was a significant interaction ($\Delta AIC >$
263 2, Table 1) between Raunkiær life form and L-value (Fig. 3). Both life form and L-
264 value significantly increased model fit, though L-value had a markedly larger effect
265 on model AIC (Table 1). The significant interaction is caused by different slopes
266 between life forms. Among life forms with the overall greatest L-value (therophytes,
267 hemicryptophytes, and chamaephytes, see Fig. 2), there was a strong positive rela-
268 tionship between L-value and SR_{even} . Parametrically bootstrapped 95% confidence
269 intervals for the slope did not overlap zero (Fig. 3). The slope was weakly positive
270 or not significantly different from zero in the most shade-adapted life forms (geo-
271 phytes and phanerophytes), albeit the patterns were distinct in these groups. There
272 were both hypostomatous ($SR_{\text{even}} \approx 0$) and amphistomatous ($SR_{\text{even}} \approx 1$) geophytes,
273 but these were distributed across L-values. In contrast, phanerophytes were nearly

274 always hypostomatous regardless of L-value.

275 **Adaxial stomatal density contributes most of the variation in** 276 **stomatal ratio**

277 Adaxial ('upper') stomatal density contributed most to the evolutionary variation
278 in stomatal ratio. The contributions of adaxial density, abaxial density, and their
279 covariance are 1.12, 0.38, and -0.5, respectively. This implies that evolutionary varia-
280 tion in adaxial stomatal density is greater than that for stomatal ratio due to positive
281 covariance between ab- and adaxial stomatal density.

282 Similarly, the phylogenetic SEM showed that changes in stomatal ratio associated
283 with L-value can be attributed mostly to evolution of adaxial stomatal density
284 (Fig. 4). Both sd_{ad} and sd_{ab} increased with L-value ($P = 1.2 \times 10^{-8}$ and 8.9×10^{-7} ,
285 respectively). However, the regression of L-value on sd_{ad} was $2\times$ that of L-value on
286 sd_{ab} (0.24 versus 0.12). Because stomatal densities were natural log-transformed, this
287 implies an increase in L-value by one leads to a 1.27-fold change in adaxial stom-
288 atal density versus a 1.13-fold change in abaxial stomatal density. The SEM also
289 showed a significant positive covariance between stomatal densities on each surface
290 ($P = 2.5 \times 10^{-10}$). These results together imply that total stomatal density increases
291 with L-value, but the response is mediated mostly by increases in adaxial stomatal
292 density.

Discussion

The ratio of stomatal densities on the abaxial (‘lower’) to that of the adaxial (‘upper’) surface varies greatly across plant species, but the adaptive significance is not clear. Comparative studies correlating stomatal ratio to ecological factors can distinguish among competing hypotheses and reveal critical experiments for future work. Previous comparative studies suggested that high light and herbaceous growth form favor amphistomy (Mott et al., 1984; Jordan et al., 2014; Muir, 2015; Bucher et al., 2017), particularly in the British flora (Salisbury, 1927; Peat and Fitter, 1994). However, none of these studies have accounted for the fact that light and growth form are often confounded – open, high light habitats are often dominated by herbs – or the fact that species are not independent because of shared evolutionary history. Here, I reanalyzed data on stomata, light tolerance, and growth form in British angiosperms using phylogenetic comparative methods. As expected, species’ light tolerance (Ellenberg light indicator or L-value) is confounded with growth form (Raunkiær life form; Fig. 2). Nevertheless, both L-value and Raunkiær life form affect stomatal ratio, but these factors also interact; the influence of L-value on stomatal ratio varies across forms. These novel findings provide further evidence that variation in stomatal ratio is adaptive and have important implications for interpreting changes in stomatal ratio through the paleo record (Jordan et al., 2014) and during domestication (Milla et al., 2013).

Adaptive significance of amphistomy

Previously, associations between light, growth form, and stomatal ratio have been interpreted in isolation as indicating that either high light and/or herbaceous growth

316 form favors amphistomy. In British angiosperms, both factors are important, though
317 statistical analyses suggest that light may be a stronger determinant than growth
318 form (Table 1). Unlike previous studies, I found a significant interaction between
319 light and growth form among British angiosperms, which suggests that amphistomy
320 may only be strongly favored when CO₂ strongly limits photosynthesis (as in open
321 habitat) *and* photosynthesis strongly limits fitness (as in herbs). This is consistent
322 with life history theory predicting that the demography of open habitat herbs is
323 strongly limited by plant growth (Franco and Silvertown, 1996). The ideal way to
324 test this would be to measure selection on stomatal ratio in a species that varied
325 quantitatively in both stomatal ratio and life history (e.g., containing both annual
326 and perennial forms). I predict that amphistomy will be favored more strongly in
327 the annual form grown under high light compared to an annual under low light
328 or a perennial in high light, and much more strongly than a perennial grown in low
329 light. Similar experiments could also be performed to test if and when light-mediated
330 plasticity in stomatal ratio is adaptive (Gay and Hurd, 1975; Mott and Michaelson,
331 1991; Fontana et al., 2017).

332 The prevalence of amphistomatous species in high light habitats supports the hy-
333 pothesis that amphistomy is an adaptation to maximize photosynthetic rates by
334 increasing CO₂ diffusion (Jones, 1985). It is also evidence against the hypothesis
335 that the principle fitness cost of amphistomy is water loss (Darwin, 1886; Foster and
336 Smith, 1986) or dehydration of palisade mesophyll (Buckley et al., 2015), though
337 these factors are likely very important in determining differential regulation of stom-
338 ata on each surface. Since evaporative demand increases under high light, under
339 these hypotheses we would expect plants in high light to be hypostomatous. Because
340 stomatal conductances on each surface can be regulated independently in response

341 to the environment (Darwin, 1898; Pospíšilová and Solárová, 1984; Smith, 1981; Re-
 342 ich, 1984; Mott and O’Leary, 1984), amphistomatous leaves likely cope with these
 343 stresses by rapidly closing adaxial stomata when water supply cannot match evapo-
 344 rative demands (Richardson et al., 2017). Instead, patterns in the British flora are
 345 at least consistent with the idea that adaxial stomata increase susceptibility to foliar
 346 pathogens (Gutschick, 1984; McKown et al., 2014). The cost of adaxial stomata
 347 may be greater in the shade because greater leaf wetness and lower ultraviolet light
 348 provide a more suitable microclimate for many foliar pathogens.

349 **Amphistomy as a proxy for open habitat**

350 These observations from the British flora partially support the hypothesis that am-
 351 phistomy can be used a proxy for open habitat in paleoenvironment reconstruction
 352 (Carpenter, 1994; Jordan et al., 2014; Carpenter et al., 2015) but also point out pre-
 353 viously unknown subtleties. These previous studies based their conclusions on data
 354 from Proteaceae, in which there is little quantitative variation in stomatal ratio;
 355 species are either completely hypostomatous ($SR_{propAd} \approx 0$) or completely amphis-
 356 tomatous ($SR_{propAd} \approx 0.5$). Stomatal ratio in British angiosperms is also bimodal
 357 (Peat and Fitter, 1994), but across many families there is also quantitative variation.
 358 Importantly, this means that quantitative variation in stomatal ratio may provide a
 359 more precise, quantitative indicator of vegetation type, rather than simply ‘open’ or
 360 ‘closed’. A quantitative relationship between L-value and stomatal ratio has already
 361 been shown for herbaceous perennials (Bucher et al., 2017), but we now know that
 362 it holds among annuals (therophytes), subshrubs (chamaephytes), and, to a lesser
 363 extent, geophytes as well (Fig. 3).

364 The weak or nonsignificant relationship between L-value and stomatal ratio in geo-
 365 phytes and phanerophytes suggests that in some cases amphistomy may not reliably
 366 indicate open habitat without further information. For example, perhaps amphis-
 367 tomatous geophytes from partially shaded habitats are spring ephemerals, so they
 368 experience high light during their growth phase, but this has not been tested. Like-
 369 wise, phanerophytes (most tall trees) are almost always hypostomatous (see also
 370 Muir (2015)). Most British phanerophytes are tall, hypostomatous trees, but the
 371 exceptions are telling. For example, the most amphistomatous phanerophyte in this
 372 dataset is *Brassica oleracea*, a short-statured biennial that has more in common
 373 physiologically with hemicryptophytes than other phanerophytes. The other am-
 374 phistomatous phanerophytes in this data set (*Populus nigra* and *Lavatera arborea*)
 375 are fast-growing pioneer species.

376 Finally, phylogenetic information should improve inferences about paleoclimates be-
 377 cause there is appreciable phylogenetic signal in stomatal ratio. The phylogenetic
 378 half-life of stomatal ratio evolution, after accounting for L-value and Raunkiær life
 379 form, is $\log(2)/\alpha = 1.5$ my (see Table 1 for maximum likelihood estimates of α , the
 380 return rate in the Ornstein-Uhlenbeck model of trait evolution). This lag time may
 381 indicate that evolving to the ‘optimum’ is constrained by the shape of the fitness
 382 landscape (Muir, 2015) or that other unmeasured factors which affect stomatal ra-
 383 tio have some phylogenetic signal. Regardless of the mechanism, this fact means
 384 that researchers may be able to use data from closely related species to improve
 385 paleoenvironment reconstruction.

386 **Why does adaxial stomatal density control stomatal ratio?**

387 Variation in stomatal ratio is determined primarily by evolution of adaxial stom-
388 atal density and is coordinated with increases in total leaf stomatal density summed
389 across both surfaces. Note here that I am referring only to evolutionary variation in
390 stomatal ratio among species; different processes may mediate within species vari-
391 ation or plastic responses. Phylogenetic analyses show that changes in stomatal
392 ratio and total stomatal density, especially in response to L-value, are predominantly
393 mediated by changes in adaxial stomatal density. This highly nonrandom pattern
394 among British angiosperms mirrors evolutionary changes wrought by domestication
395 (Milla et al., 2013); crop species tend to have higher adaxial stomatal density than
396 their wild relatives.

397 There are at least two hypotheses that could explain why adaxial stomatal density
398 is the most responsive. The first I refer to as the ‘real estate’ hypothesis. In hy-
399 postomatous plants, the lower surface is already crowded with stomata, and hence
400 plants must increase the real estate available for stomata by developing them on the
401 upper surface whenever there is selection for greater stomatal density. When stomata
402 are packed too densely on one surface, stomatal interference limits their function-
403 ing and hence may create a strong selective pressure for amphistomy (Parlange and
404 Waggoner, 1970; Dow et al., 2014).

405 I refer to the second hypothesis as the ‘coordination’ hypothesis. In this scenario,
406 ecological conditions such as high light select for both increased total stomatal density
407 and for amphistomy because these traits work well in coordination with one another.
408 For example, if stomatal density were very high on a hypostomatous plant, then CO₂
409 would be more strongly limited by the mesophyll. Adding a second parallel pathway

410 for diffusion by developing stomata on both surfaces would restore a more optimal
411 balance between stomatal and mesophyll limitations. Conversely, there would be
412 little benefit to amphistomy when total stomatal density is low because CO₂ diffusion
413 is strongly limited by stomatal resistance, and therefore photosynthetic rate is not
414 sensitive to changes in mesophyll diffusion mediated by stomatal ratio. A related
415 prediction is that increased atmospheric CO₂ may select for reduced stomatal ratio
416 and density primarily by decreasing adaxial stomatal density, but this has not been
417 well tested (but see Woodward and Bazzaz, 1988).

418 **Conclusions**

419 By revisiting this classic ecological dataset with modern phylogenetic comparative
420 methods, I have shown that amphistomy is strongly associated with both light and
421 growth form, but the interaction between these factors is also important. Fur-
422 thermore, amphistomy and high stomatal density are closely connected in species
423 from high light environments, suggesting selection for coordination between these
424 traits.

425 **Acknowledgements**

426 I thank Sally Otto, Matt Pennell, and Rob Salguero-Gómez for feedback on this
427 manuscript. I was supported by an NSERC CREATE grant.

428 Author contribution statement

429 CDM designed the study, collected data, analyzed the data, and wrote the manuscript.

References

- Bartelheimer, M. and P. Poschlod, 2016. Functional characterizations of Ellenberg indicator values—a review on ecophysiological determinants. *Functional Ecology* 30:506–516.
- Bazzaz, F., 1979. The physiological ecology of plant succession. *Annual Review of Ecology and Systematics* 10:351–71.
- Beerling, D. J. and C. K. Kelly, 1996. Evolutionary comparative analyses of the relationship between leaf structure and function. *New Phytologist* 134:35–51.
- Beerling, D. J. and D. L. Royer, 2011. Convergent Cenozoic CO₂ history. *Nature Geoscience* 4:418–420.
- Boettiger, C., G. Coop, and P. Ralph, 2012. Is your phylogeny informative? Measuring the power of comparative methods. *Evolution* 66:2240–2251.
- Bucher, S. F., K. Auerswald, C. Grün-Wenzel, S. I. Higgins, J. G. Jorge, and C. Römermann, 2017. Stomatal traits relate to habitat preferences of herbaceous species in a temperate climate. *Flora* 229:107–115.
- Buckley, T. N., G. P. John, C. Scoffoni, and L. Sack, 2015. How does leaf anatomy influence water transport outside the xylem? *Plant Physiology* 168:1616–1635.
- Carpenter, R. J., 1994. Cuticular morphology and aspects of the ecology and fossil history of North Queensland rainforest Proteaceae. *Botanical Journal of the Linnean Society* 116:249.
- Carpenter, R. J., M. K. Macphail, G. J. Jordan, and R. S. Hill, 2015. Fossil evidence

451 for open, Proteaceae-dominated heathlands and fire in the Late Cretaceous of
 452 Australia. *American Journal of Botany* 102:2092–2107.

453 Darwin, F., 1886. On the relation between the “bloom” on leaves and the distribution
 454 of the stomata. *Botanical Journal of the Linnean Society* 22:99–116.

455 ———, 1898. Observations on stomata. *Philosophical Transactions of the Royal*
 456 *Society B: Biological Sciences* 190:531–621.

457 Dow, G. J., J. A. Berry, and D. C. Bergmann, 2014. The physiological importance
 458 of developmental mechanisms that enforce proper stomatal spacing in *Arabidopsis*
 459 *thaliana*. *New Phytologist* 201:1205–1217.

460 Ellenberg, H., 1974. Indicator values of vascular plants in central Europe, *Scripta*
 461 *Geobotanica*, vol. 9. Springer-Verlag, Göttingen, Germany.

462 Felsenstein, J., 1985. Phylogenies and the comparative method. *The American*
 463 *Naturalist* 1:1–15.

464 Fior, S., P. O. Karis, G. Casazza, L. Minuto, and F. Sala, 2006. Molecular phylogeny
 465 of the Caryophyllaceae (Caryophyllales) inferred from chloroplast matk and nuclear
 466 rDNA ITS sequences. *American Journal of Botany* 93:399–411.

467 Fitter, A. and H. Peat, 1994. The ecological flora database. *Journal of Ecology*
 468 82:415–425.

469 ———, 2017. Ecological flora of the British isles. URL
 470 <http://www.ecoflora.co.uk>.

471 Fontana, M., M. Labrecque, A. Collin, and N. Bélanger, 2017. Stomatal distribu-
 472 tion patterns change according to leaf development and leaf water status in *Salix*
 473 *miyabeana*. *Plant Growth Regulation* 81:63–70.

- 474 Foster, J. and W. Smith, 1986. Influence of stomatal distribution on transpiration
475 in low-wind environments. *Plant, Cell & Environment* 9:751–759.
- 476 Franco, M. and J. Silvertown, 1996. Life history variation in plants: an exploration
477 of the fast-slow continuum hypothesis. *Philosophical Transactions: Biological Sci-*
478 *ences* 351:1341–1348.
- 479 Gay, A. and R. Hurd, 1975. The influence of light on stomatal density in the tomato.
480 *New Phytologist* 75:37–46.
- 481 Gibson, A. C., 1996. *Structure-Function Relations of Warm Desert Plants*. Springer-
482 Verlag, Berlin.
- 483 Givnish, T. J., 1987. Comparative studies of leaf form: assessing the relative roles
484 of selective pressures and phylogenetic constraints. *New Phytologist* 106:131–160.
- 485 Goolsby, E. W., J. Bruggeman, and C. Ané, 2016. Rphylopars: Phyloge-
486 netic Comparative Tools for Missing Data and Within-Species Variation. URL
487 <https://CRAN.R-project.org/package=Rphylopars>. R package version 0.2.9.
- 488 ———, 2017. Rphylopars: fast multivariate phylogenetic comparative methods for
489 missing data and within-species variation. *Methods in Ecology and Evolution*
490 8:22–27.
- 491 Gutschick, V. P., 1984. Photosynthesis model for C₃ leaves incorporating CO₂ trans-
492 port, propagation of radiation, and biochemistry 2. ecological and agricultural
493 utility. *Photosynthetica* 18:569–595.
- 494 Haberlandt, G., 1914. *Physiological Plant Anatomy*. Macmillan and Co., London.
- 495 Hill, M., C. Preston, and D. Roy, 2004. *PLANTATT - Attributes of British and Irish*

496 Plants: Status, Size, Life History, Geography and Habitats. Centre for Ecology &
497 Hydrology, Huntingdon, Cambridgeshire.

498 Ho, L. S. T. and C. Ané, 2014. Intrinsic inference difficulties for trait evolution with
499 Ornstein-Uhlenbeck models. *Methods in Ecology and Evolution* 5:1133–1146.

500 Jones, H. G., 1985. Adaptive significance of leaf development and structural responses
501 to environment. Pp. 155–173, *in* N. R. Baker, W. Davies, and C. K. Ong, eds.
502 Control of Leaf Growth, *Society for Experimental Biology Seminar Series*, vol. 27.
503 Cambridge University Press, Cambridge.

504 Jordan, G. J., R. J. Carpenter, and T. J. Brodribb, 2014. Using fossil leaves as
505 evidence for open vegetation. *Palaeogeography, Palaeoclimatology, Palaeoecology*
506 395:168–175.

507 Kelly, C. and D. Beerling, 1995. Plant life form, stomatal density and taxonomic
508 relatedness: a reanalysis of Salisbury (1927). *Functional Ecology* 9:422–431.

509 Körner, C., M. Neumayer, S. P. Menendez-Riedl, and A. Smeets-Scheel, 1989. Func-
510 tional morphology of mountain plants. *Flora* 182:353–383.

511 Lim, J., M. J. Crawley, N. De Vere, T. Rich, and V. Savolainen, 2014. A phylogenetic
512 analysis of the British flora sheds light on the evolutionary and ecological factors
513 driving plant invasions. *Ecology and Evolution* 4:4258–4269.

514 Mason, C. M., E. W. Goolsby, D. P. Humphreys, and L. A. Donovan, 2016. Phy-
515 logenetic structural equation modelling reveals no need for an ‘origin?’ of the leaf
516 economics spectrum. *Ecology letters* 19:54–61.

517 McElwain, J. C. and M. Steinthorsdottir, 2017. Paleoecology, ploidy, paleoatmo-

518 spheric composition, and developmental biology: a review of the multiple uses of
519 fossil stomata. *Plant Physiology* 174:650–664.

520 McKown, A. D., R. D. Guy, L. Quamme, J. Klápště, J. La Mantia, C. Constabel,
521 Y. A. El-Kassaby, R. C. Hamelin, M. Zifkin, and M. Azam, 2014. Association
522 genetics, geography and ecophysiology link stomatal patterning in *Populus tri-*
523 *chocarpa* with carbon gain and disease resistance trade-offs. *Molecular Ecology*
524 23:5771–5790.

525 Melotto, M., L. Zhang, P. R. Oblessuc, and S. Y. He, 2017. Stomatal defense a
526 decade later. *Plant Physiology* 174:561–571.

527 Metcalfe, C. R. and L. Chalk, 1950. *Anatomy of the dicotyledons*, Vols. 1 & 2. First
528 ed. Oxford University Press, Oxford.

529 Milla, R., N. de Diego-Vico, and N. Martín-Robles, 2013. Shifts in stomatal traits
530 following the domestication of plant species. *Journal of Experimental Botany*
531 64:3137–3146.

532 Mott, K. A., A. C. Gibson, and J. W. O’Leary, 1984. The adaptive significance of
533 amphistomatic leaves. *Plant, Cell & Environment* 5:455–460.

534 Mott, K. A. and O. Michaelson, 1991. Amphistomy as an adaptation to high light
535 intensity in *Ambrosia cordifolia* (Compositae). *American Journal of Botany* 78:76–
536 79.

537 Mott, K. A. and J. W. O’Leary, 1984. Stomatal behavior and CO₂ exchange char-
538 acteristics in amphistomatous leaves. *Plant Physiology* 74:47–51.

539 Muir, C. D., 2015. Making pore choices: repeated regime shifts in stomatal ratio.
540 *Proc. R. Soc. B* 282:20151498.

- 541 ———, 2017. Data from: Light and life form interact to shape stomatal ratio among
542 British angiosperms. URL <http://dx.doi.org/10.5061/dryad.?????>
- 543 Niinemets, Ü. and F. Valladares, 2006. Tolerance to shade, drought, and waterlog-
544 ging of temperate Northern Hemisphere trees and shrubs. *Ecological Monographs*
545 76:521–547.
- 546 Parkhurst, D. F., 1978. The adaptive significance of stomatal occurrence on one or
547 both surfaces of leaves. *The Journal of Ecology* 66:367–383.
- 548 Parkhurst, D. F. and K. A. Mott, 1990. Intercellular diffusion limits to CO₂ uptake
549 in leaves studied in air and helox. *Plant Physiology* 94:1024–1032.
- 550 Parlange, J.-Y. and P. E. Waggoner, 1970. Stomatal dimensions and resistance to
551 diffusion. *Plant Physiology* 46:337–342.
- 552 Peat, H. and A. Fitter, 1994. A comparative study of the distribution and density of
553 stomata in the British flora. *Biological Journal of the Linnean Society* 52:377–393.
- 554 Pennell, M. W., R. G. FitzJohn, and W. K. Cornwell, 2016. A simple approach for
555 maximizing the overlap of phylogenetic and comparative data. *Methods in Ecology*
556 and *Evolution* 7:751–758.
- 557 Pospíšilová, J. and J. Solárová, 1984. Environmental and biological control of diffu-
558 sive conductances of adaxial and abaxial leaf epidermes. *Photosynthetica* 18:445–
559 453.
- 560 Raunkiaer, C. C., 1934. *The Life Forms of Plants and Statistical Plant Geography*.
561 Clarendon Press, Oxford.
- 562 Reich, P., 1984. Relationships between leaf age, irradiance, leaf conductance, CO₂
563 exchange, and water-use efficiency in hybrid poplar. *Photosynthetica* 18:445–453.

- 564 Revell, L. J., 2010. Phylogenetic signal and linear regression on species data. *Methods*
565 *in Ecology and Evolution* 1:319–329.
- 566 ———, 2012. phytools: An R package for phylogenetic comparative biology (and
567 other things). *Methods in Ecology and Evolution* 3:217–223.
- 568 Richardson, F., T. J. Brodribb, and G. J. Jordan, 2017. Amphistomatic leaf sur-
569 faces independently regulate gas exchange in response to variations in evaporative
570 demand. *Tree Physiology* Pp. 1–10.
- 571 Rosseel, Y., 2012. lavaan: An R package for structural equation modeling. *Journal*
572 *of Statistical Software* 48:1–36.
- 573 Royer, D. L., 2001. Stomatal density and stomatal index as indicators of paleoatmo-
574 spheric CO₂ concentration. *Review of Palaeobotany and Palynology* 114:1–28.
- 575 Sack, L. and T. N. Buckley, 2016. The developmental basis of stomatal density and
576 flux. *Plant Physiology* 171:2358–2363.
- 577 Salguero-Gómez, R., O. R. Jones, E. Jongejans, S. P. Blomberg, D. J. Hodgson,
578 C. Mbeau-Ache, P. A. Zuidema, H. de Kroon, and Y. M. Buckley, 2016. Fast–
579 slow continuum and reproductive strategies structure plant life-history variation
580 worldwide. *Proceedings of the National Academy of Sciences of the United States*
581 *of America* 113:230–235.
- 582 Salisbury, E., 1927. On the causes and ecological significance of stomatal frequency,
583 with special reference to the woodland flora. *Philosophical Transactions of the*
584 *Royal Society of London. Series B* 216:1–65.
- 585 Salmaki, Y., S. Zarre, O. Ryding, C. Lindqvist, C. Bräuchler, G. Heubl, J. Barber,

586 and M. Bendiksby, 2013. Molecular phylogeny of tribe Stachydeae (Lamiaceae
587 subfamily Lamioideae). *Molecular Phylogenetics and Evolution* 69:535–551.

588 Scheen, A.-C., C. Brochmann, A. K. Brysting, R. Elven, A. Morris, D. E. Soltis, P. S.
589 Soltis, and V. A. Albert, 2004. Northern hemisphere biogeography of *Cerastium*
590 (Caryophyllaceae): insights from phylogenetic analysis of noncoding plastid nu-
591 cleotide sequences. *American Journal of Botany* 91:943–952.

592 Shipley, B., M. Belluau, I. Kühn, N. A. Soudzilovskaia, M. Bahn, J. Penue-
593 las, J. Kattge, L. Sack, J. Cavender-Bares, W. A. Ozinga, B. Blonder, P. M.
594 van Bodegom, P. Manning, T. Hickler, E. Sosinski, V. D. P. Pillar, and
595 V. Onipchenko, ????. Predicting habitat affinities of plant species using com-
596 monly measured functional traits. *Journal of Vegetation Science* Pp. n/a–n/a.
597 URL <http://dx.doi.org/10.1111/jvs.12554>.

598 Smith, W., 1981. Temperature and water relation patterns in subalpine understory
599 plants. *Oecologia* 48:353–359.

600 Smith, W. K., T. C. Vogelmann, E. H. DeLucia, D. T. Bell, and K. A. Shepherd,
601 1997. Leaf form and photosynthesis. *BioScience* 11:785–793.

602 Wolfe, J. A., 1971. Tertiary climatic fluctuations and methods of analysis of Tertiary
603 floras. *Palaeogeography, Palaeoclimatology, Palaeoecology* 9:27–57.

604 Woodward, F., 1987. Stomatal numbers are sensitive to increases in CO₂ from pre-
605 industrial levels. *Nature* 327:617–618.

606 Woodward, F. I. and F. Bazzaz, 1988. The responses of stomatal density to CO₂
607 partial pressure. *Journal of Experimental Botany* 39:1771–1781.

608 Wullschleger, S. D., 1993. Biochemical limitations to carbon assimilation in C₃

609 plants? A retrospective analysis of the A/Ci curves from 109 species. Journal
610 of Experimental Botany 44:907–920.

Table 1: Interaction between species' Ellenberg light indicator value (L-value) and Raunkiaer lifeform predict stomatal ratio (SR_{even}). I compared phylogenetic linear models using the Akaike Information Criterion (AIC), where $AIC = 2k - 2\log(\mathcal{L})$. k is the number of model parameters and \mathcal{L} is the model likelihood. Given a set of candidate models, the difference in AIC between a model and the lowest AIC (ΔAIC) indicates the relative fit of competing models. The correlation coefficient r^2 is another indicator of model fit. α and σ^2 are the return rate and diffusion parameters of the Ornstein-Uhlenbeck model of trait evolution.

Model: $SR_{\text{even}} \sim$	α	σ^2	r^2	k	$\log(\mathcal{L})$	AIC	ΔAIC
L-value \times Raunkiaer lifeform	0.46	0.068	0.34	12	-33.2	90.4	0
L-value \times growth form	0.46	0.07	0.32	12	-38.2	100.4	9.9
L-value + Raunkiaer lifeform	0.46	0.071	0.32	8	-40.2	96.4	6
L-value + growth form	0.51	0.08	0.31	8	-43.4	102.7	12.3
Raunkiaer lifeform	0.34	0.067	0.15	7	-79.2	172.4	82
growth form	0.35	0.069	0.13	7	-82.5	179.1	88.6
L-value	0.64	0.107	0.26	4	-59.3	126.6	36.2
null	0.29	0.067	0	3	-107.6	221.1	130.7




Figure 1: CAPTION




Figure 2: Lifeforms have different tolerances for sun and shade among British angiosperms. Each panel is the distribution of Ellenberg light indicator values on an integer scale of 1-9 for different Raunkiær life forms. Height of the bars indicate the raw proportion of species in each bin for that lifeform. The sample size for each lifeform is listed next in parentheses. The mean (open circle) and 95% confidence intervals (black line) around the mean Ellenberg light indicator value for each lifeform based on phylogenetic regression are above the histogram.




Figure 3: The effect of light on stomatal ratio depends on Raunkiaer life form. Greater Ellenberg light indicator values (L-value) are associated with greater stomatal ratio (SR_{even}) in therophytes, hemicryptophytes, and chamaephytes but not geophytes and phanerophytes. The maximum likelihood slope from phylogenetic regression is given with statistical significance based on 10⁴ parametric bootstrap samples. Numbers in parentheses next to Raunkiaer life form are the sample sizes in the final dataset. Estimated slopes (solid line) and 95% bootstrapped confidence intervals (gray polygon between dashed lines) are plotted against raw data. Points have been jittered for visual clarity.




Figure 4: Light-mediated evolution of stomatal ratio is mostly driven by increased adaxial (‘upper’) stomatal density (Panel A), whereas abaxial (‘lower’) stomatal density (Panel B) is similar across Ellenberg light indicator values (L-value x -axis). The violin plot shows stomatal density (y -axis, log-scale) as a function of L-value. The width of the grey polygons indicates the density of data. Length of grey polygon indicate the range of the data; the dot indicates the median; the thick lines indicate the 0.25 and 0.75 quantiles. Points outside the polygons are statistical outliers.

611 Supporting Information




Figure S1: Most hydrophytes are hyperstomatous, having most stomata on the adaxial ('upper') surface (high SD_{propAd}). The violin plot shows stomatal ratio as a function of Raunkiaer lifeform. The width of the grey polygons indicates the density of data. Length of grey polygon indicate the range of the data; the point indicates the median; the thick lines indicate the 0.25 and 0.75 quantiles. Sample sizes per lifeform in the dataset are given above the upper plot margin. SD_{ad} and SD_{total} stand for the stomatal density on the adaxial surface and the total leaf surface (adaxial plus abaxial), respectively.




Figure S2: CAPTION




Figure S3: Growth forms have different tolerances for sun and shade among British angiosperms. Each panel is the distribution of Ellenberg light indicator values on an integer scale of 1-9 for different growth forms. Height of the bars indicate the raw proportion of species in each bin for that lifeform. The sample size for each lifeform is listed next in parentheses. The mean (open circle) and 95% confidence intervals (black line) around the mean Ellenberg light indicator value for each lifeform based on phylogenetic regression are above the histogram.