

Light and growth form interact to shape stomatal ratio among British angiosperms

Christopher D. Muir¹

¹ Biodiversity Research Centre and Botany Department, University of British Columbia, Vancouver, British Columbia V6T 1Z4, Canada

Author for correspondence:

Christopher D. Muir

Tel: +17782284851

Email: chrisdmuir@gmail.com

University of British Columbia

6270 University Blvd.

Vancouver, BC, Canada

V6T 1Z4

Short title: Shedding light on stomatal evolution

Word count:

Summary: 190

Introduction: 1179

Materials and Methods: 1176

Results: 433

Discussion: 1504

Acknowledgement: 24

3 Figures and 1 Table, 1 Supplemental Figure

1 Summary

- 2 • In most plants, stomata are located only on the abaxial leaf surface (hypos-
3 tomy), but many plants have stomata on both surfaces (amphistomy). High
4 light and herbaceous growth form have been hypothesized to favor amphis-
5 tomy, but these hypotheses have not been rigourously tested together using
6 phylogenetic comparative methods.
- 7 • I leveraged a large dataset including stomatal ratio, Ellenberg light indica-
8 tor value, Raunkiær lifeform, and phylogenetic relationships for 372 species of
9 British angiosperms. I used phylogenetic comparative methods to test how
10 light and/or growth form influence stomatal ratio.
- 11 • High light and herbaceous growth form are correlated with amphistomy, as
12 predicted, but they also interact; the effect of light is pronounced in therophytes
13 (annuals) and perennial herbs, but muted in phanerophytes (mostly trees).
14 Interestingly, amphistomy and stomatal density evolve together in response to
15 light, suggesting coordinated selection on this trait combination.
- 16 • I show for the first time that light and growth form interact to shape variation
17 in stomatal ratio; amphistomy is advantageous in high light, but mostly for
18 herbs. These results improve our understanding of the adaptive significance of
19 stomatal ratio as well as its use as functional trait for paleoecology and crop
20 improvement.

21 **Keywords**

22 Adaptation, amphistomy, Ellenberg light indicator value, growth form, phylogenetic
23 comparative methods, Raunkiær lifeform, stomata, stomatal ratio

24 **Introduction**

25 Natural selection shapes leaf anatomy in order to optimize its photosynthetic function
26 in a given environment (Haberlandt, 1914; Givnish, 1987; Smith et al., 1997). By
27 understanding the adaptive significance of leaf anatomical variation we can learn
28 about natural history, find targets for crop improvement, and identify anatomical
29 proxies for paleoclimates preserved in the fossil record (e.g. Wolfe, 1971; Royer, 2001;
30 McElwain and Steinthorsdottir, 2017). The size, density, and distribution of stomata
31 on a leaf vary widely and impact the flux of CO₂ and water vapour (recently reviewed
32 in Sack and Buckley, 2016), as well as susceptibility to foliar pathogens that infect
33 through stomata (McKown et al., 2014; Melotto et al., 2017). Hence, stomata have
34 been especially useful in understanding plastic and evolutionary response to climate
35 change and domestication (Woodward, 1987; Beerling and Royer, 2011; Milla et al.,
36 2013).

37 While the density and size of stomata have been researched extensively (Sack and
38 Buckley, 2016, and references therein), the adaptive significance of stomatal distri-
39 bution is less well understood. Stomata are most often found only on the lower
40 leaf surface (hypostomy) but occur on both surfaces (amphistomy) in many species
41 (Metcalfe and Chalk, 1950; Parkhurst, 1978; Mott et al., 1984). Theory and ex-
42 periments demonstrate that amphistomy increases photosynthetic rates under many

43 conditions. By creating a second parallel pathway for CO₂ diffusion within the meso-
44 phyll, amphistomy optimally supplies CO₂ (Parkhurst, 1978; Gutschick, 1984; Jones,
45 1985). Amphistomy is correlated with greater CO₂ diffusion (Beerling and Kelly,
46 1996) and higher photosynthetic rates (McKown et al., 2014). These observations
47 are corroborated by experiments demonstrating that amphistomy increases maxi-
48 mum photosynthetic rates by up to 20% (Parkhurst and Mott, 1990). On the other
49 hand, amphistomy can increase transpiration (Jones, 1985; Foster and Smith, 1986;
50 Buckley et al., 2015). While transition to amphistomy is thus thought to increase
51 transpiration, empirical studies suggest greater water-use efficiency in amphistoma-
52 tous species (Bucher et al., 2017). Hence, amphistomy appears to benefit a plant's
53 carbon use relative to water loss and should be favored when CO₂ limits photo-
54 synthetic rate. The open questions are under what ecological conditions does CO₂
55 supply most strongly limit photosynthetic rate (Peat and Fitter, 1994) and when is
56 photosynthetic rate most important to fitness?

57 The leading, nonmutually exclusive hypotheses are that 1) open habitats favour
58 amphistomy because CO₂ diffusion most strongly limits photosynthetic rate under
59 high light and 2) herbaceous growth form favours amphistomy because traits that
60 maximize photosynthetic rate are often under stronger selection in herbs. Salisbury
61 (1927) first noted that amphistomy is most common in herbaceous plants from open
62 habitats (i.e., with high light) of the British flora. These observations have been
63 replicated in other studies (Mott et al., 1984; Peat and Fitter, 1994; Jordan et al.,
64 2014; Muir, 2015) and may support physiological and ecological hypotheses that CO₂
65 most strongly limits photosynthesis in high light and/or photosynthesis contributes
66 most to fitness in herbaceous plants. Under high light, CO₂ can strongly limit max-
67 imum photosynthetic rates, especially in thick leaves (Jones, 1985). Hence, having

68 stomata on both surfaces relieves this limitation by adding a second parallel pathway
69 for CO₂ diffusion. Parkhurst (1978) argued that greater leaf thickness *per se* selected
70 for amphistomy, but there is little evidence for correlations between leaf thickness
71 and stomatal ratio independent of light (Mott et al., 1984; Gibson, 1996; Muir, 2015).
72 Amphistomy is correlated with open habitat in warm desert plants of western North
73 America (Mott et al., 1984; Gibson, 1996), among the Proteaceae (Jordan et al.,
74 2014), and in continental European herbs (Bucher et al., 2017).

75 Stomatal ratio is also associated with growth form. In the British flora, Salisbury
76 (1927) found that trees and shrubs are nearly always hypostomatous, whereas herbs
77 from open habitats are amphistomatous. This pattern holds when data are averaged
78 by family to coarsely control for phylogenetic nonindependence (Peat and Fitter,
79 1994) or when using alternative classification schemes, such as Raunkiaer life form
80 (Peat and Fitter, 1994). Across plants from ~ 90 families worldwide, growth form
81 is the strongest predictor of stomatal ratio when multiple factors are estimated si-
82 multaneously and controlling for phylogenetic nonindependence (Muir, 2015). These
83 patterns are consistent with other data indicating that many herbaceous plants are
84 under strong selection for high maximum photosynthetic rates (Bazzaz, 1979; Körner
85 et al., 1989; Wullschleger, 1993).

86 Although previous comparative studies have tested whether open habitat and growth
87 form influence stomatal ratio, we do not know if these effects are independent of one
88 another. Open habitat and growth form may not be independent because open
89 habitats generally consist of more short-statured, herbaceous plants. Some authors
90 have attempted to disentangle light and growth form by contrasting herbs from
91 open and understory habitats (Salisbury, 1927). However, this is problematic if
92 phylogenetic relationships are not controlled for, because shade species may share

93 traits simply because they are more closely related to each other than they are to
94 high light species. Finally, open habitat and growth form may also interact with one
95 another. For example, amphistomy may only be favored when CO₂ strongly limits
96 photosynthetic rate (e.g. in high light) *and* photosynthetic rate strongly limits fitness
97 (e.g. in herbs).

98 To better understand the adaptive significance of stomatal ratio, I asked three main
99 questions:

- 100 1. Are light habitat and growth form correlated?
- 101 2. Do light habitat and growth form influence stomatal ratio additively, or do
102 their effects interact?
- 103 3. Is evolution of stomatal ratio mediated by changes in stomatal density on the
104 adaxial (upper) surface, abaxial (lower) surface, or both?

105 The final question is important for addressing whether amphistomy is part of a co-
106 ordinated syndrome of traits that promote higher photosynthetic rate, as both the
107 light and growth form hypotheses assume. If evolved increases in stomatal ratio are
108 mediated by shifting abaxial stomata to the adaxial surface, holding total stomatal
109 density constant, then the overall increase in CO₂ diffusion would be small. In con-
110 trast, if amphistomy evolves by increasing adaxial stomatal density while holding
111 abaxial density constant, then *total* stomatal density must increase as well. Evolu-
112 tionary coordination of amphistomy and high stomatal density would reinforce one
113 another, increasing CO₂ supply to chloroplasts more than changes in either trait
114 would in isolation.

115 To address these questions, I reanalyzed existing data on stomatal ratio, light habi-
116 tat, and growth form in British angiosperms (Salisbury, 1927; Fitter and Peat, 1994,

117 2017) using phylogenetic comparative methods. The British angiosperm flora is well
118 suited for these questions because this flora has been comprehensively surveyed for
119 many ecologically important traits, meaning it is probably the least biased survey
120 of stomatal trait variation. Salisbury's observations on stomata and ecology in the
121 British flora have heavily influenced plant ecophysiology, but many of his and subse-
122 quent authors' analyses have significant limitations because of inadequate statistical
123 methods. For example, few analyses until recently account for phylogenetic nonin-
124 dependence (Felsenstein, 1985), which can strongly influence inferences on stomatal
125 traits and growth form (Kelly and Beerling, 1995, this study did not consider light).
126 A species-level phylogeny of the entire British flora (Lim et al., 2014) now allows for
127 the first time rigorous analysis of evolutionary relationships among stomatal ratio,
128 light, and growth form.

129 **Materials and Methods**

130 Data and annotated source code to generate this manuscript are available on GitHub
131 (<https://github.com/cdmuir/britstom>) and Dryad (Muir, 2017).

132 **Data on stomatal ratio, light habitat, growth form, and phy-** 133 **logenetic relationships**

134 I obtained data on ab- and adaxial stomatal density on 395 species from British
135 Ecological Flora (Salisbury, 1927; Fitter and Peat, 1994, 2017). Following recent
136 comparative analyses (e.g. Bartelheimer and Poschlod, 2016; Salguero-Gómez et al.,
137 2016; Shipley et al.), I used Ellenberg light indicator values (Ellenberg, 1974) and

138 Raunkiaer life form (Raunkiaer, 1934) as measures of light habitat and growth form,
139 respectively. Hence, I am assuming that the species' light habitat is closely related to
140 the type of habitat (open versus closed) where that species is found. Both attributes
141 have been recently updated by taxonomic experts of the British flora (PLANTATT,
142 Hill et al. (2004)). Ellenberg light indicator values are hereafter abbreviated L-
143 value. I used a dated molecular phylogeny of the British flora (Lim et al., 2014)
144 available from TreeBASE (<http://treebase.org/>; accession number 15105). 14 species
145 (3.5%) in the dataset were not present in the phylogeny. For 8 of these species, I
146 used the position of a congeneric species as a proxy for the focal species (following
147 Pennell et al., 2016). When multiple congeneric species were present, I consulted
148 the phylogenetic literature to identify the most closely related proxy species (Scheen
149 et al., 2004; Salmaki et al., 2013). For the remaining 6 missing species, I positioned
150 them in the tree based on phylogenetic relationships to other genera or families
151 present in the tree (Fior et al., 2006). Because many phylogenetic comparative
152 methods do not allow polytomies, zero-length branches, and non-ultrametric trees, I
153 made several small adjustments to the tree. I resolved polytomies randomly using the
154 'multi2di' function in **phytools** version 0.6-00 (Revell, 2012). I added 0.02 my to all
155 zero-length branches, as this was approximately the length of the shortest nonzero
156 branch length in the tree. After these changes, I slightly altered terminal branch
157 lengths to make the tree precisely ultrametric.

158 I excluded data on hyrdrophytes (14 species) because many of these species are
159 hyperstomatous (Fig. S1) due to the fact that leaves may rest on the water's surface,
160 selecting for stomata to be present on the upper surface only. I also excluded C₄
161 (3 species) and CAM (2 species) plants. I limited this investigation to angiosperms
162 because only 4 non-angiosperms had stomata data. The final dataset contained 372

163 species (Fig. 1). The R code accompanying this paper documents these decisions
164 with citations to the relevant literature.

165 Following Muir (2015), I calculated stomatal ratio in two different ways depending
166 on what was most appropriate for the question:

$$SR_{propAd} = \frac{SD_{ad}}{SD_{total}} \quad (1)$$

$$SR_{even} = \frac{\min\{SD_{ab}, SD_{ad}\}}{\max\{SD_{ab}, SD_{ad}\}} \quad (2)$$

167 SD_{ab} and SD_{ad} are the stomatal densities on abaxial or adaxial surface, respectively.
168 $SD_{total} = SD_{ab} + SD_{ad}$. SR_{propAd} is the proportion of stomata density on the adaxial
169 surface, which is useful for discriminating among hypostomatous ($SR_{propAd} = 0$),
170 amphistomatous ($0 < SR_{propAd} < 1$), and hyperstomatous species ($SR_{propAd} = 1$).
171 SR_{even} indicates how evenly stomatal densities are distributed across both leaf sur-
172 faces. This expression is useful because several hypotheses are based on the fact that
173 a more even distribution should optimize leaf CO_2 diffusion.

174 **Testing for an association between open habitat and growth** 175 **form**

176 I tested whether Raunkiær life form was associated with L-value among British an-
177 giosperms using ANOVA with Type-2 sum of squares. I did not use phylogenetic
178 ANOVA for this test because there was no phylogenetic signal in the regression fit
179 using **phylolm** version 2.5 (Ho and Ané, 2014). See the R code accompanying this

180 paper for further detail. I predicted that species with faster life histories, especially
181 therophytes (annuals), would have greater L-values than species with slower life his-
182 tories, especially phanerophytes, which are mostly long-lived trees.

183 **Open habitat, growth form, and stomatal ratio**

184 I compared phylogenetic linear models to test whether Raunkiaer life form, L-value,
185 or interactions between them predicted SR_{even} . Unlike the analysis above, there
186 was significant phylogenetic signal in this comparison (see R code). I used SR_{even}
187 rather than SR_{propAd} as the response variable because the hypothesis is that faster
188 life history and/or high light favor more even stomatal densities on each surface.
189 I fit models using **phylolm** and extracted Akaike Information Criteria (AIC). For
190 these and subsequent analyses, I assumed an Ornstein-Uhlenbeck process model for
191 the residuals with the root character state integrated over the stationary distribu-
192 tion. I used a 10^4 parametric bootstrap samples of the full model (including main
193 effects and interactions) to calculate parameter confidence intervals (Boettiger et al.,
194 2012).

195 **Does ab- or adaxial stomatal density contribute more to stom-** 196 **atal ratio evolution?**

197 I used two related phylogenetic methods, variance decomposition and structural equa-
198 tion modeling (SEM), to assess the relative contribution of ab- versus adaxial stom-
199 atal density to light-mediated stomatal ratio evolution. First, the contribution of ab-
200 versus adaxial stomatal density can be calculated using phylogenetic variance de-

201 composition methods as derived below. Because stomatal density is highly skewed,
 202 I log-transformed values for normality:

$$\text{SR}_{\text{even}} = \frac{\text{SD}_{\text{ad}}}{\text{SD}_{\text{ab}}} \quad (3)$$

$$\log(\text{SR}_{\text{even}}) = \log(\text{SD}_{\text{ad}}) - \log(\text{SD}_{\text{ab}}) \quad (4)$$

$$\text{sr}_{\text{even}} = \text{sd}_{\text{ad}} - \text{sd}_{\text{ab}} \quad (5)$$

203 Lowercase variables (sr, sd) indicate log-transformed values. Because some species
 204 had zero adaxial stomata, I added one to all values prior to log-transformation. To
 205 make the variance decomposition calculations tractable, I have defined SR_{even} here
 206 as the ratio of ad- to abaxial stomatal density because in most cases adaxial stomatal
 207 density is lower than abaxial (see Eq. 2). This differs from analyses described above
 208 because in those I wanted to test what factors influenced the evenness of stomatal
 209 densities, regardless of which surface had higher density. With this modified form,
 210 the variance in sr_{even} can readily be decomposed into contributions of sd_{ad} , sd_{ab} , and
 211 their covariance:

$$\text{Var}(\text{sr}_{\text{even}}) = \text{Var}(\text{sd}_{\text{ad}}) + \text{Var}(\text{sd}_{\text{ab}}) - 2\text{Cov}(\text{sd}_{\text{ad}}, \text{sd}_{\text{ab}}) \quad (6)$$

212 I did not use the raw covariance, but rather estimated the phylogenetic covariance
 213 matrix between L-value, sd_{ab} , and sd_{ad} using a multivariate Ornstein-Uhlenbeck
 214 model fit in **Rphylopars** version 0.2.9 (Goolsby et al., 2016, 2017). From the co-

215 variance matrix, I estimated the contribution of abaxial density, adaxial density, and
 216 their covariance as:

$$\text{Contribution of } sd_{ad} = \frac{\text{Var}(sd_{ad})}{\text{Var}(sr_{even})} \quad (7)$$

$$\text{Contribution of } sd_{ab} = \frac{\text{Var}(sd_{ab})}{\text{Var}(sr_{even})} \quad (8)$$

$$\text{Contribution of } \text{Cov}(sd_{ad}, sd_{ab}) = \frac{\text{Cov}(sd_{ad}, sd_{ab})}{\text{Var}(sr_{even})} \quad (9)$$

217 respectively. Note that when ab- and adaxial densities positively covary, the contri-
 218 bution will be negative because this reduces the variance in stomatal ratio.

219 I also wanted to test whether light-mediated evolution of stomatal ratio acted mostly
 220 by 1) increasing adaxial stomatal density while maintaining abaxial density, or 2)
 221 keeping total stomatal density the same, but shifting a greater proportion to the adax-
 222 ial surface. The first scenario predicts that the phylogenetic regression of L-value on
 223 sd_{ad} is stronger than that for sd_{ab} . The second scenario predicts that L-value acts sim-
 224 ilarly on both and that there is a negative covariance ($\text{Cov}(sd_{ad}, sd_{ab}) < 0$). I tested
 225 these competing predictions by fitting a very simple phylogenetic SEM (see Mason
 226 et al., 2016, for a similar approach). The model uses the phylogenetic covariance
 227 matrix, as described above, to simultaneously estimate regressions of L-value on sd_{ad}
 228 and sd_{ab} while allowing covariance between them (i.e. estimating $\text{Cov}(sd_{ad}, sd_{ab})$).
 229 To fit the SEM, I used the R package **lavaan** version 0.5-23.1097 (Rosseel, 2012).
 230 I tested whether parameter estimates were significantly different from zero using
 231 z-scores.

Results

Light tolerance varies among Raunkiær life forms

Ellenberg light indicator values (L-value) differed significantly among life forms (Fig. 2; ANOVA - $F_{4,367} = 18.3$, $P = 1.1 \times 10^{-13}$). Therophytes (annuals), hemicryptophytes (perennial herbs with buds near the soil surface), and chamaephytes (subshrubs) had greater L-values than phanerophytes (large woody plants) and geophytes (perennial herbs with storage organs) (Fig. 2).

Interactions between light and Raunkiær life form determine stomatal ratio

Overall, SR_{even} increased with L-value, but there was a significant interaction ($\Delta AIC > 2$, Table 1) between Raunkiær life form and L-value (Fig. 3). Both life form and L-value significantly increased model fit, though L-value had a markedly larger effect on model AIC (Table 1). The significant interaction is caused by different slopes between life forms. Among life forms with the overall greatest L-value (therophytes, hemicryptophytes, and chamaephytes, see Fig. 2), there was a strong positive relationship between L-value and SR_{even} . Parametrically bootstrapped 95% confidence intervals for the slope did not overlap zero (Fig. 3). The slope was weakly positive or not significantly different from zero in the most shade-adapted life forms (geophytes and phanerophytes), albeit the patterns were distinct in these groups. There were both hypostomatous ($SR_{\text{even}} \approx 0$) and amphistomatous ($SR_{\text{even}} \approx 1$) geophytes, but these were distributed across L-values. In contrast, phanerophytes were nearly

253 always hypostomatous regardless of L-value.

254 **Adaxial stomatal density contributes most of the variation in** 255 **stomatal ratio**

256 Adaxial ('upper') stomatal density contributed most to the evolutionary variation
257 in stomatal ratio. The contributions of adaxial density, abaxial density, and their
258 covariance are 1.12, 0.38, and -0.5, respectively. This implies that evolutionary varia-
259 tion in adaxial stomatal density is greater than that for stomatal ratio due to positive
260 covariance between ab- and adaxial stomatal density.

261 Similarly, the phylogenetic SEM showed that changes in stomatal ratio associated
262 with L-value can be attributed mostly to evolution of adaxial stomatal density
263 (Fig. 4). Both sd_{ad} and sd_{ab} increased with L-value ($P = 1.2 \times 10^{-8}$ and 8.9×10^{-7} ,
264 respectively). However, the regression of L-value on sd_{ad} was $2\times$ that of L-value on
265 sd_{ab} (0.24 versus 0.12). Because stomatal densities were natural log-transformed, this
266 implies an increase in L-value by one leads to a 1.27-fold change in adaxial stom-
267 atal density versus a 1.13-fold change in abaxial stomatal density. The SEM also
268 showed a significant positive covariance between stomatal densities on each surface
269 ($P = 2.5 \times 10^{-10}$). These results together imply that total stomatal density increases
270 with L-value, but the response is mediated mostly by increases in adaxial stomatal
271 density.

Discussion

The ratio of stomatal densities on the abaxial (‘lower’) to that of the adaxial (‘upper’) surface varies greatly across plant species, but the adaptive significance is not clear. Comparative studies correlating stomatal ratio to ecological factors can distinguish among competing hypotheses and reveal critical experiments for future work. Previous comparative studies suggested that high light and herbaceous growth form favor amphistomy (Mott et al., 1984; Jordan et al., 2014; Muir, 2015; Bucher et al., 2017), particularly in the British flora (Salisbury, 1927; Peat and Fitter, 1994). However, none of these studies have accounted for the fact that light and growth form are often confounded – open, high light habitats are often dominated by herbs – or the fact that species are not independent because of shared evolutionary history. Here, I reanalyzed data on stomata, light tolerance, and growth form in British angiosperms using phylogenetic comparative methods. As expected, species’ light tolerance (Ellenberg light indicator or L-value) is confounded with growth form (Raunkiær life form; Fig. 2). Nevertheless, both L-value and Raunkiær life form affect stomatal ratio, but these factors also interact; the influence of L-value on stomatal ratio varies across forms. These novel findings provide further evidence that variation in stomatal ratio is adaptive and have important implications for interpreting changes in stomatal ratio through the paleo record (Jordan et al., 2014) and during domestication (Milla et al., 2013).

Adaptive significance of amphistomy

Previously, associations between light, growth form, and stomatal ratio have been interpreted in isolation as indicating that either high light and/or herbaceous growth

295 form favors amphistomy. In British angiosperms, both factors are important, though
296 statistical analyses suggest that light may be a stronger determinant than growth
297 form (Table 1). Unlike previous studies, I found a significant interaction between
298 light and growth form among British angiosperms, which suggests that amphistomy
299 may only be strongly favored when CO₂ strongly limits photosynthesis (as in open
300 habitat) *and* photosynthesis strongly limits fitness (as in herbs). This is consistent
301 with life history theory predicting that the demography of open habitat herbs is
302 strongly limited by plant growth (Franco and Silvertown, 1996). The ideal way to
303 test this would be to measure selection on stomatal ratio in a species that varied
304 quantitatively in both stomatal ratio and life history (e.g., containing both annual
305 and perennial forms). I predict that amphistomy will be favored more strongly in
306 the annual form grown under high light compared to an annual under low light
307 or a perennial in high light, and much more strongly than a perennial grown in low
308 light. Similar experiments could also be performed to test if and when light-mediated
309 plasticity in stomatal ratio is adaptive (Gay and Hurd, 1975; Mott and Michaelson,
310 1991; Fontana et al., 2017).

311 The prevalence of amphistomatous species in high light habitats supports the hy-
312 pothesis that amphistomy is an adaptation to maximize photosynthetic rates by
313 increasing CO₂ diffusion (Jones, 1985). It is also evidence against the hypothesis
314 that the principle fitness cost of amphistomy is water loss (Darwin, 1886; Foster and
315 Smith, 1986) or dehydration of palisade mesophyll (Buckley et al., 2015), though
316 these factors are likely very important in determining differential regulation of stom-
317 ata on each surface. Since evaporative demand increases under high light, under
318 these hypotheses we would expect plants in high light to be hypostomatous. Because
319 stomatal conductances on each surface can be regulated independently in response

320 to the environment (Darwin, 1898; Pospíšilová and Solárová, 1984; Smith, 1981; Re-
 321 ich, 1984; Mott and O’Leary, 1984), amphistomatous leaves likely cope with these
 322 stresses by rapidly closing adaxial stomata when water supply cannot match evapo-
 323 rative demands (Richardson et al., 2017). Instead, patterns in the British flora are
 324 at least consistent with the idea that adaxial stomata increase susceptibility to foliar
 325 pathogens (Gutschick, 1984; McKown et al., 2014). The cost of adaxial stomata
 326 may be greater in the shade because greater leaf wetness and lower ultraviolet light
 327 provide a more suitable microclimate for many foliar pathogens.

328 **Amphistomy as a proxy for open habitat**

329 These observations from the British flora partially support the hypothesis that am-
 330 phistomy can be used a proxy for open habitat in paleoenvironment reconstruction
 331 (Carpenter, 1994; Jordan et al., 2014; Carpenter et al., 2015) but also point out pre-
 332 viously unknown subtleties. These previous studies based their conclusions on data
 333 from Proteaceae, in which there is little quantitative variation in stomatal ratio;
 334 species are either completely hypostomatous ($SR_{propAd} \approx 0$) or completely amphis-
 335 tomatous ($SR_{propAd} \approx 0.5$). Stomatal ratio in British angiosperms is also bimodal
 336 (Peat and Fitter, 1994), but across many families there is also quantitative variation.
 337 Importantly, this means that quantitative variation in stomatal ratio may provide a
 338 more precise, quantitative indicator of vegetation type, rather than simply ‘open’ or
 339 ‘closed’. A quantitative relationship between L-value and stomatal ratio has already
 340 been shown for herbaceous perennials (Bucher et al., 2017), but we now know that
 341 it holds among annuals (therophytes), subshrubs (chamaephytes), and, to a lesser
 342 extent, geophytes as well (Fig. 3).

343 The weak or nonsignificant relationship between L-value and stomatal ratio in geo-
 344 phytes and phanerophytes suggests that in some cases amphistomy may not reliably
 345 indicate open habitat without further information. For example, perhaps amphis-
 346 tomatus geophytes from partially shaded habitats are spring ephemerals, so they
 347 experience high light during their growth phase, but this has not been tested. Like-
 348 wise, phanerophytes (most tall trees) are almost always hypostomatous (see also
 349 Muir (2015)). Most British phanerophytes are tall, hypostomatous trees, but the
 350 exceptions are telling. For example, the most amphistomatous phanerophyte in this
 351 dataset is *Brassica oleracea*, a short-statured biennial that has more in common
 352 physiologically with hemicryptophytes than other phanerophytes. The other am-
 353 phistomatous phanerophytes in this data set (*Populus nigra* and *Lavatera arborea*)
 354 are fast-growing pioneer species.

355 Finally, phylogenetic information should improve inferences about paleoclimates be-
 356 cause there is appreciable phylogenetic signal in stomatal ratio. The phylogenetic
 357 half-life of stomatal ratio evolution, after accounting for L-value and Raunkiaer life
 358 form, is $\log(2)/\alpha = 1.5$ my (see Table 1 for maximum likelihood estimates of α , the
 359 return rate in the Ornstein-Uhlenbeck model of trait evolution). This lag time may
 360 indicate that evolving to the ‘optimum’ is constrained by the shape of the fitness
 361 landscape (Muir, 2015) or that other unmeasured factors which affect stomatal ra-
 362 tio have some phylogenetic signal. Regardless of the mechanism, this fact means
 363 that researchers may be able to use data from closely related species to improve
 364 paleoenvironment reconstruction.

365 **Why does adaxial stomatal density control stomatal ratio?**

366 Variation in stomatal ratio is determined primarily by evolution of adaxial stom-
367 atal density and is coordinated with increases in total leaf stomatal density summed
368 across both surfaces. Note here that I am referring only to evolutionary variation in
369 stomatal ratio among species; different processes may mediate within species vari-
370 ation or plastic responses. Phylogenetic analyses show that changes in stomatal
371 ratio and total stomatal density, especially in response to L-value, are predominantly
372 mediated by changes in adaxial stomatal density. This highly nonrandom pattern
373 among British angiosperms mirrors evolutionary changes wrought by domestication
374 (Milla et al., 2013); crop species tend to have higher adaxial stomatal density than
375 their wild relatives.

376 There are at least two hypotheses that could explain why adaxial stomatal density
377 is the most responsive. The first I refer to as the ‘real estate’ hypothesis. In hy-
378 postomatous plants, the lower surface is already crowded with stomata, and hence
379 plants must increase the real estate available for stomata by developing them on the
380 upper surface whenever there is selection for greater stomatal density. When stomata
381 are packed too densely on one surface, stomatal interference limits their function-
382 ing and hence may create a strong selective pressure for amphistomy (Parlange and
383 Waggoner, 1970; Dow et al., 2014).

384 I refer to the second hypothesis as the ‘coordination’ hypothesis. In this scenario,
385 ecological conditions such as high light select for both increased total stomatal density
386 and for amphistomy because these traits work well in coordination with one another.
387 For example, if stomatal density were very high on a hypostomatous plant, then CO₂
388 would be more strongly limited by the mesophyll. Adding a second parallel pathway

389 for diffusion by developing stomata on both surfaces would restore a more optimal
390 balance between stomatal and mesophyll limitations. Conversely, there would be
391 little benefit to amphistomy when total stomatal density is low because CO₂ diffusion
392 is strongly limited by stomatal resistance, and therefore photosynthetic rate is not
393 sensitive to changes in mesophyll diffusion mediated by stomatal ratio. A related
394 prediction is that increased atmospheric CO₂ may select for reduced stomatal ratio
395 and density primarily by decreasing adaxial stomatal density, but this has not been
396 well tested (but see Woodward and Bazzaz, 1988).

397 **Conclusions**

398 By revisiting this classic ecological dataset with modern phylogenetic comparative
399 methods, I have shown that amphistomy is strongly associated with both light and
400 growth form, but the interaction between these factors is also important. Fur-
401 thermore, amphistomy and high stomatal density are closely connected in species
402 from high light environments, suggesting selection for coordination between these
403 traits.

404 **Acknowledgements**

405 I thank Sally Otto, Matt Pennell, and Rob Salguero-Gómez for feedback on this
406 manuscript. I was supported by an NSERC CREATE grant.

407 Author contribution statement

408 CDM designed the study, collected data, analyzed the data, and wrote the manuscript.

References

- Bartelheimer, M. and P. Poschlod, 2016. Functional characterizations of Ellenberg indicator values—a review on ecophysiological determinants. *Functional Ecology* 30:506–516.
- Bazzaz, F., 1979. The physiological ecology of plant succession. *Annual Review of Ecology and Systematics* 10:351–71.
- Beerling, D. J. and C. K. Kelly, 1996. Evolutionary comparative analyses of the relationship between leaf structure and function. *New Phytologist* 134:35–51.
- Beerling, D. J. and D. L. Royer, 2011. Convergent Cenozoic CO₂ history. *Nature Geoscience* 4:418–420.
- Boettiger, C., G. Coop, and P. Ralph, 2012. Is your phylogeny informative? Measuring the power of comparative methods. *Evolution* 66:2240–2251.
- Bucher, S. F., K. Auerswald, C. Grün-Wenzel, S. I. Higgins, J. G. Jorge, and C. Römermann, 2017. Stomatal traits relate to habitat preferences of herbaceous species in a temperate climate. *Flora* 229:107–115.
- Buckley, T. N., G. P. John, C. Scoffoni, and L. Sack, 2015. How does leaf anatomy influence water transport outside the xylem? *Plant Physiology* 168:1616–1635.
- Carpenter, R. J., 1994. Cuticular morphology and aspects of the ecology and fossil history of North Queensland rainforest Proteaceae. *Botanical Journal of the Linnean Society* 116:249.
- Carpenter, R. J., M. K. Macphail, G. J. Jordan, and R. S. Hill, 2015. Fossil evidence

430 for open, Proteaceae-dominated heathlands and fire in the Late Cretaceous of
 431 Australia. *American Journal of Botany* 102:2092–2107.

432 Darwin, F., 1886. On the relation between the “bloom” on leaves and the distribution
 433 of the stomata. *Botanical Journal of the Linnean Society* 22:99–116.

434 ———, 1898. Observations on stomata. *Philosophical Transactions of the Royal*
 435 *Society B: Biological Sciences* 190:531–621.

436 Dow, G. J., J. A. Berry, and D. C. Bergmann, 2014. The physiological importance
 437 of developmental mechanisms that enforce proper stomatal spacing in *Arabidopsis*
 438 *thaliana*. *New Phytologist* 201:1205–1217.

439 Ellenberg, H., 1974. Indicator values of vascular plants in central Europe, *Scripta*
 440 *Geobotanica*, vol. 9. Springer-Verlag, Göttingen, Germany.

441 Felsenstein, J., 1985. Phylogenies and the comparative method. *The American*
 442 *Naturalist* 1:1–15.

443 Fior, S., P. O. Karis, G. Casazza, L. Minuto, and F. Sala, 2006. Molecular phylogeny
 444 of the Caryophyllaceae (Caryophyllales) inferred from chloroplast matk and nuclear
 445 rDNA ITS sequences. *American Journal of Botany* 93:399–411.

446 Fitter, A. and H. Peat, 1994. The ecological flora database. *Journal of Ecology*
 447 82:415–425.

448 ———, 2017. Ecological flora of the British isles. URL
 449 <http://www.ecoflora.co.uk>.

450 Fontana, M., M. Labrecque, A. Collin, and N. Bélanger, 2017. Stomatal distribu-
 451 tion patterns change according to leaf development and leaf water status in *Salix*
 452 *miyabeana*. *Plant Growth Regulation* 81:63–70.

- 453 Foster, J. and W. Smith, 1986. Influence of stomatal distribution on transpiration
454 in low-wind environments. *Plant, Cell & Environment* 9:751–759.
- 455 Franco, M. and J. Silvertown, 1996. Life history variation in plants: an exploration
456 of the fast-slow continuum hypothesis. *Philosophical Transactions: Biological Sci-*
457 *ences* 351:1341–1348.
- 458 Gay, A. and R. Hurd, 1975. The influence of light on stomatal density in the tomato.
459 *New Phytologist* 75:37–46.
- 460 Gibson, A. C., 1996. *Structure-Function Relations of Warm Desert Plants*. Springer-
461 Verlag, Berlin.
- 462 Givnish, T. J., 1987. Comparative studies of leaf form: assessing the relative roles
463 of selective pressures and phylogenetic constraints. *New Phytologist* 106:131–160.
- 464 Goolsby, E. W., J. Bruggeman, and C. Ané, 2016. Rphylopars: Phyloge-
465 netic Comparative Tools for Missing Data and Within-Species Variation. URL
466 <https://CRAN.R-project.org/package=Rphylopars>. R package version 0.2.9.
- 467 ———, 2017. Rphylopars: fast multivariate phylogenetic comparative methods for
468 missing data and within-species variation. *Methods in Ecology and Evolution*
469 8:22–27.
- 470 Gutschick, V. P., 1984. Photosynthesis model for C₃ leaves incorporating CO₂ trans-
471 port, propagation of radiation, and biochemistry 2. ecological and agricultural
472 utility. *Photosynthetica* 18:569–595.
- 473 Haberlandt, G., 1914. *Physiological Plant Anatomy*. Macmillan and Co., London.
- 474 Hill, M., C. Preston, and D. Roy, 2004. *PLANTATT - Attributes of British and Irish*

- 475 Plants: Status, Size, Life History, Geography and Habitats. Centre for Ecology &
476 Hydrology, Huntingdon, Cambridgeshire.
- 477 Ho, L. S. T. and C. Ané, 2014. Intrinsic inference difficulties for trait evolution with
478 Ornstein-Uhlenbeck models. *Methods in Ecology and Evolution* 5:1133–1146.
- 479 Jones, H. G., 1985. Adaptive significance of leaf development and structural responses
480 to environment. Pp. 155–173, *in* N. R. Baker, W. Davies, and C. K. Ong, eds.
481 Control of Leaf Growth, *Society for Experimental Biology Seminar Series*, vol. 27.
482 Cambridge University Press, Cambridge.
- 483 Jordan, G. J., R. J. Carpenter, and T. J. Brodribb, 2014. Using fossil leaves as
484 evidence for open vegetation. *Palaeogeography, Palaeoclimatology, Palaeoecology*
485 395:168–175.
- 486 Kelly, C. and D. Beerling, 1995. Plant life form, stomatal density and taxonomic
487 relatedness: a reanalysis of Salisbury (1927). *Functional Ecology* 9:422–431.
- 488 Körner, C., M. Neumayer, S. P. Menendez-Riedl, and A. Smeets-Scheel, 1989. Func-
489 tional morphology of mountain plants. *Flora* 182:353–383.
- 490 Lim, J., M. J. Crawley, N. De Vere, T. Rich, and V. Savolainen, 2014. A phylogenetic
491 analysis of the British flora sheds light on the evolutionary and ecological factors
492 driving plant invasions. *Ecology and Evolution* 4:4258–4269.
- 493 Mason, C. M., E. W. Goolsby, D. P. Humphreys, and L. A. Donovan, 2016. Phy-
494 logenetic structural equation modelling reveals no need for an ‘origin?’ of the leaf
495 economics spectrum. *Ecology letters* 19:54–61.
- 496 McElwain, J. C. and M. Steinthorsdottir, 2017. Paleocology, ploidy, paleoatmo-

497 spheric composition, and developmental biology: a review of the multiple uses of
498 fossil stomata. *Plant Physiology* 174:650–664.

499 McKown, A. D., R. D. Guy, L. Quamme, J. Klápště, J. La Mantia, C. Constabel,
500 Y. A. El-Kassaby, R. C. Hamelin, M. Zifkin, and M. Azam, 2014. Association
501 genetics, geography and ecophysiology link stomatal patterning in *Populus tri-*
502 *chocarpa* with carbon gain and disease resistance trade-offs. *Molecular Ecology*
503 23:5771–5790.

504 Melotto, M., L. Zhang, P. R. Oblessuc, and S. Y. He, 2017. Stomatal defense a
505 decade later. *Plant Physiology* 174:561–571.

506 Metcalfe, C. R. and L. Chalk, 1950. *Anatomy of the dicotyledons*, Vols. 1 & 2. First
507 ed. Oxford University Press, Oxford.

508 Milla, R., N. de Diego-Vico, and N. Martín-Robles, 2013. Shifts in stomatal traits
509 following the domestication of plant species. *Journal of Experimental Botany*
510 64:3137–3146.

511 Mott, K. A., A. C. Gibson, and J. W. O’Leary, 1984. The adaptive significance of
512 amphistomatic leaves. *Plant, Cell & Environment* 5:455–460.

513 Mott, K. A. and O. Michaelson, 1991. Amphistomy as an adaptation to high light
514 intensity in *Ambrosia cordifolia* (Compositae). *American Journal of Botany* 78:76–
515 79.

516 Mott, K. A. and J. W. O’Leary, 1984. Stomatal behavior and CO₂ exchange char-
517 acteristics in amphistomatous leaves. *Plant Physiology* 74:47–51.

518 Muir, C. D., 2015. Making pore choices: repeated regime shifts in stomatal ratio.
519 *Proc. R. Soc. B* 282:20151498.

- 520 ———, 2017. Data from: Light and life form interact to shape stomatal ratio among
521 British angiosperms. URL <http://dx.doi.org/10.5061/dryad.?????>
- 522 Parkhurst, D. F., 1978. The adaptive significance of stomatal occurrence on one or
523 both surfaces of leaves. *The Journal of Ecology* 66:367–383.
- 524 Parkhurst, D. F. and K. A. Mott, 1990. Intercellular diffusion limits to CO₂ uptake
525 in leaves studied in air and helox. *Plant Physiology* 94:1024–1032.
- 526 Parlange, J.-Y. and P. E. Waggoner, 1970. Stomatal dimensions and resistance to
527 diffusion. *Plant Physiology* 46:337–342.
- 528 Peat, H. and A. Fitter, 1994. A comparative study of the distribution and density of
529 stomata in the British flora. *Biological Journal of the Linnean Society* 52:377–393.
- 530 Pennell, M. W., R. G. FitzJohn, and W. K. Cornwell, 2016. A simple approach for
531 maximizing the overlap of phylogenetic and comparative data. *Methods in Ecology*
532 *and Evolution* 7:751–758.
- 533 Pospíšilová, J. and J. Solárová, 1984. Environmental and biological control of diffu-
534 sive conductances of adaxial and abaxial leaf epidermes. *Photosynthetica* 18:445–
535 453.
- 536 Raunkiaer, C. C., 1934. *The Life Forms of Plants and Statistical Plant Geography*.
537 Clarendon Press, Oxford.
- 538 Reich, P., 1984. Relationships between leaf age, irradiance, leaf conductance, CO₂
539 exchange, and water-use efficiency in hybrid poplar. *Photosynthetica* 18:445–453.
- 540 Revell, L. J., 2012. phytools: An R package for phylogenetic comparative biology
541 (and other things). *Methods in Ecology and Evolution* 3:217–223.

- 542 Richardson, F., T. J. Brodribb, and G. J. Jordan, 2017. Amphistomatic leaf sur-
543 faces independently regulate gas exchange in response to variations in evaporative
544 demand. *Tree Physiology* Pp. 1–10.
- 545 Rosseel, Y., 2012. lavaan: An R package for structural equation modeling. *Journal*
546 *of Statistical Software* 48:1–36.
- 547 Royer, D. L., 2001. Stomatal density and stomatal index as indicators of paleoatmo-
548 spheric CO₂ concentration. *Review of Palaeobotany and Palynology* 114:1–28.
- 549 Sack, L. and T. N. Buckley, 2016. The developmental basis of stomatal density and
550 flux. *Plant Physiology* 171:2358–2363.
- 551 Salguero-Gómez, R., O. R. Jones, E. Jongejans, S. P. Blomberg, D. J. Hodgson,
552 C. Mbeau-Ache, P. A. Zuidema, H. de Kroon, and Y. M. Buckley, 2016. Fast-
553 slow continuum and reproductive strategies structure plant life-history variation
554 worldwide. *Proceedings of the National Academy of Sciences of the United States*
555 *of America* 113:230–235.
- 556 Salisbury, E., 1927. On the causes and ecological significance of stomatal frequency,
557 with special reference to the woodland flora. *Philosophical Transactions of the*
558 *Royal Society of London. Series B* 216:1–65.
- 559 Salmaki, Y., S. Zarre, O. Ryding, C. Lindqvist, C. Bräuchler, G. Heubl, J. Barber,
560 and M. Bendiksby, 2013. Molecular phylogeny of tribe Stachydeae (Lamiaceae
561 subfamily Lamioideae). *Molecular Phylogenetics and Evolution* 69:535–551.
- 562 Scheen, A.-C., C. Brochmann, A. K. Brysting, R. Elven, A. Morris, D. E. Soltis, P. S.
563 Soltis, and V. A. Albert, 2004. Northern hemisphere biogeography of *Cerastium*

564 (Caryophyllaceae): insights from phylogenetic analysis of noncoding plastid nu-
565 cleotide sequences. *American Journal of Botany* 91:943–952.

566 Shipley, B., M. Belluau, I. Kühn, N. A. Soudzilovskaia, M. Bahn, J. Penue-
567 las, J. Kattge, L. Sack, J. Cavender-Bares, W. A. Ozinga, B. Blonder, P. M.
568 van Bodegom, P. Manning, T. Hickler, E. Sosinski, V. D. P. Pillar, and
569 V. Onipchenko, ????. Predicting habitat affinities of plant species using com-
570 monly measured functional traits. *Journal of Vegetation Science* Pp. n/a–n/a.
571 URL <http://dx.doi.org/10.1111/jvs.12554>.

572 Smith, W., 1981. Temperature and water relation patterns in subalpine understory
573 plants. *Oecologia* 48:353–359.

574 Smith, W. K., T. C. Vogelmann, E. H. DeLucia, D. T. Bell, and K. A. Shepherd,
575 1997. Leaf form and photosynthesis. *BioScience* 11:785–793.

576 Wolfe, J. A., 1971. Tertiary climatic fluctuations and methods of analysis of Tertiary
577 floras. *Palaeogeography, Palaeoclimatology, Palaeoecology* 9:27–57.

578 Woodward, F., 1987. Stomatal numbers are sensitive to increases in CO₂ from pre-
579 industrial levels. *Nature* 327:617–618.

580 Woodward, F. I. and F. Bazzaz, 1988. The responses of stomatal density to CO₂
581 partial pressure. *Journal of Experimental Botany* 39:1771–1781.

582 Wullschleger, S. D., 1993. Biochemical limitations to carbon assimilation in C₃
583 plants? A retrospective analysis of the A/Ci curves from 109 species. *Journal*
584 *of Experimental Botany* 44:907–920.

Table 1: Interaction between species' Ellenberg light indicator value (L-value) and Raunkiaer lifeform predict stomatal ratio (SR_{even}). I compared phylogenetic linear models using the Akaike Information Criterion (AIC), where $\text{AIC} = 2k - 2\log(\mathcal{L})$. k is the number of model parameters and \mathcal{L} is the model likelihood. Given a set of candidate models, the difference in AIC between a model and the lowest AIC (ΔAIC) indicates the relative fit of competing models. The correlation coefficient r^2 is another indicator of model fit. α and σ^2 are the return rate and diffusion parameters of the Ornstein-Uhlenbeck model of trait evolution.

Model: $\text{SR}_{\text{even}} \sim$	α	σ^2	r^2	k	$\log(\mathcal{L})$	AIC	ΔAIC
L-value \times lifeform	0.46	0.068	0.34	12	-33.2	90.4	0
L-value + lifeform	0.46	0.071	0.32	8	-40.2	96.4	6
L-value	0.64	0.107	0.26	4	-59.3	126.6	36.2
lifeform	0.34	0.067	0.15	7	-79.2	172.4	82
null	0.29	0.067	0	3	-107.6	221.1	130.7




Figure 1: CAPTION




Figure 2: Lifeforms have different tolerances for sun and shade among British angiosperms. Each panel is the distribution of Ellenberg light indicator values on an integer scale of 1-9 for different Raunkiær life forms. Height of the bars indicate the raw proportion of species in each bin for that lifeform. The sample size for each lifeform is listed next in parentheses. The mean (open circle) and 95% confidence intervals (black line) around the mean Ellenberg light indicator value for each lifeform based on phylogenetic regression are above the histogram.




Figure 3: The effect of light on stomatal ratio depends on Raunkiaer life form. Greater Ellenberg light indicator values (L-value) are associated with greater stomatal ratio (SR_{even}) in therophytes, hemicryptophytes, and chamaephytes but not geophytes and phanerophytes. The maximum likelihood slope from phylogenetic regression is given with statistical significance based on 10^4 parametric bootstrap samples. Numbers in parentheses next to Raunkiaer life form are the sample sizes in the final dataset. Estimated slopes (solid line) and 95% bootstrapped confidence intervals (gray polygon between dashed lines) are plotted against raw data. Points have been jittered for visual clarity.




Figure 4: Light-mediated evolution of stomatal ratio is mostly driven by increased adaxial (‘upper’) stomatal density (Panel A), whereas abaxial (‘lower’) stomatal density (Panel B) is similar across Ellenberg light indicator values (L-value x -axis). The violin plot shows stomatal density (y -axis, log-scale) as a function of L-value. The width of the grey polygons indicates the density of data. Length of grey polygon indicate the range of the data; the dot indicates the median; the thick lines indicate the 0.25 and 0.75 quantiles. Points outside the polygons are statistical outliers.

585 **Supporting Information**




Figure S1: Most hydrophytes are hyperstomatous, having most stomata on the adaxial ('upper') surface (high SD_{propAd}). The violin plot shows stomatal ratio as a function of Raunkiaer lifeform. The width of the grey polygons indicates the density of data. Length of grey polygon indicate the range of the data; the point indicates the median; the thick lines indicate the 0.25 and 0.75 quantiles. Sample sizes per lifeform in the dataset are given above the upper plot margin. SD_{ad} and SD_{total} stand for the stomatal density on the adaxial surface and the total leaf surface (adaxial plus abaxial), respectively.